

Name: Valteri berlijanto enzo trajkunovic

Student Number: 2434404

Module Code & Title: LCHS4008 introduction to public health

Lecturer's Name: Vivian Osuchukwu

Assignment Title: Dementia: Monitoring, Control, and Community Impact

Introduction, Definition, Signs and Symptoms, and Incidence

Dementia is a progressive neurological condition that affects cognitive functioning, memory, and the ability to carry out everyday activities. It is not a single disease but a syndrome caused by various underlying conditions, most commonly Alzheimer's disease (NHS, 2024). Dementia primarily affects older adults, although it is not a normal part of ageing.

Common signs and symptoms include memory loss, confusion, difficulty with communication, impaired judgement, and changes in mood and behaviour. As the condition progresses, individuals may struggle with basic tasks such as eating, dressing, and recognising familiar people (World Health Organization, 2023). These symptoms can significantly reduce independence and quality of life.

Dementia represents a major global health concern, with over 55 million people currently living with the condition worldwide and nearly 10 million new cases diagnosed each year (World Health Organization, 2023). In the UK, approximately 900,000 individuals are affected, and this number is projected to rise to 1.6 million by 2040 (Alzheimer's Society, 2024). These figures highlight the growing severity and impact of dementia on healthcare systems and society.

Main Body

Approaches to Measure and Monitor Dementia

A variety of approaches are used to measure and monitor dementia, combining clinical assessments, medical imaging, and emerging technologies. One of the most widely used methods is cognitive assessment. Tools such as the Mini-Mental State Examination (MMSE) and the Montreal Cognitive Assessment (MoCA) are commonly used to assess memory, attention, and language skills (NHS, 2024). These tests are practical and cost-effective, allowing early screening in primary care. However, they can be influenced by a patient's educational background and may not always detect early-stage dementia, highlighting a limitation in their reliability.

Neuroimaging techniques also play a significant role in diagnosis and monitoring. Magnetic Resonance Imaging (MRI) and Computed Tomography (CT) scans are used to identify structural changes in the brain, such as shrinkage or damage caused by vascular dementia (World Health Organization, 2023). Positron Emission Tomography (PET) scans can detect abnormal protein deposits associated with Alzheimer's disease. While these methods provide more accurate and objective data, they are costly and may not be accessible in all healthcare settings, particularly in under-resourced areas.

In addition, biomarkers are increasingly used to support diagnosis. These involve analysing blood or cerebrospinal fluid for proteins linked to dementia (Alzheimer's Society, 2024). This approach has the advantage of detecting the disease before symptoms become severe. However, it is still developing and is not yet widely available in routine practice.

Furthermore, digital technologies such as wearable devices and smart home monitoring systems are becoming more common. These tools can track behaviour, movement, and sleep patterns, allowing continuous monitoring (NHS, 2024). Although they improve safety and early intervention, concerns around privacy, cost, and user acceptance may limit their use. Overall, no single method is sufficient, and a combination of approaches is required for accurate monitoring.

Control and Management of Dementia

Although dementia cannot currently be cured, several strategies are used to control symptoms and slow disease progression. Pharmacological treatments are commonly prescribed, including cholinesterase inhibitors such as donepezil and NMDA receptor antagonists like memantine (NHS, 2024). These medications can improve communication between brain cells and temporarily reduce symptoms. However, they do not stop the progression of the disease, meaning their effectiveness is limited.

Non-pharmacological approaches are also essential in managing dementia. Cognitive stimulation therapy, physical activity, and social engagement have been shown to improve mental wellbeing and delay cognitive decline (World Health Organization, 2023). These approaches are particularly beneficial because they have fewer side effects compared to medication. However, their success often depends on patient participation and access to services.

Prevention strategies focus on reducing risk factors such as smoking, obesity, and high blood pressure. Maintaining a healthy lifestyle, including a balanced diet and regular exercise, can lower the risk of developing dementia (Alzheimer's Society, 2024). While these measures are effective in reducing risk, they cannot fully prevent the condition, particularly in individuals with genetic predisposition.

Early diagnosis is another key aspect of control. Identifying dementia at an early stage allows individuals to access treatment, plan for the future, and maintain independence for longer. However, delays in diagnosis are common due to stigma and lack of awareness, which can negatively impact outcomes. Therefore, a combination of medical treatment, lifestyle changes, and early intervention is required to effectively manage dementia.

Consequences of Dementia on Individuals and the Community

Dementia has significant consequences for individuals, families, and the wider community. For individuals, the condition leads to a progressive loss of independence, affecting their ability to perform daily tasks such as cooking, dressing, and managing finances. This often results in emotional distress, anxiety, and depression (NHS, 2024). As the disease progresses, individuals may require full-time care, which can reduce their quality of life.

Families and caregivers are also heavily impacted. Caring for someone with dementia can be physically and emotionally demanding, often leading to stress, burnout, and financial strain. Many caregivers reduce working hours or leave employment altogether, resulting in loss of income (Alzheimer's Society, 2024). This highlights the wider economic impact of the condition.

At a community level, dementia places increasing pressure on healthcare and social care services. The growing number of cases in the UK has led to higher demand for GP services, hospital care, and long-term residential care (World Health Organization, 2023). This creates challenges for resource allocation and funding within the healthcare system.

Additionally, dementia contributes to social isolation and stigma. Individuals may withdraw from social activities due to embarrassment or difficulty communicating, while a lack of public understanding can lead to discrimination. This demonstrates that dementia is not only a medical issue but also a significant social concern that affects communities as a whole.

Health and Social Care Support Services

A wide range of health and social care services are required to support individuals living with dementia. Primary healthcare services, particularly general practitioners (GPs), play a key role in early diagnosis and ongoing monitoring. Community nurses also provide essential home-based care, supporting individuals in maintaining independence (NHS, 2024).

Specialist services such as memory clinics offer comprehensive assessments and diagnosis. These clinics involve multidisciplinary teams, including neurologists, psychiatrists, and psychologists, who develop personalised care plans. While these services are highly effective, access can be limited due to long waiting times.

Social care services are crucial in supporting daily living. Home care services assist with tasks such as personal hygiene, meal preparation, and medication management. For individuals in advanced stages, residential and nursing homes provide 24-hour care. Although these services are essential, they can be costly and may not be accessible to all individuals.

Support for caregivers is equally important. Respite care services provide temporary relief, allowing caregivers to rest and reduce stress. Support groups and counselling services also help individuals cope with the emotional challenges of caring for someone with dementia (Alzheimer's Society, 2024). However, awareness and availability of these services can vary.

Voluntary organisations and charities play a vital role in providing information, helplines, and community support programmes. Initiatives such as dementia-friendly communities aim to improve inclusion and reduce stigma. Additionally, assistive technologies such as medication reminders and GPS tracking devices enhance safety and independence.

Overall, effective support requires a coordinated approach involving healthcare professionals, social services, families, and community organisations. However, challenges such as funding limitations and unequal access must be addressed to ensure all individuals receive adequate care.

Conclusion

In conclusion, dementia is a complex and progressive condition that presents significant challenges for individuals, families, and healthcare systems worldwide. As discussed, a

variety of approaches are used to measure and monitor dementia, including cognitive assessments, neuroimaging, and emerging technologies, each with its own strengths and limitations. While there is currently no cure, a combination of pharmacological treatments, lifestyle interventions, and early diagnosis can help manage symptoms and improve quality of life.

The impact of dementia extends beyond the individual, affecting caregivers, communities, and the wider economy through increased healthcare demands and social care needs. This highlights the importance of accessible and effective support services, including healthcare provision, social care, and community-based support.

Overall, addressing dementia requires a coordinated and holistic approach, with continued investment in research, early intervention, and support systems to ensure individuals affected by the condition can live with dignity and appropriate care.

Reference List

- [Alzheimer's Research UK – Dementia Statistics Hub](#)
Alzheimer's Research UK (2024) *Dementia Statistics Hub*.
- [Alzheimer's Society – Dementia Leading Cause of Death](#)
Alzheimer's Society (2024) *Dementia is the UK's leading cause of death*.
- [Alzheimer's Society – UK Dementia Statistics](#)
Alzheimer's Society (2024) *How many people have dementia in the UK?*
- [UK Government Dementia Data and Intelligence](#)
GOV.UK (2026) *Dementia Data and Intelligence*.
- [NHS Dementia Overview](#)
NHS (2024) *Dementia*.
- [WHO Dementia Fact Sheet](#)
World Health Organization (2023) *Dementia Fact Sheet*.

