



Prifysgol Cymru
Y Drindod Dewi Sant
University of Wales
Trinity Saint David

Name: Swapna Khanam

Student ID: 2510691

Module Code: LCHS4008

Module: Introduction to Public Health

Assignment Title: A Public Health Report on Stroke: Incidence, Prevention, Impact, and Care in the UK

Lecturer: Arif Ahmed

Submitted on: 4/17/2026

Table of Contents

1	Introduction.....	1
2	Main Body.....	1
2.1	Approaches to Measure and Monitor Stroke	1
2.2	Control and Prevention of Stroke.....	2
2.3	Consequences of Stroke on Individuals and Communities	3
2.4	Health and Social Care Support Services	3
3	Conclusion.....	4
4	References	4

1 Introduction

Stroke is a dangerous and potentially fatal medical disorder in which the blood supply to a portion of the brain is cut off, resulting in brain cell damage or death (World Health Organisation, 2023). Strokes are classified into two types: ischemic stroke, which occurs when a blood artery becomes blocked, and haemorrhagic stroke, which occurs when there is bleeding in or around the brain. Common indications and symptoms include sudden weakness or numbness (especially on one side of the body), trouble speaking, confusion, vision problems, dizziness, and severe headache (NHS, 2023). Stroke is a significant health issue worldwide.

One of the biggest global health issues is stroke. Stroke accounts for about 11% of all deaths globally, making it the second most common cause of death, according to the World Health Organisation (2023). Every year, almost 100,000 people in the UK suffer a stroke, which continues to be a major cause of mortality and long-term impairment (Stroke Association, 2022). These figures highlight the substantial toll that stroke has on people, societies, and medical systems.

2 Main Body

2.1 Approaches to Measure and Monitor Stroke

Accurate measuring and monitoring techniques are critical for effective public health stroke management. One of the most common ways is epidemiological surveillance, which entails gathering, analysing, and interpreting data on stroke incidence and prevalence. The Sentinel Stroke National Audit Programme (SSNAP) in the United Kingdom collects comprehensive data on stroke care, including hospital admissions, treatment outcomes, and patient recovery (Stroke Association, 2022). This enables healthcare providers to assess performance and enhance the quality of services. Smith et al. (2021) found that post-stroke monitoring improves the detection of arrhythmia and reduces long-term complications.

Another essential way is to use mortality and morbidity data. Mortality data show the number of people who died as a result of a stroke, whereas morbidity data show how many people suffer from stroke-related disabilities. These figures assist policymakers in understanding the scope of the problem and allocating healthcare resources efficiently (Public Health England, 2018).

Risk factor monitoring is also important. Stroke is highly linked to modifiable risk factors such as hypertension, smoking, obesity, poor diet, and physical inactivity. Data acquired by institutions like NHS Digital and Public Health England allows for the identification of high-risk populations and the development of tailored preventative efforts (NHS Digital, 2021).




Figure 1. Post-Stroke Arrhythmia Monitoring Pathway and Clinical Outcomes (Smith et al., 2021)

Screening and early detection strategies also help to monitor stroke risk. The NHS Health Check programme screens people aged 40 to 74 for cardiovascular risk factors such as blood pressure and cholesterol levels. Early detection of these hazards enables timely intervention and minimises the likelihood of stroke (NHS, 2023).

Overall, these approaches are consistent with public health initiatives focused on measuring, monitoring, and managing illness occurrence in communities.

2.2 Control and Prevention of Stroke

Stroke management includes both primary and secondary preventative methods. Primary prevention aims to reduce risk factors before the disease develops. Lifestyle changes are necessary in this regard. According to Public Health England (2016), individuals should be encouraged to adopt healthy behaviours such as eating a balanced diet, exercising regularly, quitting smoking, and limiting alcohol intake.

Since hypertension is the biggest risk factor for stroke, managing blood pressure with medication and routine monitoring is essential (WHO, 2023). Similarly, anticoagulants and cholesterol-lowering medications like statins are used to stop blood clots from forming.

Campaigns for public health are crucial for prevention as well. Initiatives that encourage physical exercise and discourage sedentary behaviour, for instance, help reduce the incidence of strokes. The significance of lifestyle behaviours emphasised in the lecture materials is reflected in these tactics.

Those who have already had a stroke or are at high risk are the target audience for secondary prevention. To avoid recurrence, this entails long-term medication, lifestyle

modifications, and routine medical follow-up. Rapid response programs, like the FAST (Face, Arms, Speech, Time) initiative, greatly lower death and disability by improving early detection and prompt treatment (NHS, 2023).

Furthermore, public health policies targeting health inequalities are required. People at a disadvantage on the social gradient have less access to healthcare and are more likely to live and work in unhealthy environments, exposing them to risk factors for disease (Marmot, 2010). So, reducing inequalities is vital to preventing disease.

2.3 Consequences of Stroke on Individuals and Communities

Stroke has a significant impact on people, families, and communities. Stroke frequently causes long-term physical impairments in individuals, such as paralysis, trouble speaking, and cognitive decline. An individual's capacity to carry out everyday tasks and preserve independence may be severely diminished by these deficits (Stroke Association, 2022).

Stroke has major psychological implications in addition to physical ones. Due to changes in their lifestyle and physical capabilities, many survivors suffer from mental discomfort, anxiety, and depression. This demonstrates the wider effects of illness on wellbeing and mental health.

There are also clear social repercussions. Financial instability may result from stroke survivors' inability to return to the workforce. Social isolation is a regular occurrence, especially for people with severe disabilities. Family members frequently take on caregiving responsibilities, which can be financially and emotionally taxing.

Stroke has a large impact on community health and social care costs, with the expense of treatment, rehabilitation, and long-term care straining NHS resources. The increasing demand for social care services highlights why it is important that integrated healthcare systems are in place.

Health inequalities exacerbate these problems. People from the lower socio-economic classes have a higher incidence of stroke and a greater impact. There are disparities in life expectancies, access to care, and exposure (Public Health England, 2018).

2.4 Health and Social Care Support Services

Individuals suffering from stroke require a complete variety of health and social care services. Acute care is the initial step of treatment. Rapid identification and intervention in dedicated stroke centres are crucial for increasing survival rates and lowering long-term impairment. When provided promptly, treatments like thrombolysis and thrombectomy can considerably improve outcomes (NHS, 2023).

For recuperation, rehabilitation treatments are crucial. Physiotherapists, occupational therapists, psychologists, and speech-language pathologists collaborate in multidisciplinary teams to help patients restore their cognitive and physical capacities. Depending on how severe the stroke was, rehabilitation could take months or years.

In long-term care, community-based services are essential. These consist of support groups, outpatient clinics, and home care services. Social care services help people with everyday tasks, preserving their freedom and enhancing their quality of life.

Programs for health promotion and education are also crucial. Public health initiatives increase knowledge of stroke prevention techniques and symptoms. For instance, the FAST campaign promotes early detection and prompt medical care, which can save lives.

Addressing health disparities is also a top priority. In order to lessen health disparities, the Marmot Review (2010) highlights the significance of enhancing living circumstances, access to healthcare, and social support. Preventive, early intervention, and community support policies are crucial.

Lastly, caregiver assistance is essential. To effectively handle their responsibilities, family members who frequently offer long-term care need access to financial aid, emotional support, and training.

3 Conclusion

In conclusion, stroke is a major public health problem that impacts the country and the globe. It impacts individuals, families, and healthcare systems; is a leading cause of mortality; and results in long-term disability. Knowledge of the burden of disease and quality of care is greatly dependent on effective measurement and monitoring systems, such as national registries and epidemiological studies.

Preventative measures, including lifestyle change, medical treatment, and public health policies, are effective ways to reduce the occurrence of strokes among people, but do you know that the effects of these diseases are not limited to physical health; in fact, it hit at social interaction, mental wellbeing, and financial stability of a person.

To address the needs of impacted people, comprehensive health and social care services are required, including acute treatment, rehabilitation, and community assistance. In order to guarantee fair access to care and enhance results, it is also crucial to address health disparities. To effectively manage stroke and lessen its effects on society, a coordinated and comprehensive public health approach is necessary.

4 References

Marmot, M. (2010). *Fair society, healthy lives: the Marmot Review*. London: Institute of Health Equity.

NHS (2023) *Stroke*. Available at: <https://www.nhs.uk/conditions/stroke/> (Accessed: 6 April 2026).

NHS Digital (2021) *Statistics on obesity, physical activity and diet*. Available at: <https://digital.nhs.uk/data-and-information/publications/statistical/statistics-on-obesity-physical-activity-and-diet> (Accessed: 6 April 2026).

Public Health England (2016) *Health matters: getting every adult active every day*. London: PHE.

Public Health England (2018) *Health profile for England*. London: PHE.

Stroke Association (2022). *State of the Nation: Stroke statistics*. London: Stroke Association.

Smith, J., Brown, A. and Patel, R. (2021) *Clinical and health economic evaluation of a post-stroke arrhythmia monitoring service*. *BMJ Open*, 11(3), pp. 1–10.

World Health Organisation (2023) *Stroke*. Available at: <https://www.who.int/news-room/fact-sheets/detail/stroke> (Accessed: 6 April 2026).