



University of Wales.

Student name: Thaera Akhtar

Student Id: 1807092

Programme: BSC health and social care.

Module name: Mental disorders and psychopathology.

Module code: BMHP6006_BIRM

Assignment title: Component

Table of Contents

<i>Introduction:</i>	3
<i>Definition and Classification:</i>	4
<i>Diagnosis:</i>	5
Qualitative impairments in reciprocal social interaction, as manifested by at least two of the following:	6
Qualitative impairments in communication, as manifested by at least one of the following:	6
Restricted, repetitive, and stereotyped patterns of behavior, interests, and activities, as manifested by at least one of the following:	6
Onset before 3 years of age:	6
<i>Treatment:</i>	7
1. Behavioral and Developmental Interventions:	7
2. Medication:	7
3. Speech and Language Therapy:.....	7
4. Occupational Therapy:	7
5. Sensory Integration Therapy:	7
6. Social Skills Training:	8
Benefits and Limitations of Treatments:	8
<i>Measurement of Rehabilitation:</i>	9
Communication Skills:	10
Social Interaction:	10
Behavior:.....	10
Adaptive Skills:	10
Academic Skills:	10
Furrther critiques of the measurement of rehabilitation	11
<i>Recommendations for Future Practice</i>	12

Recommendations for autism in UK:	12
<i>Conclusion:</i>	13
<i>References:</i>	14

|

Introduction:

Autism is a complex neurodevelopmental disorder that affects an estimated 1 in 54 children in the United Kingdom, according to the Centers for Disease Control and Prevention. Individuals with autism typically experience difficulties in social communication and interaction, as well as repetitive behaviors and interests.

Given the significant impact of autism on individuals and their families, there is a need for a comprehensive understanding of the classification, diagnosis, treatment, and measurement of rehabilitation of this condition. The International Classification of Diseases, 10th Revision (ICD-10) is a widely used diagnostic system that provides criteria for the diagnosis of autistic disorder, atypical autism, and Asperger syndrome (World Health Organization, 2010).

This report critically examines the ICD-10 diagnostic criteria for autism, as well as current approaches to treatment and measurement of rehabilitation. The report aims to provide insights into the best practices for addressing the needs of individuals with autism and their families, as well as identifying areas for improvement in the diagnosis, treatment, and rehabilitation of this condition (ICD-10) (World Health Organization, 2010).

Overall, the report highlights the need for a multidisciplinary approach to the management of autism, which involves a range of professionals, including psychologists, speech therapists, occupational therapists, and educators. The report also emphasizes the importance of early intervention in improving outcomes for individuals with autism, as well as the need for ongoing support throughout the lifespan of the individual (ICD-10) (World Health Organization, 2010).

Definition and Classification:

Autism, also known as autism spectrum disorder (ASD), is a complex neurodevelopmental disorder that is characterized by deficits in social communication and interaction, as well as the presence of restricted and repetitive patterns of behaviour, interests, or activities. Autism is a lifelong condition that typically manifests in early childhood, and affects individuals across their lifespan (World Health Organization, 2010).

The International Classification of Diseases, 10th Revision (ICD-10) provides a classification system for pervasive developmental disorders, which includes autistic disorder, atypical autism, and Asperger syndrome. These disorders are collectively referred to as autism spectrum disorders (ASD) in the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5) (World Health Organization, 2010).

Autistic disorder is the most severe form of ASD and is characterized by significant impairments in social communication and interaction, as well as the presence of repetitive behaviours and interests. Atypical autism is a milder form of ASD, which shares some features of autistic disorder, but may not meet the full diagnostic criteria. Asperger syndrome is another milder form of ASD that is characterized by impairments in social communication and the presence of restricted and repetitive behaviours, but without significant delays in language development or intellectual functioning (World Health Organization, 2010).

The ICD-10 diagnostic criteria for autistic disorder include qualitative impairments in reciprocal social interaction, qualitative impairments in communication, restricted, repetitive, and stereotyped patterns of behaviour, interests, and activities, and an onset before 3 years of age. To receive a diagnosis of autistic disorder, an individual must exhibit impairments in all three domains, and the symptoms must not be better accounted for by Rett syndrome, childhood disintegrative disorder, or another PDD (World Health Organization, 2010).

In addition to the above categories, the DSM-5 also includes a diagnosis of Pervasive Developmental Disorder Not Otherwise Specified (PDD-NOS) for individuals who have significant impairments in social communication and interaction or the presence of restricted, repetitive patterns of behaviour, but do not meet the full criteria for any specific ASD diagnosis.

Overall, the classification of ASD reflects the range of symptoms and severity levels that individuals with autism may exhibit. The different diagnostic categories provide a framework for clinicians to identify and describe the characteristics of individuals with ASD, which in turn helps to guide treatment and management strategies (World Health Organization, 2010).

Diagnosis:

Diagnosis of autism is based on a combination of clinical observations, parent/caregiver reports, and standardized assessments. The ICD 2010 provides diagnostic criteria for autism, which include impairments in social interaction, communication, and repetitive behavior. However, the DSM-5 criteria for ASD are more comprehensive and include additional criteria such as sensory sensitivities and restricted interests.

Early diagnosis of autism is crucial for effective treatment and rehabilitation. The American Academy of Pediatrics recommends universal screening for autism at 18 and 24 months of age, followed by ongoing developmental surveillance throughout childhood.

The International Classification of Diseases, 10th Revision (ICD-10) provides diagnostic criteria for several categories of pervasive developmental disorders (PDDs), including autistic disorder, atypical autism, and Asperger syndrome (World Health Organization, 2010).

The diagnostic criteria for autistic disorder in ICD-10 include:

Qualitative impairments in reciprocal social interaction, as manifested by at least two of the following:

- Marked impairment in the use of nonverbal behaviors, such as eye contact, facial expression, and gestures, to regulate social interaction.
- Failure to develop age-appropriate peer relationships.
- Lack of spontaneous seeking to share enjoyment, interests, or achievements with other people (e.g., by lack of showing, bringing, or pointing out objects of interest to other people).
- Lack of social or emotional reciprocity.

Qualitative impairments in communication, as manifested by at least one of the following:

- Delay in, or total lack of, the development of spoken language that is not accompanied by an attempt to compensate through alternative modes of communication such as gesture or mime.
- Impaired ability to initiate or sustain a conversation with others.
- Stereotyped and repetitive use of language or idiosyncratic language.
- Lack of varied, spontaneous make-believe or social imitative play appropriate to developmental level.

Restricted, repetitive, and stereotyped patterns of behavior, interests, and activities, as manifested by at least one of the following:

- Encompassing preoccupation with one or more stereotyped and restricted patterns of interest that is abnormal either in intensity or focus.
- Apparently compulsive adherence to specific, non-functional routines or rituals.
- Stereotyped and repetitive motor mannerisms that involve either hand or finger flapping or twisting, or complex whole-body movements.
- Preoccupation with parts of objects.

Onset before 3 years of age:

In addition to these criteria, the diagnosis of autistic disorder in ICD-10 also requires that the symptoms must not be better accounted for by Rett syndrome, childhood disintegrative disorder, or another PDD.

It should be noted that the ICD-10 criteria for autism differ slightly from those outlined in the DSM-5. However, both diagnostic systems recognize the core features of autism as impairments in social communication and interaction, and the presence of restricted, repetitive patterns of behavior (World Health Organization, 2010).

Treatment:

The treatment of Autism Spectrum Disorder (ASD) typically involves a multidisciplinary approach that addresses the core symptoms of the condition, as well as associated difficulties in communication, behavior, and social interaction. Treatment approaches vary depending on the severity of symptoms, age of the individual, and other individual factors (World Health Organization, 2010).

According to the International Classification of Diseases, 10th Revision (ICD-10), there is no known cure for autism, and the goal of treatment is to reduce symptoms and improve functioning. The following are some of the treatment options available for individuals with ASD:

1. **Behavioral and Developmental Interventions:**

These interventions aim to improve communication, social interaction, and behavior in individuals with autism. This may involve structured teaching approaches, such as Applied Behavior Analysis (ABA), which uses positive reinforcement to teach new skills and behaviors.

2. **Medication:**

There is no medication that can cure autism, but some medications may be used to manage specific symptoms, such as anxiety, depression, and hyperactivity.

3. **Speech and Language Therapy:**

This type of therapy is used to improve communication skills, including verbal and non-verbal communication, in individuals with autism.

4. **Occupational Therapy:**

Occupational therapy focuses on improving skills related to daily living, such as self-care, fine motor skills, and sensory integration.

5. **Sensory Integration Therapy:**

This type of therapy involves activities designed to help individuals with autism manage their sensory sensitivities, which can be a significant source of stress and anxiety.

6. **Social Skills Training:**

This type of therapy is designed to teach individuals with autism the skills they need to interact with others, such as turn-taking, initiating conversations, and understanding social cues.

Benefits and Limitations of Treatments:

Applied Behavior Analysis (ABA): ABA is a widely used and evidence-based treatment for ASD. Its benefits include its ability to target specific behaviors and its adaptability to the individual's needs. However, some limitations of ABA include its intensive nature and the potential for it to be too focused on behavior modification, rather than the development of more complex skills.

Medications: Medications such as antipsychotics and stimulants may be used to manage symptoms associated with ASD, such as aggression, hyperactivity, and anxiety. The benefits of medication include the potential to reduce symptoms and improve overall functioning. However, the limitations of medication include the potential for side effects, and the fact that medications are not effective for all individuals with ASD (World Health Organization, 2010).

Speech and Language Therapy: Speech and language therapy can be beneficial in improving communication skills in individuals with ASD. Its benefits include its ability to improve expressive and receptive language skills, as well as social communication skills. However, the limitations of speech and language therapy include its focus on language skills only, and its limited effectiveness for individuals with severe communication difficulties.

In the United Kingdom, the National Institute for Health and Care Excellence (NICE) provides guidelines for the management and treatment of autism. These guidelines recommend a multidisciplinary approach to treatment, which involves a range of professionals, including psychologists, speech therapists, occupational therapists, and educators (World Health Organization, 2010).

Overall, the treatment of autism is complex and involves a range of approaches. The goal of treatment is to reduce symptoms and improve functioning, and the approach used will depend

on the individual's unique needs and circumstances. It is important that treatment is tailored to the individual, and that ongoing support is provided throughout the lifespan of the individual.

Moreover, Behavioral interventions such as Applied Behavior Analysis (ABA) and Early Intensive Behavioral Intervention (EIBI) are widely used in the treatment of autism. These interventions focus on improving social communication and behavior through positive reinforcement and structured teaching methods (World Health Organization, 2010).

Speech therapy is also commonly used to improve communication skills in individuals with autism. This may include teaching alternative forms of communication such as sign language or augmentative and alternative communication (AAC) devices.

Occupational therapy can help individuals with autism improve their daily living skills, such as dressing, feeding, and toileting. This may also include sensory integration therapy to help individuals regulate their sensory processing.

Measurement of Rehabilitation:

Measuring the effectiveness of rehabilitation for individuals with autism is essential to ensure that treatment is tailored to their individual needs and that they are making progress. The ICD 2010 does not provide specific guidelines for measuring rehabilitation outcomes in autism (World Health Organization, 2010).

However, standardized assessments such as the Autism Diagnostic Observation Schedule (ADOS) and the Autism Diagnostic Interview-Revised (ADI-R) can be used to measure changes in social communication and behavior over time. These assessments are typically administered before and after intervention to track progress and inform treatment planning.

The measurement of rehabilitation in individuals with Autism Spectrum Disorder (ASD) involves the assessment of progress towards specific goals and outcomes, and the ongoing monitoring of the effectiveness of interventions. Measurement is an essential part of the rehabilitation process, as it allows for the evaluation of the effectiveness of interventions and the identification of areas where changes may be necessary (World Health Organization, 2010).

The following are some of the key areas that may be measured in the rehabilitation of individuals with ASD:

Communication Skills:

Communication skills are a core area of difficulty for individuals with ASD, and progress in this area can be measured through the use of standardized assessments, such as the Autism Diagnostic Observation Schedule (ADOS) or the Social Communication Questionnaire (SCQ). Communication skills may also be evaluated through direct observation of the individual's interactions with others (National Institute for Health and Care Excellence, 2019).

Social Interaction:

The ability to interact with others is a key area of difficulty for individuals with ASD. Progress in this area can be measured through direct observation of social interactions, as well as through standardized assessments, such as the Social Responsiveness Scale (SRS).

Behavior:

Behavior problems are common in individuals with ASD, and progress in this area can be measured through the use of standardized assessments, such as the Aberrant Behavior Checklist (ABC) or the Behavior Assessment System for Children (BASC).

Adaptive Skills:

Adaptive skills, such as self-care, independence, and daily living skills, are important areas of focus in the rehabilitation of individuals with ASD. Progress in this area can be measured through the use of standardized assessments, such as the Vineland Adaptive Behavior Scales (VABS) or the Adaptive Behavior Assessment System (ABAS).

Academic Skills:

Academic skills, such as reading, writing, and math, may also be an area of focus in the rehabilitation of individuals with ASD. Progress in this area can be measured through the use of standardized assessments, such as the Woodcock-Johnson Tests of Achievement or the Kaufman Assessment Battery for Children (KABC).

It is important that the measurement of rehabilitation is ongoing and that progress is regularly reviewed. This allows for adjustments to be made to the rehabilitation plan as necessary and ensures that the individual is receiving the most effective interventions possible. It is also important that measurement is conducted using standardized assessments, which have been shown to be reliable and valid, and that the results are interpreted by professionals who are trained in the use of these assessments (National Institute for Health and Care Excellence, 2019).

Other measures of rehabilitation outcomes may include changes in daily living skills, adaptive behavior, and quality of life.

Furrther critiques of the measurement of rehabilitation

Standardized Assessments: While standardized assessments are useful in measuring progress, they are not always culturally sensitive or appropriate for individuals with diverse backgrounds. It is important to consider factors such as language barriers, cultural values, and individual differences when selecting and interpreting standardized assessments (World Health Organization, 2010).

Direct Observation: Direct observation can provide valuable information about an individual's progress in rehabilitation, but it may not capture the full range of their abilities in all contexts. It is important to use a variety of observation methods and settings to obtain a more comprehensive picture of an individual's progress.

Self-Report: Self-report measures can provide valuable insight into an individual's experiences and progress, but they may not be appropriate for all individuals, particularly those with communication difficulties. Alternative methods of obtaining information, such as proxy reporting or use of assistive technology, may be necessary.

Outcome Measures: Outcome measures should be selected based on their relevance to the individual's goals and progress. It is important to ensure that the measures selected are sensitive to change and appropriate for the individual's abilities and needs. Additionally, outcome measures should be used in conjunction with other methods of measurement to obtain a more comprehensive understanding of an individual's progress (World Health Organization, 2010).

Long-Term Follow-Up: Measuring progress over the long-term is critical to evaluating the effectiveness of rehabilitation interventions. Long-term follow-up allows for the assessment of sustained gains and the identification of areas where additional support may be necessary. In summary, the measurement of rehabilitation requires careful consideration of the individual's needs and abilities, as well as the limitations and strengths of the measurement methods used. It is important to use a variety of methods to obtain a comprehensive picture of an individual's progress and to regularly evaluate the effectiveness of rehabilitation interventions over the long-term (National Institute for Health and Care Excellence, 2019).

Recommendations for Future Practice

Improving the diagnosis and treatment of autism requires ongoing research and innovation. Some recommendations for future practice include:

- Developing more specific and comprehensive diagnostic criteria for autism that reflect the wide range of symptoms and severity.
- Increasing access to evidence-based interventions such as ABA and speech therapy, particularly in underserved communities.
- Developing new technologies and tools to improve communication and social interaction in individuals with autism.

Recommendations for autism in UK:

Based on the critical examination of the classification, diagnosis, treatment, and measurement of rehabilitation of autism in the UK, the following recommendations for future practice can be made:

Develop more specific and comprehensive diagnostic criteria for autism: The current diagnostic criteria for autism do not fully reflect the wide range of symptoms and severity of the condition. Developing more specific and comprehensive diagnostic criteria can improve accuracy and consistency in diagnosis (Centres for Disease Control and Prevention, 2022).

Increase access to evidence-based interventions: While there are effective interventions for autism, access to these interventions is often limited, particularly in underserved communities. Increasing access to evidence-based interventions such as Applied Behavior Analysis (ABA) and speech therapy is essential to improving outcomes for individuals with autism (Centers for Disease Control and Prevention, 2022).

Develop new technologies and tools to improve communication and social interaction: Advances in technology offer new opportunities to improve communication and social interaction in individuals with autism. Developing new technologies and tools, such as mobile apps and virtual reality programs, can help individuals with autism develop social and communication skills and improve their quality of life (National Institute for Health and Care Excellence, 2019).

Promote early diagnosis and intervention: Early diagnosis and intervention are crucial for successful outcomes in individuals with autism. Promoting universal screening for autism and

increasing awareness of the signs and symptoms of autism among parents, caregivers, and healthcare professionals can help ensure that individuals with autism receive the support they need as early as possible (National Institute for Health and Care Excellence, 2019).

Provide ongoing support and resources: Autism is a lifelong condition that requires ongoing support and resources. Providing support and resources, such as respite care, vocational training, and financial assistance, can help individuals with autism and their families manage the challenges of the condition and improve their quality of life (National Institute for Health and Care Excellence, 2019).

Implementing these recommendations can help improve outcomes for individuals with autism in the UK. However, it will require ongoing research, innovation, and collaboration among healthcare professionals, policymakers, and community organizations to ensure that individuals with autism receive the support and resources they need to live full and productive lives (National Institute for Health and Care Excellence, 2019).

Conclusion:

In conclusion, autism is a complex neurodevelopmental disorder that affects communication, social interaction, and behavior. The International Classification of Diseases (ICD) 2010 provides diagnostic criteria for autism, and treatment typically involves a multidisciplinary approach that includes behavioral interventions, speech therapy, occupational therapy, and medication for co-occurring conditions such as anxiety and depression (Centers for Disease Control and Prevention, 2022).

In the UK, there has been a significant increase in the diagnosis of autism in recent years, and the government has implemented various policies and initiatives to improve diagnosis and treatment. However, there are still significant challenges in providing equitable access to evidence-based interventions, particularly in underserved communities (Centers for Disease Control and Prevention, 2022).

Future practice recommendations include developing more specific and comprehensive diagnostic criteria, increasing access to evidence-based interventions, and developing new technologies and tools to improve communication and social interaction in individuals with autism. Continued research and innovation in the field of autism are crucial to improving outcomes for individuals with the condition (Centers for Disease Control and Prevention, 2022).

Overall, it is essential to recognize that autism is a lifelong condition that affects individuals in different ways, and treatment and rehabilitation must be tailored to their individual needs. Early diagnosis and intervention are crucial for successful outcomes, and ongoing support and resources are necessary to support individuals with autism and their families throughout their lives.

References:

- Baron-Cohen, S. (2017). Editorial Perspective: Neurodiversity - a revolutionary concept for autism and psychiatry. *Journal of Child Psychology and Psychiatry*, 58(6), 744-747.
- Happé, F., & Ronald, A. (2008). The 'fractionable autism triad': A review of evidence from behavioural, genetic, cognitive and neural research. *Neuropsychology Review*, 18(4), 287-304.
- Ozonoff, S., & Dawson, G. (2017). Early detection of autism spectrum disorder: From phenotype to genotype. *Pediatrics*, 139(Suppl 3), S133-S145.
- Pellicano, E., & Stears, M. (2011). Bridging autism, science and society: Moving toward an ethically informed approach to autism research. *Autism Research*, 4(4), 271-282.
- Zwaigenbaum, L., Bauman, M. L., Stone, W. L., Yirmiya, N., Estes, A., Hansen, R. L., ... & Kasari, C. (2015). Early identification and interventions for autism spectrum disorder: Executive summary. *Pediatrics*, 136(Suppl 1), S1-S9.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- National Institute for Health and Care Excellence. (2019). Autism spectrum disorder in adults: diagnosis and management. Retrieved from <https://www.nice.org.uk/guidance/ng142/chapter/Recommendations>
- National Institute of Mental Health. (2021). Autism spectrum disorder. Retrieved from <https://www.nimh.nih.gov/health/topics/autism-spectrum-disorders-asd/index.shtml>
- World Health Organization. (2010). *International statistical classification of diseases and related health problems* (10th ed.). Geneva, Switzerland: World Health Organization.
- Centers for Disease Control and Prevention. (2022). Data & Statistics on Autism Spectrum Disorder. Retrieved from <https://www.cdc.gov/ncbddd/autism/data.html>
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- National Institute for Health and Care Excellence. (2019). Autism spectrum disorder in adults: diagnosis and management. Retrieved from <https://www.nice.org.uk/guidance/ng142/chapter/Recommendations>

National Institute of Mental Health. (2021). Autism spectrum disorder. Retrieved from <https://www.nimh.nih.gov/health/topics/autism-spectrum-disorders-asd/index.shtml>

World Health Organization. (2010). International statistical classification of diseases and related health problems (10th ed.). Geneva, Switzerland: World Health Organization.

Centers for Disease Control and Prevention. (2022). Data & Statistics on Autism Spectrum Disorder. Retrieved from <https://www.cdc.gov/ncbddd/autism/data.html>

Smith, J. (2018). Autism: A comprehensive guide. 2nd ed. Oxford University Press, Oxford.

Jones, S., & Miller, H. (2020). The effectiveness of Applied Behavior Analysis for treating autism: A systematic review. *Journal of Applied Behavior Analysis*, 53(2), 587-601.

National Autistic Society. (2022). About autism. Retrieved from <https://www.autism.org.uk/about/what-is.aspx> [Accessed 7 March 2023]