

BRIEF REPORT

How Do Electronic Cigarettes Affect Cravings to Smoke or Vape? Parsing the Influences of Nicotine and Expectancies Using the Balanced-Placebo Design

Amanda M. Palmer and Thomas H. Brandon

University of South Florida and H. Lee Moffitt Cancer Center & Research Institute, Tampa, Florida

Objective: Although electronic cigarettes (e-cigarettes) are frequently initiated for smoking cessation, results from the first two clinical trials testing this suggest that the perceived benefits of vaping may be influenced by non-nicotine factors, including cognitive outcome expectancies. The current study investigated the separate and combined effects of nicotine delivery and outcome expectancies on cravings for cigarettes and e-cigarettes using a balanced-placebo experiment. **Method:** Drug dosage (contains nicotine or not) was crossed with instructional set (told nicotine or non-nicotine) during ad lib e-cigarette use sessions by 128 current e-cigarette users (52 identifying as current cigarette smokers or “dual users”). It was hypothesized that reduction in craving for both cigarettes and e-cigarettes following e-cigarette administration would be driven primarily by the instructional set manipulation, reflecting the influence of outcome expectancies. **Results:** As hypothesized, among dual users, a main effect of instructional set emerged on reductions in craving to smoke cigarettes, with participants who were told that their e-cigarette contained nicotine reporting greater craving reduction ($p = .046$). With respect to reduced cravings for e-cigarettes, we found an interaction between drug dose and instructional set ($p = .02$) such that nicotine e-cigarettes reduced cravings more than non-nicotine e-cigarettes only among participants told to expect nicotine. **Conclusions:** Findings suggest that cognitive expectancies contribute to the acute effects of e-cigarettes on craving, which may provide guidance for their potential as smoking cessation aids.

What is the public health significance of this article?

This study demonstrated that use of an electronic cigarette may reduce nicotine craving (i.e., desire to smoke or vape) via nonpharmacological routes, including beliefs about nicotine, rather than simply via nicotine delivery itself. This finding has implications for understanding e-cigarette use as well as its potential as a smoking cessation aid.

Keywords: e-cigarette, smoking, expectancies, balanced-placebo, nicotine

Recently, there has been a shift in the landscape of tobacco use with the introduction of novel products such as electronic cigarettes (e-cigarettes), which are portable, battery-powered devices containing a heating element that aerosolizes a liquid solution. Using e-cigarettes is often referred to as *vaping*, and users may

self-identify as *vapers*. It has been estimated that 2.4% of U.S. adults use e-cigarettes regularly, with the vast majority being current or former smokers (Zhu, Zhuang, Wong, Cummins, & Tedeschi, 2017). Vapers report that their primary reason for using e-cigarettes is to quit or reduce smoking of traditional, combustible

Amanda M. Palmer and Thomas H. Brandon, Department of Psychology, University of South Florida, and Department of Health Outcomes and Behavior, H. Lee Moffitt Cancer Center & Research Institute, Tampa, Florida.

Amanda M. Palmer and Thomas H. Brandon acknowledge Alessandra Santa-Ana, Walid Hazem, Alina Hoehn, Ursula Martinez, John Correa, and Lancia Darville for their assistance in conducting this project, as well as Mark Goldman, Joseph Vandello, and Vani Nath Simmons for their consultation.

This research was supported in part by funding from the University of South Florida, the National Institute on Drug Abuse Grant R01 DA037961,

and the Proteomics Core at the H. Lee Moffitt Cancer Center & Research Institute, a comprehensive cancer center designated by the National Cancer Institute (P30 CA076292). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Thomas H. Brandon has had past research support from Pfizer, Inc.

Correspondence concerning this article should be addressed to Thomas H. Brandon, Department of Health Outcomes and Behavior, H. Lee Moffitt Cancer Center & Research Institute, 4115 East Fowler Avenue, Tampa, FL 33617. E-mail: thomas.brandon@moffitt.org

cigarettes (e.g., Siegel, Tanwar, & Wood, 2011). To date, only two double-blind randomized controlled trials (RCTs) of e-cigarettes for smoking cessation have been published. One study (Bullen et al., 2013) randomly assigned transdermal nicotine patches, nicotine e-cigarettes, and placebo (non-nicotine) e-cigarettes to participants interested in quitting smoking. Another trial (Caponnetto et al., 2013) randomized nontreatment seeking participants to receive one of two nicotine doses or placebo e-cigarettes. Although both trials indicated that e-cigarettes were effective in promoting smoking reduction or cessation, no significant differences were found based on nicotine content.

Cravings to use a drug are often considered the final common pathway in theoretical models of drug use motivation (Baker, Morse, & Sherman, 1986). In the case of tobacco smoking, the FDA-approved medications (NRTs, bupropion, and varenicline) have all been shown to reduce cue- and abstinence-induced cravings to smoke in laboratory paradigms (Ferguson & Shiffman, 2009). In addition, research has found that e-cigarette use reduces cravings to smoke and to vape, with mixed evidence regarding the role of nicotine per se (Dawkins, Turner, Hasna, & Soar, 2012; Perkins, Karelitz, & Michael, 2017). There is also some evidence that the sensorimotor aspects of vaping (i.e., its similarity to smoking behavior) contributes to craving reduction (Van Heel, Van Gucht, Vanbrabant, & Baeyens, 2017).

Drug use and addictive behaviors may also be influenced by outcome expectancies, which are learned, cognitive intervening variables. Drug-related expectancies refer to the degree that individuals expect positive and negative outcomes from drug use. Expectancies have been predictive of initiation, maintenance, cessation, and relapse to alcohol, tobacco, and other substances (Brandon, Juliano, & Copeland, 1999; Goldman, 1999). Research on expectancies for e-cigarette use has been limited to survey data (e.g., Harrell, Marquinez, et al., 2015).

Prior studies have indicated that expectancies can sometimes influence immediate drug use behaviors and outcomes to a greater degree than drug dosage itself (Kirsch, 1985)—often referred to as *the placebo effect*. This phenomenon has been studied through simultaneous expectancy and pharmacological manipulation using the balanced-placebo design (BPD), initially to study the effects of alcohol (Hull & Bond, 1986; Marlatt, Demming, & Reid, 1973) and more recently with tobacco and NRT (Dar & Barrett, 2014; Juliano & Brandon, 2002). This paradigm utilizes a 2 × 2 factorial design in which drug type (active or placebo) is crossed with instructional set (told active or placebo). From this, the effects of both the pharmacologic properties of the drug and expectancies about the drug can be independently evaluated as causal influences. In general, the results from these studies demonstrate that active drug delivery appears to have primary influence over physiological or objective domains, whereas drug expectancies may

influence more emotionally salient or subjective domains, such as craving. These effects have been observed in both laboratory studies and more naturalistic field studies (Dar & Barrett, 2014). Thus, the BPD can be used to test the independent and synergistic effects of nicotine delivery and nicotine expectancies upon craving reduction.

The primary goal of the present study was to investigate the effects of nicotine and expectancies on cravings to smoke and cravings to vape. Current e-cigarette users were randomized to use e-cigarettes that contained either nicotine or non-nicotine solutions, and were independently instructed that the e-cigarette contained nicotine or non-nicotine, resulting in four experimental conditions as illustrated in Table 1. It was hypothesized that instructional set should produce differences in craving reduction such that told nicotine would produce greater reduction in craving than told no-nicotine, reflecting the role of nicotine-related expectancies. There were no a priori hypotheses regarding main effects of nicotine content or interactions.

Method

Participants

Participants were 130 individuals recruited primarily from flyers at local vape shops (see Table 2). Participants were screened by telephone for the following eligibility criteria: (1) ≥18 years old; (2) current e-cigarette users (daily nicotine solution use for ≥30 days); (3) history of cigarette smoking (≥100 lifetime cigarettes; ≥1 cigarette/day for ≥30 days); (4) no current e-cigarette cessation attempt; and (5) not currently pregnant, attempting to get pregnant, or nursing.

Experimental Procedures

Eligible participants were asked to abstain from using e-cigarettes and combustible cigarettes for three hours prior to the session. To increase adherence, participants were told that a breath carbon monoxide (CO) reading would be administered. Upon arrival, research staff obtained written informed consent, and then collected the CO sample ($M = 12.63$, $SD = 8.63$ ppm among smokers; $M = 6.00$, $SD = 4.21$ ppm among former smokers). Participants were then randomized to condition and their e-cigarette solution was prepared by a researcher with no participant contact, ensuring a double-blind for the drug manipulation. Randomization used a 4-block pattern with stratification based on sex, cigarette smoking status (current [defined by smoking >1 cigarette per week] or former), and flavor preference (tobacco, menthol, or fruit).

Table 1
Experimental Design

Drug content	Instructional Set	
	E-cigarette contains nicotine	E-cigarette does not contain nicotine
Nicotine	True positive	False negative, anti-placebo
Non-nicotine	False positive, placebo	True negative

Table 2
Participant Demographics ($N = 128$)

Variable	Description	M (SD) or %
Age	(range: 18–76 years)	36.40 (13.79)
Sex	Female	38%
Race	Black/African American	12%
	White/European Origin	83%
	Other	6%
Ethnicity	Hispanic/Latino	16%
Marital status	Single	60%
Sexual orientation	Identify as LGBT+	13%
Education	High school or less	25%
	Some college	30%
	Tech school/associate's	29%
	4-year college degree or beyond	16%
Income	Under \$10,000	22%
	\$10,000–\$29,999	28%
	\$30,000 or more	49%

Note. No significant differences between conditions were found for any of the variables.

Participants completed demographic and baseline measures, followed by the first administration of craving measures. Participants were then provided an e-cigarette with instructions and labeling consistent with the instructional set conditions. They were instructed to take at least 10 puffs over the 10 min session. Following the ad lib session, the craving measures were readministered. (Secondary outcome variables—affect, appetite, reinforcement, and attention—were then collected and will be reported elsewhere).

Apparatus

Participants were provided with an eGo-style 3.6–4.2 V, 1100 mAh battery with a 2.8-Ohm, 510-style clearomizer. This device contained an LCD display showing number of puffs. This e-cigarette style is considered “second generation,” which deliver nicotine more consistently and are preferred among experienced vapers compared with first generation “cig-a-like” styles (e.g., Dawkins, Kimber, Puwanesarasa, & Soar, 2015).

The solution used was a 50% vegetable glycerin (VG), 50% propylene glycol (PG) liquid. Target nicotine content was either 0 mg/ml or 12 mg/ml, with the latter having produced similar plasma nicotine concentrations (as measured in venous blood samples) as traditional cigarettes (Ramôa et al., 2016). Participants were given the choice of tobacco, menthol, or fruit flavors. This solution was a custom-made “research blend” (Avail Vapor, LLC). The solutions were retested using mass spectrometry and liquid chromatography, which verified that the non-nicotine solutions contained 0 mg/ml nicotine. Final concentrations of the nicotine solutions were 11.2 mg/ml (tobacco), 10.3 mg/ml (menthol), and 10.0 mg/ml (fruit).

Assessments

Baseline assessments. Participants completed questionnaires capturing basic demographic information, smoking and vaping history, and device preference. Dependence on e-cigarettes was measured using the Penn State Electronic Cigarette Dependence Index (Foulds et al., 2015; $\alpha = .70$), whereas cigarette dependence

was measured with the Fagerström Test for Nicotine Dependence (Heatherton, Kozlowski, Frecker, & Fagerström, 1991; $\alpha = .56$ for current smokers).

Craving to smoke and craving to use e-cigarettes. A three-item adaptation of the Questionnaire of Smoking Urges-Brief (QSU; Cox, Tiffany, & Christen, 2001; Kozlowski, Pillitteri, Sweeney, Whitfield, & Graham, 1996) was used to assess craving to smoke ($\alpha = .90$ – $.95$) and craving to vape ($\alpha = .91$ – $.94$). The items, assessing “desire,” “craving,” and “wanting,” were rated 0 to 6, yielding scores from 0 to 24.

Nicotine dosing estimate. As a check on the instructional set manipulation, participants completed a brief questionnaire post ad lib session asking them to estimate the nicotine dose of the provided e-cigarette (0 mg/ml, 6 mg/ml, 12 mg/ml, 18 mg/ml, or 24 mg/ml).

Data Analysis Plan

Factorial 2×2 analyses of variances (ANOVAs) and chi-square tests were used to test for baseline differences between factorial conditions (drug, instructional set). T tests were used to test for baseline differences by sex and smoking status. Next, 2 (drug) $\times$ 2 (instructional set) $\times$ 2 (sex) ANOVAs were used to test the main effects and interactions of the drug and instructional set manipulations on pre-to-post changes in craving. Given previous sex differences found on both general nicotine effects (Perkins, 1996) and e-cigarette outcome expectancies (Copp, Collins, Dar, & Barrett, 2015; Piñeiro et al., 2016), sex was included as a third factor.

Results

Preliminary Analyses

Two participants were removed (instructional noncompliance, incorrect randomization) from final analyses ($N = 128$). Drug $\times$ Instructional Set ANOVAs failed to reveal significant differences between conditions on any demographic, baseline, pretest variables, or puff count. No differences were found between sexes on pread lib outcome variables. Current smokers and former smokers were compared on several descriptive variables, as shown in Table 3.

Craving Reduction

Cravings to smoke. Among current smokers ($n = 52$), the hypothesized main effect of instructional set was observed on change in craving to smoke as measured by the QSU, $F(1, 44) = 4.21$, $p = .046$, $\eta^2 = 0.09$. Greater craving reduction was found among those told they received nicotine ($M = 7.92$, $SD = 6.59$) than among those told they did not receive nicotine ($M = 4.25$, $SD = 5.31$; see Table 4 and Figure 1). No main effect of drug was found, and there were no interactions between instruction, drug, or sex.

Craving to vape. Using the full sample, no main effects of drug, instructional set, or sex emerged, but a significant Drug $\times$ Instructional Set interaction was found on craving to vape, $F(1, 120) = 5.56$, $p = .020$, $\eta^2 = 0.04$ (see Figure 2). Post hoc analyses revealed a significant effect of nicotine dose only when participants were told to expect nicotine, $t(62) = -2.75$, $p = .008$, $d =$

Table 3

Smoking and Vaping Characteristics of the Full Sample and Broken Down by Current Smokers (Dual Users) and Former Smokers

Variable	Full sample (<i>N</i> = 128)	Current smokers (<i>N</i> = 52)	Former smokers (<i>N</i> = 76)	<i>p</i>
Mean cigarettes per day on days smoked (<i>SD</i>) ^a	13.33 (11.31)	8.02 (8.57)	16.96 (11.56)	<.001
Mean days cigarettes smoked per week (<i>SD</i>) ^a	6.01 (1.93)	5.01 (2.35)	6.70 (1.15)	<.001
Mean number of daily e-cigarette uses (<i>SD</i>)	36.50 (53.36)	26.66 (42.40)	43.91 (59.60)	<i>ns</i>
% reporting vaping continuously all day (<i>SD</i>)	47%	35%	66%	.001
Personal device used ^b				
Disposable, "cig-a-like"	11%	14%	5%	<i>ns</i>
Refillable tank system	9%	14%	11%	<i>ns</i>
Tank system with modifiable electric components	66%	52%	75%	<i>ns</i>
Mean nicotine content of solution in personal device (mg/ml)	8.80 (7.51)	10.57 (8.07)	7.71 (6.98)	.046
Flavor used most often				
Tobacco	11%	17%	7%	<i>ns</i>
Menthol	21%	21%	21%	<i>ns</i>
Fruit	41%	40%	42%	<i>ns</i>
Other (e.g. custard, dessert, beverages)	23%	12%	30%	.013
Reported e-cigarette initiation to quit smoking	68%	50%	80%	<.001
Flavor requested for ad-lib session				
Tobacco	12%	17%	8%	<i>ns</i>
Menthol	25%	25%	25%	<i>ns</i>
Fruit	63%	58%	67%	<i>ns</i>
Mean EDCI	10.05 (4.66)	8.44 (5.10)	11.15 (4.00)	.001
Mean FTND ^a	4.50 (2.89)	3.69 (3.28)	5.05 (2.45)	.008

Note. *ns* = not significant; no significant differences were found between conditions on these variables. EDCI = Penn State Electronic Cigarette Dependence Inventory; FTND = Fagerström Test for Nicotine Dependence.

^a Among former smokers, cigarette quantity, frequency, and dependence reflect reported levels prior to quitting smoking. ^b Personal device used was coded from participants' self-reported device brand/model. Above *p* values represent significant differences between current and former smokers.

.69. Additional paired-comparisons revealed that the true positive group showed significantly greater reductions in craving to vape than the placebo group, $t(61) = -2.57, p = .013, d = .65$, and the anti-placebo group, $t(62) = -2.75, p = .008, d = .69$, and marginally greater than the true negative group $t(63) = -1.96, p = .05, d = .49$.

Nicotine Dosing Estimate

A 2×2 ANOVA revealed main effects of both instruction, $F(1, 124) = 47.17, p < .001$ and drug content, $F(1, 124) = 15.71, p < .001$ on estimated nicotine content. Told nicotine produced higher nicotine estimates ($M = 10.98, SD = 6.64$ vs. $M = 4.12, SD = 5.24$), as did actual nicotine content ($M = 9.50, SD = 7.56$ vs. $M = 5.50, SD = 5.47$).

To further explore the relationship between perceived nicotine dose and craving relief, correlations between these variables were

examined. Among smokers, higher nicotine dose estimates were associated with greater cigarette craving reduction, $r(50) = .37, p = .007$. Among the full sample, nicotine dose estimate was not associated with e-cigarette craving reduction, $r(126) = .15, ns$.

Discussion

This study aimed to address specific motivational factors involved with e-cigarette use, primarily their immediate ability to ameliorate cravings. This study is, to our knowledge, the first fully crossed BPD conducted to parse the independent and synergistic influences of both nicotine delivery and expectancies on craving-related outcomes of e-cigarette use.

Among dual users, the hypothesized main effect of instructional set was observed upon reduction in cravings to smoke, which suggests that the craving reduction was driven by participants' expectancies about the effects of nicotine rather than the pharma-

Table 4

Manipulation Effects on Craving: Nicotine, Instruction, and Interactions

Variable	Condition means				Marginal means drug content		Marginal means instructional set		<i>F</i> (Nicotine)	<i>F</i> (Instruction)	<i>F</i> (Nicotine × Instruction)
	True positive	Placebo	Anti-placebo	True negative	Nicotine	No nicotine	Told nicotine	Told no nicotine			
QSU-Smoke	7.75	8.08	3.93	4.57	5.69	6.19	7.92 ^a	4.25 ^a	.15	4.21*	.02
Modified QSU-Vape	8.00 ^{a,b}	3.68 ^a	3.84 ^b	4.82	5.92	4.26	5.87	4.34	1.73	1.31	5.56*

Note. QSU = three-item version of Questionnaire of Smoking Urges-Brief. Positive difference scores represent reductions in value from pre- to post-tests. Shared superscripts indicate significant differences in cell means:

^a $p < .05$. ^b $p < .01$.

* $p < .05$.

colological properties of nicotine. The correlation between nicotine dosing estimate and craving reduction provides further support for the role of expectancies. These results are consistent with contemporary models and research indicating that drug delivery alone is insufficient for explaining drug craving and its alleviation (e.g., Baker, Piper, McCarthy, Majeskie, & Fiore, 2004). Findings are also consistent with previous BPDs with cigarettes (Juliano & Brandon, 2002), and with a non-BPD study showing that nicotine instructional set influenced craving reduction following use of nicotine-free e-cigarettes (Copp et al., 2015).

Notably, the main effect of instructional set rather than drug content is consistent with the findings from the two extant clinical trials of e-cigarettes for smoking cessation, which both failed to find statistically superior cessation outcomes from nicotine versus placebo products (Bullen et al., 2013; Caponnetto et al., 2013), and it suggests that any therapeutic benefit of e-cigarettes may derive, at least in part, from users' cognitive expectancies about them.

Whereas the findings on cravings to smoke address the potential of e-cigarettes as a substitute for smoking, the effects upon cravings to vape address the maintenance of e-cigarette use itself. Here no main effects were found, but an interaction was found between the two experimental factors that indicated the greatest craving reduction when participants were accurately told that they were receiving nicotine (i.e., true positive). The contrast with the findings on craving to smoke suggest that nicotine delivered via e-cigarettes may reduce cravings for the same product, but it may not transfer to a different nicotine-delivery product, combustible cigarettes.

The results of this study should be considered within the context of some methodological limitations. Both instructional set and, to a lesser degree, actual nicotine dose were found to have influenced participants' estimates of the dose they had received. The former finding indicates that the instructional set manipulation was successful. The latter is not surprising, yet reflects a common challenge with the BPD (Dar & Barrett, 2014) that must be considered when interpreting BPD results. Although we collected puff count, additional topography or blood nicotine measures would have provided more complete data on nicotine delivery. Several of the measures used in this study were adapted from validated cigarette




Figure 1. Manipulation effects on desire to smoke among current smokers ($n = 52$). Main effect of instructional set significant at $p = .046$. Error bars are standard error of the mean.




Figure 2. Manipulation effects on desire to vape among full sample ($N = 128$). Interaction $p = .020$. Error bars are standard error of the mean.

measures, but have not yet been validated specifically for e-cigarettes. Inclusion of a behavioral outcome might have enhanced the self-report findings. Additionally, it should be noted that the analysis of craving to vape utilized a larger sample, which may have yielded more stable sample statistics with greater power. Finally, the majority of participants used modifiable e-cigarette systems, which may deliver nicotine more effectively than the standard device used in the study.

Emerging evidence suggests that e-cigarettes may be an effective tool for smoking cessation (Zhu et al., 2017). The results of the present study suggest that nicotine delivery may not be necessary for the acute management of cravings to smoke via vaping. Thus, the possibility of further harm-reduction through the elimination or reduction of nicotine content without sacrificing e-cigarettes' potential efficacy for smoking cessation is promising. Public health campaigns and clinicians could endorse alternative expectancies about the benefits of vaping upon cigarette craving, reducing the emphasis on nicotine per se. Moreover, in other domains, placebo medications have retained their efficacy even after their content has been revealed to patients (i.e., open-label placebo; Kaptchuk et al., 2010). Future studies should extend this line of research beyond abstinence-induced craving to those induced by negative affect and conditioned stimuli (i.e., "smoking triggers").

References

- Baker, T. B., Morse, E., & Sherman, J. E. (1986). The motivation to use drugs: A psychobiological analysis of urges. *Nebraska Symposium on Motivation*, 34, 257–323.
- Baker, T. B., Piper, M. E., McCarthy, D. E., Majeskie, M. R., & Fiore, M. C. (2004). Addiction motivation reformulated: An affective processing model of negative reinforcement. *Psychological Review*, 111, 33–51. <http://dx.doi.org/10.1037/0033-295X.111.1.33>
- Brandon, T. H., Juliano, L. M., & Copeland, A. L. (1999). Expectancies for tobacco smoking. In I. Kirsch (Ed.), *How expectancies shape experience* (pp. 263–299). Washington, DC: American Psychological Association. <http://dx.doi.org/10.1037/10332-011>
- Bullen, C., Howe, C., Laugesen, M., McRobbie, H., Parag, V., Williman, J., & Walker, N. (2013). Electronic cigarettes for smoking cessation: A randomised controlled trial. *Lancet*, 382, 1629–1637. [http://dx.doi.org/10.1016/S0140-6736\(13\)61842-5](http://dx.doi.org/10.1016/S0140-6736(13)61842-5)

- Caponnetto, P., Campagna, D., Cibella, F., Morjaria, J. B., Caruso, M., Russo, C., & Polosa, R. (2013). Efficiency and Safety of an eElectronic cigAreTte (ECLAT) as tobacco cigarettes substitute: A prospective 12-month randomized control design study. *PLoS ONE*, 8, e66317. <http://dx.doi.org/10.1371/journal.pone.0066317>
- Copp, S. R., Collins, J. L., Dar, R., & Barrett, S. P. (2015). The effects of nicotine stimulus and response expectancies on male and female smokers' responses to nicotine-free electronic cigarettes. *Addictive Behaviors*, 40, 144–147. <http://dx.doi.org/10.1016/j.addbeh.2014.09.013>
- Cox, L. S., Tiffany, S. T., & Christen, A. G. (2001). Evaluation of the brief questionnaire of smoking urges (QSU-brief) in laboratory and clinical settings. *Nicotine & Tobacco Research*, 3, 7–16. <http://dx.doi.org/10.1080/14622200020032051>
- Dar, R., & Barrett, S. P. (2014). The effects of beliefs regarding drug assignment in experimental and field studies of nicotine delivery devices: A review. *Journal of Psychopharmacology*, 28, 1071–1079. <http://dx.doi.org/10.1177/0269881114548295>
- Dawkins, L., Kimber, C., Puwanesarasa, Y., & Soar, K. (2015). First-versus second-generation electronic cigarettes: Predictors of choice and effects on urge to smoke and withdrawal symptoms. *Addiction*, 110, 669–677. <http://dx.doi.org/10.1111/add.12807>
- Dawkins, L., Turner, J., Hasna, S., & Soar, K. (2012). The electronic-cigarette: Effects on desire to smoke, withdrawal symptoms and cognition. *Addictive Behaviors*, 37, 970–973. <http://dx.doi.org/10.1016/j.addbeh.2012.03.004>
- Ferguson, S. G., & Shiffman, S. (2009). The relevance and treatment of cue-induced cravings in tobacco dependence. *Journal of Substance Abuse Treatment*, 36, 235–243. <http://dx.doi.org/10.1016/j.jsat.2008.06.005>
- Foulds, J., Veldheer, S., Yingst, J., Hrabovsky, S., Wilson, S. J., Nichols, T. T., & Eissenberg, T. (2015). Development of a questionnaire for assessing dependence on electronic cigarettes among a large sample of ex-smoking E-cigarette users. *Nicotine & Tobacco Research*, 17, 186–192. <http://dx.doi.org/10.1093/ntr/ntu204>
- Goldman, M. S. (1999). Expectancy operation: Cognitive–neural models and architectures. In I. Kirsch (Ed.), *How expectancies shape experience* (pp. 263–299). Washington, DC: American Psychological Association.
- Gottlieb, A. M., Killen, J. D., Marlatt, G. A., & Taylor, C. B. (1987). Psychological and pharmacological influences in cigarette smoking withdrawal: Effects of nicotine gum and expectancy on smoking withdrawal symptoms and relapse. *Journal of Consulting and Clinical Psychology*, 55, 606–608. <http://dx.doi.org/10.1037/0022-006X.55.4.606>
- Harrell, P. T., Marquinez, N. S., Correa, J. B., Meltzer, L. R., Unrod, M., Sutton, S. K., . . . Brandon, T. H. (2015). Expectancies for cigarettes, e-cigarettes, and nicotine replacement therapies among e-cigarette users (aka vapers). *Nicotine & Tobacco Research*, 17, 193–200. <http://dx.doi.org/10.1093/ntr/ntu149>
- Heatherington, T. F., Kozlowski, L. T., Frecker, R. C., & Fagerström, K.-O. (1991). The Fagerström Test for Nicotine Dependence: A revision of the Fagerström Tolerance Questionnaire. *British Journal of Addiction*, 86, 1119–1127. <http://dx.doi.org/10.1111/j.1360-0443.1991.tb01879.x>
- Hull, J. G., & Bond, C. F., Jr. (1986). Social and behavioral consequences of alcohol consumption and expectancy: A meta-analysis. *Psychological Bulletin*, 99, 347–360. <http://dx.doi.org/10.1037/0033-2909.99.3.347>
- Juliano, L. M., & Brandon, T. H. (2002). Effects of nicotine dose, instructional set, and outcome expectancies on the subjective effects of smoking in the presence of a stressor. *Journal of Abnormal Psychology*, 111, 88–97. <http://dx.doi.org/10.1037/0021-843X.111.1.88>
- Kapchuk, T. J., Friedlander, E., Kelley, J. M., Sanchez, M. N., Kokkotou, E., Singer, J. P., . . . Lembo, A. J. (2010). Placebos without deception: A randomized controlled trial in irritable bowel syndrome. *PLoS ONE*, 5, e15591. <http://dx.doi.org/10.1371/journal.pone.0015591>
- Kirsch, I. (1985). Response expectancy as a determinant of experience and behavior. *American Psychologist*, 40, 1189–1202. <http://dx.doi.org/10.1037/0003-066X.40.11.1189>
- Kozlowski, L. T., Pillitteri, J. L., Sweeney, C. T., Whitfield, K. E., & Graham, J. W. (1996). Asking questions about urges or cravings for cigarettes. *Psychology of Addictive Behaviors*, 10, 248–260. <http://dx.doi.org/10.1037/0893-164X.10.4.248>
- Marlatt, G. A., Demming, B., & Reid, J. B. (1973). Loss of control drinking in alcoholics: An experimental analogue. *Journal of Abnormal Psychology*, 81, 233–241. <http://dx.doi.org/10.1037/h0034532>
- Perkins, K. A. (1996). Sex differences in nicotine versus nonnicotine reinforcement as determinants of tobacco smoking. *Experimental and Clinical Psychopharmacology*, 4, 166–177. <http://dx.doi.org/10.1037/1064-1297.4.2.166>
- Perkins, K. A., Karelitz, J. L., & Michael, V. C. (2017). Effects of nicotine versus placebo e-cigarette use on symptom relief during initial tobacco abstinence. *Experimental and Clinical Psychopharmacology*, 25, 249–254. <http://dx.doi.org/10.1037/pha0000134>
- Piñeiro, B., Correa, J. B., Simmons, V. N., Harrell, P. T., Menzie, N. S., Unrod, M., . . . Brandon, T. H. (2016). Gender differences in use and expectancies of e-cigarettes: Online survey results. *Addictive Behaviors*, 52, 91–97. <http://dx.doi.org/10.1016/j.addbeh.2015.09.006>
- Ramôa, C. P., Hiler, M. M., Spindle, T. R., Lopez, A. A., Karaoghlanian, N., Lipato, T., . . . Eissenberg, T. (2016). Electronic cigarette nicotine delivery can exceed that of combustible cigarettes: A preliminary report. *Tobacco Control*, 25, e6–e9. <http://dx.doi.org/10.1136/tobaccocontrol-2015-052447>
- Siegel, M. B., Tanwar, K. L., & Wood, K. S. (2011). Electronic cigarettes as a smoking-cessation: Tool results from an online survey. *American Journal of Preventive Medicine*, 40, 472–475. <http://dx.doi.org/10.1016/j.amepre.2010.12.006>
- Van Heel, M., Van Gucht, D., Vanbrabant, K., & Baeyens, F. (2017). The importance of conditioned stimuli in cigarette and e-cigarette craving reduction by e-cigarettes. *International Journal of Environmental Research and Public Health*, 14, 193–211. <http://dx.doi.org/10.3390/ijerph14020193>
- Zhu, S. H., Zhuang, Y. L., Wong, S., Cummins, S. E., & Tedeschi, G. J. (2017). E-cigarette use and associated changes in population smoking cessation: Evidence from U.S. current population surveys. *British Medical Journal*, 358, j3262. <http://dx.doi.org/10.1136/bmj.j3262>

Received September 15, 2017

Revision received February 18, 2018

Accepted February 26, 2018 ■