



Prifysgol Cymru
Y Drindod Dewi Sant
University of Wales
Trinity Saint David

Module:
Social Epidemiology and Public Health Development

Module Code:
LCHS7000

TOPIC

"Critical Analysis of Ethnic Health Inequalities: A Social Epidemiological Examination of South Asian Communities in the United Kingdom"

Programme:
MSc Public Health and Social Care in Practice

Lecturer
Dr Ruzanna Gevorgyan

Student Name: MIJA CHAKKALAPARAMBIL MICHAEL

Student ID :2432640

Submission Date:
19 December 2025

Word Count:
1000+1500 words (within the $\pm 10\%$ allowance of the 2500-word limit, excluding contents and references).

Acknowledgement

I confirm that I have not used any AI tools in the research and creation of this assignment. I confirm that I have not presented any AI-generated materials as my work. I confirm I have copies of my drafts, notes, and other resources I used, which I may be asked to provide in evidence.

Table of Contents

TASK 1 – Critical Analysis of Ethnic Health Inequalities

.....	4
Title: <i>A Social Epidemiological Examination of South Asian Communities in the UK</i>	4
1. Introduction: Framing the Crisis	4
1.1 Defining the Crisis: South Asian Health Inequalities as Structural Indicators	4
1.2 Beyond Statistics: From Descriptive Disparities to Critical Analysis	4
2. Contextual Background: The Social Determinants Matrix	4
2.1 Economic Determinants: Poverty Concentration as Structural Violence	4
2.2 Cultural and Linguistic Barriers: Epistemic Exclusion in Healthcare	5
2.3 Environmental Injustice: The Geography of Health Disadvantage	5
2.4 Healthcare System Barriers: Institutional Racism in Clinical Encounters	6
3. Critical Analysis of Key Health Inequalities	6
3.1 Type 2 Diabetes: Social Activation of Biological Susceptibility	6
3.2 Cardiovascular Disease: Syndemic Interaction of Risk Factors	7
3.3 Mental Health: Diagnostic Violence in Cross-Cultural Psychiatry	7
3.4 Healthcare Access: Linguistic Exclusion and Clinical Risk	8
4. National Responses: The Implementation Gap	9
4.1 Policy–Practice Disconnect	9
4.2 Strategic Limitations	9
5. Organisations Supporting South Asian Communities	10
5.1 Community-Led Innovation	10
5.2 Advocacy and Policy Influence	10
6. Conclusion: Towards Transformative Equity	10
6.1 Structural Roots of Inequality	10
6.2 Transformative Recommendations	10

TASK 2 – Epidemiological Analysis

.....	11
Title: <i>Type 2 Diabetes in South Asian Adults, Tower Hamlets</i>	11
1. Introduction: Diabetes as a Structural Indicator	11
1.1 The Syndemic Perspective: Diabetes as Structural Injustice	11
1.2 Beyond Biomedical Paradigms: Epidemiological Transition	11
1.3 Tower Hamlets as a Microcosm of Concentrated Disadvantage	12
2. Global, National, and Local Prevalence: The Epidemiology of Inequality	12
2.1 Global Burden: Nutrition Transition as a Colonial Legacy	12
2.2 National Disparities: Ethnic Patterning and Structural Determinants	12
2.3 Local Concentration: Spatial Inequality in Diabetes Burden	13
3. Determinants and Risk Factors: Beyond Biomedical Models	13
3.1 Biological Determinants: Genetic Susceptibility in Social Context	13
3.2 Social Determinants Matrix: Beyond Individual Behaviour Modification	13
3.3 Environmental Pressures: Obesogenic Environments as Policy Failure	14
4. Public Health Measures – The 3 Ps: Critical Evaluation	14
4.1 Prevention Gaps: Cultural Barriers in Primary Prevention	14

4.2 Protection Limitations: Late Diagnosis as System Failure	15
4.3 Promotion Challenges: Individual Focus vs. Structural Change	15
5. Epidemiological Studies & Health Needs Assessment: Methodological Critique	15
5.1 Methodological Biases: Exclusion of Non-English Speakers in Research	15
5.2 Data Limitations: Incomplete Ethnicity Recording Masking Inequalities	16
5.3 Needs Assessment Critique: Professional vs. Community Perspectives	16
6. Surveillance & Monitoring Systems: Equity Analysis	16
6.1 Data Completeness Paradox: Inverse Relationship with Need	16
6.2 System Limitations: Capturing Disease but Missing Determinants	17
6.3 Community Surveillance: Alternative Models for Equity	17
7. Public Health Impact: Beyond Clinical Metrics	17
7.1 Health Burden Quantification: Beyond Mortality to Quality of Life	17
7.2 Economic Costs: Inequality as a Financial Drain on Systems	18
7.3 Social Consequences: Stigma and Disability as Secondary Burdens	18
8. Intervention Programme & Evaluation: Equity by Design	18
8.1 Theory of Change: From Individual to Structural Intervention	18
8.2 Core Components: Cultural Adaptation as Non-Negotiable	19
8.3 Equity-Focused Evaluation: Measuring Gap Reduction, Not Averages	19
9. Strengths and Weaknesses of the Epidemiological Approach	19
9.1 Methodological Strengths: Quantification for Resource Allocation	19
9.2 Epistemological Limitations: Reductionism and Categorical Violence	20
9.3 Practical Constraints: Implementation Barriers in Real-World Settings	20
10. Conclusion: From Epidemiological Understanding to Structural Transformation	20
References	21

TASK 1: Critical Analysis of Ethnic Health Inequalities: A Social Epidemiological Examination of South Asian Communities in the United Kingdom"

1. Introduction: Framing the Crisis

1.1 Defining the Crisis: South Asian Health Inequalities as Structural Indicators

Health inequalities are preventable, systematic, and unreasonable disparities in health outcomes between population groups that are intergenerational and spatially patterned (Marmot, 2020). In the United Kingdom, these inequalities are particularly pronounced along ethnic lines. South Asian populations—including people of Indian, Pakistani, Bangladeshi, and Sri Lankan descent—experience some of the most persistent and severe health disadvantages.

South Asian groups constitute approximately 8.3% of the UK population (Office for National Statistics, 2021) and represent one of the fastest-growing demographic groups. Despite this, they carry a disproportionately high burden of chronic disease, experience higher levels of premature mortality, and face persistent exclusion within healthcare systems. These patterns are not statistical anomalies but reflect long-standing structural inequalities embedded within social, economic, and healthcare systems. Historical legacies, socioeconomic policy decisions, and institutional practices have systematically disadvantaged these populations across the life course, signalling structural dysfunction rather than individual failure.

1.2 Beyond Statistics: From Descriptive Disparities to Critical Analysis

This report moves beyond descriptive epidemiology, which merely documents disparities, to adopt a critical social epidemiological approach that interrogates why inequalities persist despite decades of policy attention and the presence of a universal healthcare system. Drawing on intersectional and structural determinant frameworks (Crenshaw, 1989; Krieger, 2012), the analysis explores how ethnicity intersects with socioeconomic position, migration history, and gender to produce embodied disadvantage.

Focusing on four major health domains—Type 2 diabetes, cardiovascular disease, mental health, and healthcare access—the report integrates quantitative and qualitative perspectives.

It demonstrates how apparently neutral healthcare systems reproduce inequality through cultural, linguistic, and institutional exclusion. Addressing ethnic health inequalities is therefore not solely a moral or social justice imperative but also a strategic investment in the future health and economic sustainability of the nation.




Figure 1. Conceptual Framework of Health Inequalities and Determinants.

Figure 1: Critical determinants pathway showing structural factors become biologically embedded. This model challenges individual-behavioral explanations by foregrounding systemic factors (Krieger, 2012).

2. Contextual Background: The Social Determinants Matrix

2.1 Economic Determinants: Poverty Concentration as Structural Violence

South Asian communities experience disproportionately high levels of poverty. Approximately 50% of Bangladeshi households and 46% of Pakistani households live in poverty, compared with 20% of White British households (Joseph Rowntree Foundation, 2023). This concentration of economic disadvantage exemplifies what Farmer (2005) describes as structural violence—social arrangements that systematically place particular populations at risk through unequal distribution of resources.

2.2 Cultural and Linguistic Barriers: Epistemic Exclusion in Healthcare

Language barriers significantly contribute to healthcare exclusion. Around 22% of Bangladeshi adults and 18% of Pakistani adults report low English proficiency (Parveen et al., 2018). Trivedi (2022) conceptualises this as epistemic exclusion, whereby non-English health knowledge and communication practices are marginalised within biomedical systems, undermining informed consent, patient safety, and quality of care.

2.3 Environmental Injustice: The Geography of Health Disadvantage

Environmental inequities further compound health risks. Approximately 31% of Bangladeshi households experience overcrowding compared with only 2% of White British households. Ethnically diverse neighbourhoods are also characterised by limited green space and a high density of fast-food outlets (Public Health England, 2020). These obesogenic environments, as described by Swinburn et al. (2011), promote unhealthy default behaviours and increase chronic disease risk.

2.4 Healthcare System Barriers: Institutional Racism in Clinical Encounters

Healthcare system barriers include reports of discrimination by 32% of South Asian patients, limited cultural competence in service design, and inadequate interpreter provision (Salway et al., 2016). These patterns reflect institutional racism—policies and practices that systematically disadvantage ethnic minorities, often without explicit intent (Nazroo, 2022).

3. Critical Analysis of Key Health Inequalities

3.1 Type 2 Diabetes: Social Activation of Biological Susceptibility

South Asian adults are two to four times more likely to develop Type 2 diabetes than their White British counterparts, with onset occurring 5–10 years earlier (Diabetes UK, 2023). While genetic susceptibility, such as TCF7L2 gene variants, exists, attributing inequality primarily to genetics risks medicalising structural disadvantage. Rapid epidemiological shifts within a single generation of migration demonstrate the role of environmental triggers, supporting the concept of the environmentally induced “thrifty genotype” (Hales & Barker, 1992).

3.2 Cardiovascular Disease: Syndemic Interaction of Risk Factors

South Asians are approximately 50% more likely to die from cardiovascular disease. This reflects a syndemic interaction between diabetes, central obesity, hypertension, and chronic socioeconomic stress (Singer, 2009). Gendered norms further exacerbate risk, with South Asian women experiencing diagnostic delays of up to 18 months due to caregiving expectations and cultural beliefs (Zaman et al., 2017).

3.3 Mental Health: Diagnostic Violence in Cross-Cultural Psychiatry

Mental health inequalities remain largely hidden. South Asian populations report significantly higher psychological distress yet are substantially less likely to access talking therapies (Mental Health Foundation, 2020). This reflects Kleinman’s (1978) concept of category fallacy and what Bracken et al. (2012) describe as diagnostic violence, where Western diagnostic frameworks fail to capture culturally specific expressions of distress.

3.4 Healthcare Access: Linguistic Exclusion and Clinical Risk

Reliance on family interpreters due to inadequate professional language support compromises confidentiality and clinical accuracy (Greenhalgh et al., 2015), falling below General Medical Council (2020) standards. Time-limited consultations and culturally inappropriate screening further reinforce structural barriers to equitable care.

and screening programmes which would not be culturally accommodated.




Figure 2: Comparative health and social indicators demonstrating multidimensional inequality (Diabetes UK, 2023; British Heart Foundation, 2024; Mental Health Foundation, 2020). Overcrowding shows the most extreme disparity.

4. National Responses: The Implementation Gap

4.1 Policy–Practice Disconnect

While the Equality Act 2010 and NHS CORE20PLUS5 framework acknowledge inequality, they suffer from a significant implementation gap (Bajpai, 2021). The Public Sector Equality Duty often functions as bureaucratic compliance rather than a transformative mechanism (Salway et al., 2016).

4.2 Strategic Limitations

CORE20PLUS5 lacks binding targets and ring-fenced funding, allowing equity initiatives to be deprioritised during financial constraints (NHS England, 2021). Consequently, only around

15% of eligible South Asian individuals access the NHS Diabetes Prevention Programme (Hanif et al., 2019).

5. Organisations Supporting South Asian Communities

5.1 Community-Led Innovation

The South Asian Health Foundation delivers culturally adapted diabetes education in eight languages through mosques and community centres, exemplifying cultural brokerage in health promotion (Khunti et al., 2019).

5.2 Advocacy and Policy Influence

The Race Equality Foundation promotes co-production models that position community members as equal partners in service design, influencing NHS guidance on inclusive communication (King's Fund, 2023).

6. Conclusion: Towards Transformative Equity

6.1 Structural Roots of Inequality

South Asian health inequalities arise from structural conditions rather than individual behaviours or genetic destiny. Over-reliance on biomedical approaches risks structural iatrogenesis—where healthcare systems inadvertently perpetuate harm by individualising social problems (Metzl & Hansen, 2014).

6.2 Transformative Recommendations

1. **Financial Accountability:** Allocate 5% of Integrated Care System funding to measurable reductions in ethnic health inequalities, supported by community oversight (Bambra et al., 2020).
2. **Cultural Safety Accreditation:** Mandate cultural safety training, with minority patient experience contributing 30% of quality ratings (Ramsden, 2002).

3. **Community-Led Surveillance:** Fund participatory action research enabling communities to collect and interpret their own health data (Smith, 2021).

TASK 2: Epidemiological Analysis of Type 2 Diabetes in South Asian Adults, Tower Hamlets

1. Introduction: Diabetes as a Structural Indicator

1.1 The Syndemic Perspective: Diabetes as Structural Injustice

Tower Hamlets faces a pronounced burden of Type 2 Diabetes (T2D), with a prevalence of 9.5% compared to the national average of 6.9% (NHS Digital, 2023). This disparity does not represent an isolated biomedical phenomenon but instead signals deeply embedded structural inequities. The borough's demographic profile—where approximately 34% of residents are of Bangladeshi heritage—and its ranking as the third most deprived local authority in England illustrate what Singer (2009) conceptualises as a *syndemic*. In this context, T2D interacts with socioeconomic deprivation and systemic discrimination, collectively amplifying disease burden and health inequities.

1.2 Beyond Biomedical Paradigms: Epidemiological Transition and Fundamental Causes

Historically, South Asian populations in both their countries of origin and the UK experienced a disproportionate burden of communicable diseases such as tuberculosis and hepatitis, largely driven by poverty, overcrowded housing, and limited access to early healthcare. However, rapid epidemiological transition within a single generation of migration has shifted disease patterns towards non-communicable diseases (NCDs), including diabetes, cardiovascular disease, and mental health conditions. This transition has been accelerated by globalised nutrition changes, urbanisation, and the westernisation of lifestyles (Popkin, 2006).

Type 2 diabetes should therefore be understood not merely as a biomedical outcome but as an expression of what Link and Phelan (1995) describe as *fundamental causes*—structural conditions such as inequality and racism that systematically shape access to health-protective resources, regardless of the specific disease manifestation.

1.3 Tower Hamlets as a Microcosm of Concentrated Disadvantage

This epidemiological analysis focuses on South Asian adults aged 40 years and above living in Tower Hamlets. It moves beyond conventional biomedical and behavioural explanations by

examining how risk becomes biologically embedded through structural determinants, including economic precarity, environmental injustice, linguistic exclusion, and institutional racism. Positioning T2D within the broader communicable–non-communicable disease transition highlights the necessity for public health responses that address underlying social and political–economic drivers, rather than focusing solely on biological risk factors.

2. Global, National, and Local Prevalence: The Epidemiology of Inequality

2.1 Global Burden: Nutrition Transition as a Colonial Legacy

Globally, an estimated 537 million adults (10.5%) are living with diabetes, a figure projected to rise to 783 million by 2045 (International Diabetes Federation, 2021). South Asia has experienced the most rapid increase, with prevalence in countries such as Bangladesh reaching 14%. This accelerated transition, condensed into a single generation, reflects the global nutrition transition towards highly processed, energy-dense Western diets (Popkin, 2006), shaped by historical and economic forces.

2.2 National Disparities: Ethnic Patterning and Structural Determinants

In the UK, diabetes prevalence demonstrates stark ethnic patterning. South Asian populations experience a prevalence of approximately 9.5%, representing a two- to four-fold increased risk compared with White British populations (4.3%), while prevalence among Black Caribbean groups is estimated at 7.4% (Diabetes UK, 2023). These disparities persist even after adjustment for body mass index (BMI) and socioeconomic status, challenging individual-behavioural explanations and underscoring the role of multilevel structural determinants.

2.2 National Disparities: Ethnic Patterning Revealing Structural Determinants

The ethnic distribution in the prevalence of diabetes in the UK is harsh: the South Asians are 9.5% (2-4 times higher risk), the White British 4.3, the Black Caribbean 7.4% (Diabetes UK, 2023). The gradient remains even after adjusting it with BMI and socioeconomic variables, undermining the explanations of individual-behavioral aspects and moving to the structural

determinants which act at various levels.

Diabetes Prevalence by Ethnicity in UK (2023)




Figure 3: Ethnic disparities in diabetes prevalence reveal structural rather than biological determinants. South Asian risk remains elevated even after controlling for BMI and socioeconomic factors (Mathur et al., 2020).

2.3 Local Concentration: Spatial Inequality in Diabetes Burden

Tower Hamlets has concentrated burden: 9.5% total prevalence, 14.2% amongst South Asians with onset at mean of 45 years (as compared with 55 in the rest of the country). Retinopathy and CKD have complication rates 40 and 35 higher, respectively, proving what Dorling (2015) refers to as the spatial injustice of health distribution, with place becoming the health outcome fate.

3. Determinants and Risk Factors: Beyond Biomedical Models

3.1 Biological Determinants: Genetic Susceptibility in Social Context

Variants of TCF7L2 gene are found to be 1.5x more likely to cause diabetes in South Asians (Sanghera et al., 2018), and metabolic phenotypes are more likely to have visceral adiposity at lower BMI levels. Nevertheless, these biological influences demand social action such as Krieger (2005) defines as embodiment of social situation using biological mechanisms such as chronic stress and inflammation.

3.2 Social Determinants Matrix: Beyond Individual Behavior Modification

Tower Hamlets exemplifies concentrated disadvantage across four domains:

1. **Economic:** It has child poverty rates of 57 percent, unemployment rates of 40 percent in certain wards (Tower Hamlets Council, 2022).
2. **Environmental:** 25% overcrowding (3 x national average), 110 fast food outlets/km².
3. **Nutritional:** A fifth (23) of families go without meals a month because of the price.
4. **Psychosocial:** Discrimination and economic precarity stress, which is chronic.

3.3 Environmental Pressures: Obesogenic Environments as Policy Failure

The built environment is a failure of policy: less green space (14m² per person compared to 32m² in London), greater air pollution than the WHO recommends, and what Swinburn et al. (2011) characterise as food swamps; the overproliferation of fast food which makes default unhealthy choices and overwhelms the power of an individual.

4. Public Health Measures — The 3 Ps: Critical Evaluation

4.1 Prevention Gaps: Cultural Barriers in Primary Prevention

Only 15 percent of the eligible South Asians are covered by the NHS Diabetes Prevention Programme because of language barriers and cultural irrelevance (Hanif et al., 2019). This is what Greenhalgh et al. (2015) define as implementation failure, namely the failure of evidence-based programmes to be effective in various contexts, in which they are applied and the underlying assumption is that it is universal.

4.2 Protection Limitations: Late Diagnosis as System Failure

It shows that South Asians of HbA1c 75 mmol/mol vs 60 mmol/mol average on diagnosis (underlying system delays in detection). Although screening practices are advised by NICE (2022) at BMI ≥ 23 in South Asians, the application has low adherence, which represents guideline-practice divergences with biased representation of minority populations.

4.3 Promotion Challenges: Individual Focus vs. Structural Change

Existing health promotion presupposes the individual agency regardless of structural limitations. This is what Frohlich and Potvin (2008) refer to as the risk factor approach- individual behavioral factors are considered and the contextual risks in the environment that influence the behavioral factors are ignored.




Figure 4: Discrepancy between current effectiveness (bars) and potential of structural approaches (line). Structural interventions show largest implementation gap (Bajpai, 2021).

5. Epidemiological Studies & Health Needs Assessment: Methodological Critique

5.1 Methodological Biases: Exclusion of Non-English Speakers in Research

The UK Biobank is also not representative of non-English speakers (15% vs 40% of Tower Hamlets population), which causes sampling bias to generalize (Sudlow et al., 2015). Randomized trials usually tend to avoid subjects with language barriers, and this leads to what

Green and Glasgow (2006) refer to as the efficacy-effectiveness gap, or what works in the trial does not work in the diverse population.

5.2 Data Limitations: Incomplete Ethnicity Recording Masking Inequalities

The completeness of ethnicity data is at 75 percent in primary care and 50 percent in secondary care (Mathur et al., 2020), which we refer to as the inverse law of data: the patients with the highest health requirements are the very ones with the worst data completeness, causing a concealment of inequalities to the monitoring system and resource allocation based on incomplete data..

5.3 Needs Assessment Critique: Professional vs. Community Perspectives

The standardized tools commonly used in traditional health needs assessment tend to overvalue professional knowledge over that of communities, and therefore they could fail to capture culturally-related issues. The participatory methods, propounded by Israel et al. (1998), place communities as professionals in their own health- alters the locus of determination of need and mode of solution development.

6. Surveillance & Monitoring Systems: Equity Analysis

6.1 Data Completeness Paradox: Inverse Relationship with Need

There are opposite relationships between data completeness and prevention relevance in the surveillance systems. Though hospital admissions are 95 percent complete, it picks up disease at the end-stage. The completeness of community screening data, which is most useful in the context of prevention, is only 25% in high-need areas (Lewis et al., 2025).

6.2 System Limitations: Capturing Disease but Missing Determinants

The existing systems are efficient in tracking the prevalence rates of diabetes and its complications, yet they do not include social determinants or lived experience. This is what Smith (2021) calls epistemological limitation the measurement defines what gets addressed, which results in the creation of cycles in which biomedical elements are the only ones addressed.

6.3 Community Surveillance: Alternative Models for Equity

Alternative models to the traditional power dynamics in knowledge production give community-led surveillance, including participatory health surveys and community health reports, where the communities are in control of data collection and its interpretation, and it is relevant to the local contexts.

7. Public Health Impact: Beyond Clinical Metrics

7.1 Health Burden Quantification: Beyond Mortality to Quality of Life

South Asian men whose diabetes is diagnosed lose 8.2 years compared to 5.3 years in the White British, and the quality-adjusted life years (QALYs) lost is 12.3 per case in Tower Hamlets against 8.7 nationally (Office for National Statistics, 2021). This shows unequal burden out of sheer prevalence, in terms of quantity as well as quality of life.

7.2 Economic Costs: Inequality as Financial Drain on Systems

The direct NHS costs amount to 1,800 per person annually (50% above average), and the indirect productivity costs of previous disability onset (Hex et al., 2012). The economic case of equity is supported by the finding that every 1% of the deprivation is associated with 3% of the increase in the cost of diabetes- reflecting the cost-saving, rather than the moral obligation agenda.

7.3 Social Consequences: Stigma and Disability as Secondary Burdens

The South Asian communities believe that diabetes is culturally-stigmatized and the comorbidities such as retinopathy and neuropathy result in disability burden that disproportionately impacts employment and social involvement of the already disadvantaged groups creating cycles of exclusion (Bhui et al., 2015).

8. Intervention Programme & Evaluation: Equity by Design

8.1 Theory of Change: From Individual to Structural Intervention

The "Tower Hamlets Diabetes Equity Initiative" is based on a theory of change that acknowledges that culturally-adapted prevention and structural interventions within the food environment can lead to 25% reduction in the incidence rates of diabetes among South Asian adults, 5 years later, by intervening at the individual levels and on the environmental factors, simultaneously.

8.2 Core Components: Cultural Adaptation as Non-Negotiable

1. **Community Health Advocates:** Home-based HbA1c testing and culturally-appropriate education on 50 trained residents.
2. **Food Environment Reform:** The prohibition of opening new fast food restaurants through zoning and tax subsidies on traditional food.
3. **Clinical Pathway Enhancement:** BMI automatic referrals 23kg/m^2 and ensures professional interpreter services.

8.3 Equity-Focused Evaluation: Measuring Gap Reduction, Not Averages

The assessment shifts to the evaluation of decreased disparity of South Asian and White British people (Braveman, 2014). The traditional clinical measures are not used alone and power shift indicators and community-defined measures of success are used to make sure that what affected communities value are the focus of interventions.

9. Strengths & Weaknesses of the Epidemiological Approach

9.1 Methodological Strengths: Quantification for Resource Allocation

Epidemiology has the power to accurately estimate the burden, stratify risks, and track trends, which is required to allocate resources more precisely in resource-constrained systems (Rothman et al., 2008). It offers strong arguments that deserve attention and funding of the policy because it makes inequalities visible and measurable due to its quantitative rigor.

9.2 Epistemological Limitations: Reductionism and Categorical Violence

There is a danger of the epidemiological approach having what Smith (2021) has called epistemological violence, which consists of it turning intricate social suffering into statistical variations that deny the experience of lived suffering. Ethnic classification is a way of homogenizing different communities whereas biomedical framing is centered on disease, rather than wellbeing and structural transformation.

9.3 Practical Constraints: Implementation Barriers in Real-World Settings

The obstacles to implementation in Tower Hamlets are 25% population mobility, 40% lack of digital access and capacities in primary care services, which pose difficulties to implementation due to the often unaccounted epidemiological models (Exworthy et al., 2002). Diabetes is hardly ever a disease untouched by multimorbidity and it needs holistic approaches and not disease-specific approaches which epidemiology is finding hard to embrace.

10. Conclusion: An Epidemiological Understanding to Structural Transformation.

This epidemiological discussion of Type 2 Diabetes among South Asian adults in Tower Hamlets is an indication of a crisis that is not only a matter of genetic fate or personal decision making, but a systematic functioning of structural determinants: concentrated deprivation, obesogenic environments, linguistic marginalization and institutional discrimination. The conventional surveillance and intervention paradigms are quantifying the disparity and tend to recreate it because they fail to address these underlying factors and community voice.

It is necessary to change the direction towards political-economic intervention instead of biomedical management. The suggested Tower Hamlets Diabetes Equity Initiative provides an example of such a change that aims to address both clinical pathways and food environment using community co-production. Its success, though, requires going out of pilot projects to entrenched, responsible policy change - ring-fenced funding, required cultural safety and community-based data sovereignty.

Finally, the decrease in disproportionate burden of T2D is not only a clinical or populace health objective, but a social justice challenge. It requires that epidemiology should be decolonised, that is, it should be turned into what it measures to become an agent of the structural redistribution of power, resource and health

10. Conclusion: An Epidemiological Understanding to Structural Transformation.

This epidemiological discussion of Type 2 Diabetes among South Asian adults in Tower Hamlets is an indication of a crisis that is not only a matter of genetic fate or personal decision making, but a systematic functioning of structural determinants: concentrated deprivation, obesogenic environments, linguistic marginalization and institutional discrimination. The conventional surveillance and intervention paradigms are quantifying the disparity and tend to recreate it because they fail to address these underlying factors and community voice.

It is necessary to change the direction towards political-economic intervention instead of biomedical management. The suggested Tower Hamlets Diabetes Equity Initiative provides an example of such a change that aims to address both clinical pathways and food environment using community co-production. Its success, though, requires going out of pilot projects to entrenched, responsible policy change - ring-fenced funding, required cultural safety and community-based data sovereignty.

Finally, the decrease in disproportionate burden of T2D is not only a clinical or populace health objective, but a social justice challenge. It requires that epidemiology should be decolonised, that is, it should be turned into what it measures to become an agent of the structural redistribution of power, resource and health.




Figure 5: Surveillance system limitations creating cycles of invisible inequality. Data completeness paradox leads to resource misallocation away from highest-need populations (Mathur et al., 2020).

REFERENCES

1. Bajpai, R. (2021) *The implementation gap in health equity policies*. Journal of Health Services Research & Policy, 26(2), pp.123-130. <https://doi.org/10.1177/1355819620984200>
2. Bambra, C., Lynch, J. and Smith, K.E. (2020) **The Unequal Pandemic: COVID-19 and Health Inequalities**. Bristol: Policy Press. <https://doi.org/10.51952/9781447361257>
3. Bhui, K., Lenguerrand, E., Maynard, M.J., Stansfeld, S.A. and Harding, S. (2015) *Does cultural integration explain a mental health advantage for adolescents?* International Journal of Epidemiology, 44(3), pp.968-979. <https://doi.org/10.1093/ije/dyv070>
4. Bracken, P., Thomas, P., Timimi, S., Asen, E., Behr, G., Beuster, C., Bhunnoo, S., Browne, I., Chhina, N., Double, D., Downer, S., Evans, C., Fernando, S., Garland, M.R., Hopkins, W., Huws, R., Johnson, B., Martindale, B., Middleton, H. and Moncrieff, J. (2012) *Psychiatry beyond the current paradigm*. British Journal of Psychiatry, 201(6), pp.430-434. <https://doi.org/10.1192/bjp.bp.112.109447>
5. Braveman, P. (2014) *What is health equity: and how does a life-course approach take us further toward it?* Maternal and Child Health Journal, 18(2), pp.366-372. <https://doi.org/10.1007/s10995-013-1226-9>
6. British Heart Foundation (2024) *Heart and Circulatory Disease Statistics 2024*. London: BHF. <https://www.bhf.org.uk/what-we-do/our-research/heart-statistics>
7. Crenshaw, K. (1989) *Demarginalizing the intersection of race and sex: A black feminist critique of antidiscrimination doctrine, feminist theory and antiracist politics*. University of Chicago Legal Forum, 1989(1), pp.139-167. <https://chicagounbound.uchicago.edu/uclf/vol1989/iss1/8>

8. Diabetes UK (2023) *Diabetes Prevalence by Ethnicity 2023*. London: Diabetes UK. <https://www.diabetes.org.uk/professionals/position-statements-reports/statistics>
9. Dorling, D. (2015) *Inequality and the 1%*. London: Verso Books. <https://www.versobooks.com/products/492-inequality-and-the-1>
10. Exworthy, M., Blane, D. and Marmot, M. (2002) *Tackling health inequalities in the United Kingdom: The progress and pitfalls of policy*. Health Services Research, 38(6), pp.1905-1922. <https://doi.org/10.1111/1475-6773.01036>
11. Farmer, P. (2005) *Pathologies of Power: Health, Human Rights, and the New War on the Poor*. Berkeley: University of California Press. <https://www.ucpress.edu/book/9780520243262/pathologies-of-power>
12. Frohlich, K.L. and Potvin, L. (2008) *Transcending the known in public health practice: The inequality paradox: The population approach and vulnerable populations*. American Journal of Public Health, 98(2), pp.216-221. <https://doi.org/10.2105/AJPH.2007.114777>
13. General Medical Council (2020) *Good Medical Practice*. London: GMC. <https://www.gmc-uk.org/ethical-guidance/ethical-guidance-for-doctors/good-medical-practice>
14. Green, L.W. and Glasgow, R.E. (2006) *Evaluating the relevance, generalization, and applicability of research: Issues in external validation and translation methodology*. Evaluation & the Health Professions, 29(1), pp.126-153. <https://doi.org/10.1177/0163278705284445>
15. Greenhalgh, T., Voisey, C. and Robb, N. (2015) *Interpreted consultations as 'business as usual'? An analysis of organisational routines in general practices*. Sociology of Health & Illness, 37(3), pp.430-444. <https://doi.org/10.1111/1467-9566.12208>
16. Hales, C.N. and Barker, D.J.P. (1992) *Type 2 (non-insulin-dependent) diabetes mellitus: the thrifty phenotype hypothesis*. Diabetologia, 35(7), pp.595-601. <https://doi.org/10.1007/BF00400248>
17. Hanif, W., Khunti, K., Bellary, S., Bharaj, H.S., Karamat, M.A., Patel, K. and Singh, B.M. (2019) *Type 2 diabetes in the UK South Asian population: An update from the*

- South Asian Health Foundation. *Diabetic Medicine*, 36(8), pp.952-962. <https://doi.org/10.1111/dme.13933>
18. Hex, N., Bartlett, C., Wright, D., Taylor, M. and Varley, D. (2012) *Estimating the current and future costs of Type 1 and Type 2 diabetes in the UK, including direct health costs and indirect societal and productivity costs*. *Diabetic Medicine*, 29(7), pp.855-862. <https://doi.org/10.1111/j.1464-5491.2012.03698.x>
 19. International Diabetes Federation (2021) *IDF Diabetes Atlas, 10th edition*. Brussels: IDF. <https://diabetesatlas.org/>
 20. Israel, B.A., Schulz, A.J., Parker, E.A. and Becker, A.B. (1998) *Review of community-based research: Assessing partnership approaches to improve public health*. *Annual Review of Public Health*, 19, pp.173-202. <https://doi.org/10.1146/annurev.publhealth.19.1.173>
 21. Joseph Rowntree Foundation (2023) *UK Poverty 2023: The Essential Guide to Understanding Poverty in the UK*. York: JRF. <https://www.jrf.org.uk/report/uk-poverty-2023>
 22. Khunti, K., Kumar, S. and Brodie, J. (2019) *Diabetes UK and South Asian Health Foundation recommendations on diabetes research priorities for British South Asians*. *Diabetic Medicine*, 36(8), pp.942-951. <https://doi.org/10.1111/dme.13962>
 23. King's Fund (2023) *Tackling health inequalities on a system-wide basis*. London: The King's Fund. <https://www.kingsfund.org.uk/publications/tackling-health-inequalities-system-wide-basis>
 24. Kleinman, A. (1978) *Concepts and a model for the comparison of medical systems as cultural systems*. *Social Science & Medicine*, 12(2B), pp.85-93. [https://doi.org/10.1016/0160-7987\(78\)90014-5](https://doi.org/10.1016/0160-7987(78)90014-5)
 25. Krieger, N. (2005) *Embodiment: A conceptual glossary for epidemiology*. *Journal of Epidemiology & Community Health*, 59(5), pp.350-355. <https://doi.org/10.1136/jech.2004.024562>
 26. Krieger, N. (2012) *Methods for the scientific study of discrimination and health: An ecosocial approach*. *American Journal of Public Health*, 102(5), pp.936-944. <https://doi.org/10.2105/AJPH.2011.300544>

27. Lewis, G.A., Hughes, D.M., Irving, G., Wilding, J. and Hardy, K. (2025) *Association between diabetes self-management education attendance, hospital admissions and mortality in type 2 diabetes: A cohort analysis protocol*. Diabetes, Obesity & Metabolism, 27(5), pp.2377-2386. <https://doi.org/10.1111/dom.16257>
28. Link, B.G. and Phelan, J. (1995) *Social conditions as fundamental causes of disease*. Journal of Health and Social Behavior, 35, pp.80-94. <https://doi.org/10.2307/2626958>
29. Marmot, M. (2020) *Health Equity in England: The Marmot Review 10 Years On*. London: Institute of Health Equity. <https://www.health.org.uk/publications/reports/the-marmot-review-10-years-on>
30. Mathur, R., Hull, S.A., Badrick, E. and Robson, J. (2020) *Ethnic inequalities in the management and outcomes of diabetes in the UK: A scoping review*. Diabetic Medicine, 37(3), pp.427-435. <https://doi.org/10.1111/dme.14185>
31. Mental Health Foundation (2020) *Mental Health Statistics: Ethnicity*. London: MHF. <https://www.mentalhealth.org.uk/statistics/mental-health-statistics-ethnicity>
32. Metzl, J.M. and Hansen, H. (2014) *Structural competency: Theorizing a new medical engagement with stigma and inequality*. Social Science & Medicine, 103, pp.126-133. <https://doi.org/10.1016/j.socscimed.2013.06.032>
33. Nazroo, J.Y. (2022) *Health and Social Inequalities in Ethnic Minority Groups in the UK*. Manchester: Manchester University Press. <https://doi.org/10.7765/9781526160598>
34. NHS Digital (2023) *National Diabetes Audit 2022-23*. Leeds: NHS Digital. <https://digital.nhs.uk/data-and-information/publications/statistical/national-diabetes-audit>
35. NHS England (2021) *NHS Long Term Plan: Implementing CORE20PLUS5*. London: NHS England. <https://www.england.nhs.uk/about/equality/equality-hub/national-healthcare-inequalities-improvement-programme/core20plus5/>
36. NICE (2022) *Type 2 diabetes: prevention in people at high risk*. London: National Institute for Health and Care Excellence. <https://www.nice.org.uk/guidance/ph38>
37. Office for National Statistics (2021) *Population estimates by ethnic group, England and Wales: 2011 to 2021*. London:

ONS. <https://www.ons.gov.uk/peoplepopulationandcommunity/populationandmigration/populationestimates/articles/populationestimatesbyethnicgroupandreligionenglandandwales/2019>

38. Parveen, S., Morrison, V. and Robinson, C.A. (2018) *Ethnic differences in the caregiver role: A qualitative study of perception and experiences of caring for a relative with heart failure*. *Health & Social Care in the Community*, 26(3), pp.453-460. <https://doi.org/10.1111/hsc.12538>
39. Popkin, B.M. (2006) *Global nutrition dynamics: The world is shifting rapidly toward a diet linked with noncommunicable diseases*. *American Journal of Clinical Nutrition*, 84(2), pp.289-298. <https://doi.org/10.1093/ajcn/84.2.289>
40. Public Health England (2020) *Health profile for England 2020*. London: PHE. <https://www.gov.uk/government/publications/health-profile-for-england-2020>
41. Ramsden, I. (2002) *Cultural safety and nursing education in Aotearoa and Te Waipounamu*. PhD Thesis, Victoria University of Wellington. <https://doi.org/10.26686/wgtn.16946428>
42. Rothman, K.J., Greenland, S. and Lash, T.L. (2008) *Modern Epidemiology*. 3rd ed. Philadelphia: Lippincott Williams & Wilkins.
43. Salway, S., Mir, G., Turner, D., Ellison, G.T.H., Carter, L. and Gerrish, K. (2016) *Obstacles to "race equality" in the English National Health Service: Insights from the healthcare commissioning arena*. *Social Science & Medicine*, 152, pp.102-110. <https://doi.org/10.1016/j.socscimed.2016.01.031>
44. Sanghera, D.K., Been, L.F., Ralhan, S., Wander, G.S., Mehra, N.K., Singh, J., Mulvihill, J.J., Aston, C.E. and Sanghera, N.T. (2018) *TCF7L2 polymorphisms and type 2 diabetes in South Asians*. *Diabetes*, 67(3), pp.487-492. <https://doi.org/10.2337/db17-0855>
45. Singer, M. (2009) *Introduction to Syndemics: A Systems Approach to Public and Community Health*. San Francisco: Jossey-Bass.
46. Smith, L.T. (2021) *Decolonizing Methodologies: Research and Indigenous Peoples*. 3rd ed. London: Zed Books. <https://www.zedbooks.net/shop/book/decolonizing-methodologies/>

47. Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P., Green, J., Landray, M., Liu, B., Matthews, P., Ong, G., Pell, J., Silman, A., Young, A., Sprosen, T., Peakman, T. and Collins, R. (2015) *UK Biobank: An open access resource for identifying the causes of a wide range of complex diseases of middle and old age*. PLoS Medicine, 12(3), e1001779. <https://doi.org/10.1371/journal.pmed.1001779>
48. Swinburn, B.A., Sacks, G., Hall, K.D., McPherson, K., Finegood, D.T., Moodie, M.L. and Gortmaker, S.L. (2011) *The global obesity pandemic: Shaped by global drivers and local environments*. The Lancet, 378(9793), pp.804-814. [https://doi.org/10.1016/S0140-6736\(11\)60813-1](https://doi.org/10.1016/S0140-6736(11)60813-1)
49. Tower Hamlets Council (2022) *Joint Strategic Needs Assessment 2022: Diabetes Report*. London: Tower Hamlets Council. https://www.towerhamlets.gov.uk/lgnl/health_and_social_care/public_health/jsna.aspx
50. Trivedi, P. (2022) *Language access as a right, not a privilege: Overcoming linguistic barriers in healthcare*. Journal of Medical Ethics, 48(4), pp.221-225. <https://doi.org/10.1136/medethics-2020-107146>
51. Zaman, M.J., Philipson, P., Chen, R., Farag, A., Shipley, M., Marmot, M.G. and Luben, R.N. (2017) *South Asians and coronary heart disease: A review of the evidence on risk factors, access to care and outcomes*. European Heart Journal, 38(14), pp.1066-1074. <https://doi.org/10.1093/eurheartj/ehx050>