

Name Shabina Kousar

Student ID 2009301

INTRODUCTION -

Mycobacterium tuberculosis infection, in particular, is what causes tuberculosis in humans. People who have pulmonary or laryngeal TB can cough, sneeze, or even just converse without producing airborne droplets. Inhaled infectious droplets settle in the alveoli, where macrophages take up the bacilli, starting a chain of events that either stops the infection from spreading or causes it to become active (Frieden and others 2003). After being ingested by macrophages, M. tuberculosis replicates slowly but consistently and travels to the hilar lymph nodes via the lymphatic system. Cell-mediated immunity, which is connected to a positive tuberculin test, usually appears two to eight weeks after infection in infected individuals. Granulomas are formed by activated macrophages and T lymphocytes, which stop bacilli from multiplying and spreading further. The infection stays contained within the granuloma unless a later failure in cell-mediated immunity happens.

Strategies used in the UK to measure the tuberculosis

Even if the number of cases has recently dropped by 32% from a peak of 8,280 in 2011 to 5,664 cases in 2021, England's TB rates remain unacceptably high compared to other Western European nations. This decline is mainly attributable to a drop in the proportion of cases originating outside of the UK. However, the most current data from 2021 revealed a slowdown of the drop in case numbers (1% versus 10% year-over-year earlier), with just a slight decline in cases of UK births and no decline in cases of non-UK births for TB rates by UK local authority. TB is unequally distributed, with some subgroups being disproportionately afflicted, such as recent immigrants and those with socioeconomic risk factors. With 39% of all cases in England, London has the highest infection burden, followed by the West Midlands at 13%. Nearly 75% of cases are discovered in individuals who were born abroad in nations where TB is more prevalent; the bulk of these individuals are from South Asia (51% of the non-UK-born TB population) and sub-Saharan Africa (22 percent). In the UK, the National Health Service (NHS) and Public Health England (PHE) work together in an integrated manner. All cases of TB disease that may be contagious are reported to PHE and are then screened by clinical TB nursing teams, who identify household members and other

close contacts within NHS facilities. 4,758 TB cases were reported in England in 2022. Since a long time ago, it has been national policy to screen household members and other close contacts of these cases for TB in order to find any further cases of the disease as well as any TB-infected people who would benefit from treatment.

Strategies used in the UK to monitor

Provisional TB data reports for England are currently released on a monthly and quarterly basis. These enable a quicker evaluation of TB activity at the national and local levels. To replace and enhance current systems, PHE has provided funding for the creation of a new TB surveillance system (NTBS). PHE Field services have access to early releases, which give more specific live information from TB notifications. The NTBS will begin to be fully implemented in 2021, and this will include direct access to TB clinics.

The PHE TB Fingertips tool can be used to view the TB Strategy monitoring indicators, which were created by PHE and released on World TB Day 2015 to allow interactive data viewing at the local authority and CCG levels. Each year, these are updated.

The 2016 publication "Health matters - lowering the burden of TB" included new TB infographics, blogs, a presentation set, and a video.

Since "Fingertips" did not specifically track each of the 10 "areas for action," "Strategy Progress Measures" were created to track their implementation at the national and TB Control Board levels. 9 of 22 progress measures had been met by the end of the Strategy's second year by the end of 2016, and this number rose yearly to 18 of 24 in 2019.

Based on information from the yearly TB report, annual TB infographics were created. Along with the yearly TB report, these are provided by the national and regional TB Control Boards. To address the gaps in public health and clinical services, such as the enormous reservoir of latent TB infection, it is necessary to provide clear accountability mechanisms at local and national levels, develop suitable commissioning frameworks, and meet the needs of underserved communities.

Strategies used in the UK to control the tuberculosis

Among the nations of western Europe, the UK has the second-highest tuberculosis prevalence rate. More cases of tuberculosis are treated in London clinics each year than in the capital cities of all other western European countries combined. Currently, tuberculosis rates in the UK are over five times greater than those in the USA.

With examples of cases acquired in other countries, the UK's lack of progress in tuberculosis control not only poses a threat to domestic public health but also an international embarrassment leading to illnesses in other low-incidence countries is the UK countries. Public

Health England spearheaded a coalition of partners in creating the national Tuberculosis Oversight Group as a venue for sharing innovation and best practices across local, regional, and national health authorities in response to this intolerable trend.

With the establishment of tuberculosis control boards and thorough cohort reviews in numerous locations as a result of these conversations, tuberculosis has become a top concern for Public Health England. However, not all regions of England have adopted improved tuberculosis control strategies, and there is still unacceptable heterogeneity in the standard of clinical and public health interventions. A collaborative tuberculosis strategy has been created by Public Health England and NHS England in collaboration with key stakeholders to address this issue. The approach intends to reduce tuberculosis year after year with the ultimate goal of curing the illness as a public health issue. These objectives would be accomplished by taking action in ten key areas, including bettering access and early diagnosis; providing high-quality diagnostic, treatment, and care services; better contact tracing and immunization of vulnerable people; combating drug resistance; controlling tuberculosis in underserved populations; screening and treating new immigrants for latent tuberculosis infection; effective surveillance and monitoring; and workforce planning. The plan acknowledges the global nature of tuberculosis and supports global efforts to control the illness as well as the need for research that would enable the WHO's aim of universal eradication. The disease's social and economic causes, as well as the availability of quality public health services.

To achieve a persistent decline, issues with and clinical services across England will need to be addressed. The collaborative tuberculosis strategy's success will serve as a crucial litmus test for whether and how vital organizations can collaborate to bring about the improvements in health and social services needed to effectively stem the tide of tuberculosis in England.