

Name: Jahmai Simone Howe

Student ID: 2507679

Module Code: LCHS4008

Module: Report on a Health Issue

Lecture: Deborah Odeh

Introduction

A stroke is a serious medical condition that happens when the blood supply to part of the brain is suddenly interrupted, either due to a blockage (ischaemic stroke) or a burst blood vessel (haemorrhagic stroke). This results in brain cell damage because the brain is not receiving enough oxygen. Common signs and symptoms include facial drooping, weakness in one arm, slurred speech, confusion, and problems with balance or coordination. In many cases, these symptoms appear suddenly, which is why stroke is considered a medical emergency that requires immediate attention.

Stroke is a major global health issue. Worldwide, around 12.2 million people experience a stroke each year, with over 110 million people living with its effects (World Stroke Organisation, 2022, cited in Cardiff University, 2023). In the UK, approximately 100,000 strokes occur annually, and over 1.1 million people live with stroke (British Heart Foundation, 2023). This demonstrates how common the condition is and the pressure it places on health and social care services. It also emphasises the importance of early detection, treatment, and prevention in reducing both human and economic impact.

Main Body

There are several ways used to measure and monitor stroke at both individual and population levels. At a national level, surveillance systems such as hospital records, audits, and public health data are used to track how many people are affected, mortality rates, and long-term outcomes (NHS, 2023). These systems help healthcare services identify trends and patterns, allowing resources to be used more effectively and care to be improved.

For example, stroke registries and national audits collect detailed information on patient outcomes, recovery progress, and treatment effectiveness. In the UK, the Sentinel Stroke National Audit Programme (SSNAP) is used to assess the quality of stroke care and patient outcomes (Royal College of Physicians, 2023). Monitoring also involves identifying risk factors such as high blood pressure, diabetes, and atrial fibrillation, which are strongly linked to stroke risk (NHS, 2023). This enables healthcare professionals to focus on prevention and

early intervention. Global monitoring systems also help to show the overall impact of stroke and support prevention strategies (World Stroke Organization, 2022).

On an individual level, clinical assessments such as CT or MRI scans, blood pressure monitoring, and neurological examinations are used to diagnose stroke and track recovery. These methods allow healthcare professionals to make accurate and timely decisions about treatment.

Stroke is managed through a combination of prevention, early treatment, and long-term care. Prevention focuses on reducing risk factors before a stroke occurs. This includes promoting a healthy lifestyle such as maintaining a balanced diet, taking regular exercise, reducing alcohol intake, and stopping smoking (Heart Research Institute UK, 2024). These changes help to reduce conditions like high blood pressure and high cholesterol, which are major risk factors. Research suggests that up to 80% of strokes could be prevented by managing these risks effectively (Heart Research Institute UK, 2024).

Medication is also important in controlling risk factors such as blood pressure, cholesterol, and blood clotting. Secondary prevention is used after someone has already experienced a stroke or a transient ischaemic attack (TIA), to reduce the risk of another stroke (NICE, 2022). This may include anticoagulants and statins, alongside ongoing monitoring.

Early treatment plays a crucial role in improving outcomes. Rapid access to emergency care, such as thrombolysis or thrombectomy, can reduce brain damage and improve recovery (NHS, 2023). The quicker treatment is given, the better the chances of reducing long-term disability.

The impact of stroke on individuals is often long-lasting. Physically, many people experience paralysis, mobility problems, and difficulties with speech and swallowing. Cognitive issues are also common, with around 22% of stroke survivors experiencing cognitive impairment, and some developing depression or anxiety (NIHR, 2022). These effects can significantly reduce independence and quality of life.

Stroke can also affect people socially. Individuals may struggle to communicate or take part in everyday activities, which can lead to isolation. Many people are unable to return to work, resulting in financial difficulties and increased reliance on support services. Families and carers are also affected, often experiencing stress and increased responsibilities (Stroke Association, 2023).

At a wider level, stroke places a significant demand on health and social care services. In England, it is responsible for thousands of deaths each year and causes long-term disability for many individuals (British Heart Foundation, 2023). The cost of care is also high, including hospital treatment, rehabilitation, and ongoing support. These costs are expected to increase due to an ageing population and rising risk factors such as obesity and diabetes (NHS, 2023). Inequalities are also evident, with higher rates of stroke in more deprived areas and certain ethnic groups, highlighting the need for targeted support.

A range of health and social care services are required to support people after a stroke. Following a stroke, patients often need care in a hospital, usually within a specialised stroke unit. Rehabilitation is a key part of recovery and includes physiotherapy to improve movement, occupational therapy to support daily living, and speech and language therapy to assist with communication (NHS, 2023). Rehabilitation may continue for a long period and can take place at home or within the community.

Community support plays an important role in helping individuals regain independence. Organisations such as the Stroke Association provide emotional support, communication help, and rehabilitation programmes for both patients and their families (Stroke Association, 2023). Social care services also provide support with personal care, home adaptations, and access to local services.

Another key consideration is the role of rehabilitation in supporting long-term recovery. Rehabilitation focuses not only on physical recovery but also on helping individuals adjust to changes in their daily lives. Recovery can take time, and progress will vary depending on the severity of the stroke and the individual's overall health (NHS, 2023). Some people may recover quickly, while others may need long-term support.

A person-centred approach is essential because each individual has different needs and goals. Care plans should be tailored to the person, taking into account their abilities, communication needs, and emotional wellbeing. Involving individuals in their own care can improve motivation and lead to a more positive recovery.

Family involvement is also important in rehabilitation. Families and carers often provide essential support, but this can place emotional and physical pressure on them. Support services are therefore needed to help carers manage their responsibilities. Providing carers with advice, training, and emotional support can improve outcomes for both the patient and their family (Stroke Association, 2023).

Another important aspect of stroke care is early intervention and prevention programmes. Healthcare services aim to identify individuals who are at higher risk of stroke and provide

support before a stroke occurs. This can include regular health checks and monitoring risk factors such as blood pressure and cholesterol (NHS, 2023). Early intervention helps reduce the likelihood of a stroke and ensures that risk factors are managed effectively over time.

Public health campaigns also play a key role in raising awareness about stroke and its warning signs. Educating the public about symptoms encourages people to seek medical help quickly, which can significantly improve survival rates and recovery outcomes. Campaigns such as FAST are especially important because the faster a stroke is treated, the less damage is caused to the brain (NHS, 2023)

In addition, early recognition of stroke symptoms plays a crucial role in improving outcomes. Public awareness campaigns such as FAST (Face, Arms, Speech, Time) help people identify the warning signs of a stroke quickly and seek urgent medical help. This reduces delays in treatment and can significantly limit brain damage, improving the chances of recovery and reducing long-term complications.

Conclusion

In conclusion, stroke is a serious health condition that has a major impact on individuals, families, and society. It can be measured and monitored using hospital records, national audits, and clinical assessments, which help healthcare professionals understand how many people are affected and how well treatments are working. Effective monitoring allows services to improve care and respond more quickly to patient needs.

Stroke is managed through prevention, early treatment, and ongoing care, all of which help to reduce the risk of further strokes and support recovery. However, delays in treatment can lead to more severe disability, showing the importance of acting quickly. Stroke can also result in long-term effects such as physical disability, communication difficulties, and emotional challenges, which can affect independence and quality of life.

This can place pressure on families and health and social care services. Overall, stroke requires a coordinated and joined-up approach to ensure that individuals receive the best possible care and support throughout their recovery.

Reference List

British Heart Foundation (2023) *UK factsheet: Stroke statistics*. Available at:

<https://www.bhf.org.uk/what-we-do/our-research/heart-statistics>

British Heart Foundation (2023) *Stroke statistics*. Available at:

<https://www.bhf.org.uk/information-support/conditions/stroke>

Cardiff University (2023) *Forecasted prevalence and incidence of long-term conditions*. Available at:

<https://orca.cardiff.ac.uk/id/eprint/160640/>

National Institute for Health and Care Excellence (NICE) (2022) *Stroke and transient ischaemic attack in over 16s: diagnosis and initial management*. Available at:

<https://www.nice.org.uk/guidance/cg68>

NHS (2023) *Stroke*. Available at:

<https://www.nhs.uk/conditions/stroke/>

Stroke Association (2023) *Life after stroke*. Available at:

<https://www.stroke.org.uk>

World Stroke Organization (2022) *Global stroke facts and figures*. Available at:

<https://www.world-stroke.org/world-stroke-day-campaign/facts-and-figures>

NHS (2023) *Stroke*. Available at:

<https://www.nhs.uk/conditions/stroke/>

Stroke Association (2023) *Life after stroke*. Available at:

<https://www.stroke.org.uk>