

Cover Page

Student's Name: Segun Victor Ikudehinbu

Student ID: 212 0079

Module Name: Introduction to Public Health

Module code: LCHS4004

Lecturer Name: Bianca Sanfilippo

Introduction

Infectious diseases are caused by microorganisms, such as bacteria, viruses, fungi and parasites. These organisms can be spread through contact with an infected person, or through contact with contaminated food, water or surfaces. Infectious diseases can cause a wide range of symptoms and effects, from mild to severe (Kotra, 2007).

Approaches and Strategies to measure outbreak

Measuring an infectious disease outbreak, such as TB, requires a combination of approaches including epidemiological surveillance, laboratory testing, and contact tracing (Impouma et al., 2020). Epidemiological surveillance involves identifying new cases of TB by collecting data from healthcare settings or other sources promptly to identify trends and assess the effectiveness of control measures. Laboratory testing is necessary to diagnose active TB in individuals who present with symptoms and make sure they receive the appropriate treatment. Contact tracing involves interviewing those who have been identified as having been in contact with someone infected with that illness.

Additionally, public health officials must put effective measures in place to prevent further transmission of the disease, such as ensuring individuals with TB are provided with appropriate treatment and support, educating the public about the disease, maintaining good hygiene standards, and taking precautions such as wearing face masks.

Approaches and Strategies to Monitor infectious outbreak

One of the main strategies for monitoring an infectious disease outbreak is surveillance. This involves regular collection and analysis of data about cases or outbreaks, which can help to identify trends in incidence, geographic distribution, risk factors, and preventive measures that could be taken (Threats, 2007). It also allows for early detection to help contain the spread of the

illness. Active case finding can also be employed to detect new cases; this includes contact tracing and using public health teams to actively search for individuals who may have been exposed or infected.

Other approaches include laboratory testing, such as smear microscopy and polymerase chain reaction (PCR) tests. Vaccines may also be used to prevent infection among vulnerable populations.

Main approaches and strategies to control infectious diseases

In the case of tuberculosis (TB) infection, the main objectives are to reduce transmission and facilitate treatment. This can be achieved through a combination of strategies such as public health education, contact tracing and screening, social support, and improved access to healthcare (Nakatani, 2016). Public health education is important for raising awareness about TB symptoms, diagnosis, and treatment options. Contact tracing and screening can help identify those who may have come in contact with a person infected with TB, allowing for early intervention. Social support can be critical in helping patients access medical care and adhere to their treatment plans. Finally, access to healthcare should be improved to ensure that those suffering from TB receive prompt diagnosis and appropriate treatment (Anderson et al., 2013).

UK policies and strategies

The UK government has a range of policies and strategies available to maintain public health. These include increasing awareness of the risk factors associated with TB, providing access to screening and treatment for those at high risk, and introducing targeted immunization programs (Public Health England, 2019). In addition, the government can increase funding for research into new treatments and diagnostic tests. Furthermore, initiatives such as the National Tuberculosis

Elimination Program (NTEP) can offer support to those living with the condition and raise awareness of its potential dangers. Finally, a collaboration between local authorities, health professionals, and community organizations is essential to effectively reduce the risk of TB for all members of society (Lipman & White, 2015). Such collaborations are likely to include sharing best practices and working together to ensure TB services are provided to those who need them.

If UK government public health policies have aided or challenged agencies and local communities

The UK government's public health policies have been a vital tool in helping tackle the spread of infectious diseases such as tuberculosis (TB). For example, the policy of contact tracing allows for early identification and treatment of those infected to reduce further transmission (Gray et al., 2006). Furthermore, targeted vaccinations are provided to marginalized communities which are at higher risk of contracting TB. Additionally, healthcare professionals are required to adhere to specific guidelines when a suspected TB case is reported.

However, the UK government's public health policies have not been without challenges, particularly regarding dealing with infectious outbreaks in local communities. For instance, the lack of access to adequate healthcare resources can make it difficult for individuals living in poverty to receive the medical treatment they need. Furthermore, language barriers can hinder communication between healthcare professionals and those from diverse backgrounds, making it difficult to provide accurate diagnoses and advice (Suk & Semenza, 2011)

First action when addressing the case study

When addressing the outbreak in the case study, the first action should be to assess Mr. Patel's risk factors for tuberculosis (TB). This includes taking a detailed history of his travel and living arrangements, as well as any contact with people who are known to have TB or at high risk

of having it (CDC, 2019). It is also important to consider any underlying health conditions he may have, such as HIV or diabetes, which can increase his risk of developing TB.

Once the risk factors are identified, the next action is to perform a physical examination to assess any signs or symptoms that could be suggestive of TB. This could include checking for a fever, coughing up blood-tinged sputum, or enlarged lymph nodes. It is also important to assess any chest X-ray results, if available.

Finally, a TB test should be administered to confirm the diagnosis and determine if it is active or latent (Auguste et al., 2021). This should include a sputum culture and drug susceptibility tests to determine the best treatment plan. Once the diagnosis is confirmed, the appropriate action should be taken to prevent the spread of the disease.

Conclusion

In conclusion, the UK government's public health policies have been instrumental in supporting and challenging agencies and local communities when dealing with infectious outbreaks such as tuberculosis (TB). When addressing a suspected TB case, the first action should be to assess any risk factors and perform a physical examination. A test should then be administered to confirm the diagnosis and determine the best treatment plan.

References

- Anderson, C., White, J., Abubakar, I., Lipman, M., Tamne, S., Anderson, S. R., deKoningh, J., & Dart, S. (2013). Raising standards in UK TB control: introducing cohort review: Table 1. *Thorax*, 69(2), 187–189. <https://doi.org/10.1136/thoraxjnl-2013-203751>
- Auguste, P. E., Mistry, H., McCarthy, N. D., Sutcliffe, P. A., & Clarke, A. E. (2021). Cost-effectiveness of testing for latent tuberculosis infection in people with HIV. *AIDS*, 36(1), 1–9. <https://doi.org/10.1097/qad.0000000000003060>
- CDC. (2019). *TB Risk Factors*. Centers for Disease Control and Prevention. <https://www.cdc.gov/tb/topic/basics/risk.htm>
- Gray, S., Pilkington, P., Pencheon, D., & Jewell, T. (2006). Public Health in the UK: Success or Failure? *Journal of the Royal Society of Medicine*, 99(3), 107–111. <https://doi.org/10.1177/014107680609900309>
- Impouma, B., Roelens, M., Williams, G. S., Flahault, A., Codeço, C. T., Moussana, F., Farham, B., Hamblion, E. L., Mboussou, F., & Keiser, O. (2020). Measuring Timeliness of Outbreak Response in the World Health Organization African Region, 2017–2019. *Emerging Infectious Diseases*, 26(11), 2555–2564. <https://doi.org/10.3201/eid2611.191766>
- Kotra, L. P. (2007). *Infectious Diseases - an overview* | *ScienceDirect Topics*. www.sciencedirect.com. <https://www.sciencedirect.com/topics/immunology-and-microbiology/infectious-diseases>
- Lipman, M., & White, J. (2015). Collaborative tuberculosis strategy for England. *BMJ*, 350(feb19 16), h810–h810. <https://doi.org/10.1136/bmj.h810>
- Nakatani, H. (2016). Global Strategies for the Prevention and Control of Infectious Diseases and Non-Communicable Diseases. *Journal of Epidemiology*, 26(4), 171–178. <https://doi.org/10.2188/jea.je20160010>

Public Health England. (2019, September 10). *PHE infectious diseases strategy*. GOV.UK.

<https://www.gov.uk/government/publications/phe-infectious-diseases-strategy>

Suk, J. E., & Semenza, J. C. (2011). Future Infectious Disease Threats to Europe. *American Journal of Public Health*, 101(11), 2068–2079. <https://doi.org/10.2105/AJPH.2011.300181>

Threats, I. of M. (US) F. on M. (2007). Strategies for Disease Containment. In www.ncbi.nlm.nih.gov.

National Academies Press (US).

<https://www.ncbi.nlm.nih.gov/books/NBK54163/#:~:text=Surveillance%20strategies%20include%20rapid%20diagnosis>