



Module name: Introduction to Public Health

Module code: LCHS4008

Student name: Gabriela Harissis

Student ID: 2509668

Lecturer's name: Arif Ahmed

Introduction

Memory, thinking, behavior, and the capacity to carry out daily tasks are all impacted by dementia, a degenerative neurological disorder. It is a syndrome brought on by a variety of illnesses, most frequently Alzheimer's disease, rather than a single sickness. The World Health Organization states that dementia is caused by brain cell destruction that results in a reduction in cognitive functioning that goes beyond what is typical of aging (World Health Organization, 2023).

Memory loss, disorientation, communication difficulties, poor judgment, and behavioral or emotional changes are typical signs and symptoms. Forgetting recent events, becoming lost in familiar surroundings, or having trouble following conversations are some of the early symptoms (NHS, 2023a).

Dementia is a significant global health issue. Around 57 million people worldwide were living with dementia in 2021, with nearly 10 million new cases each year (World Health Organization, 2023). In the UK, approximately 982,000 people are living with dementia, and this number is expected to rise to 1.4 million by 2040 (NHS Digital, 2023). These figures highlight the growing severity of dementia as a public health concern.

Approaches Used to Measure and Monitor Dementia

Dementia is measured and monitored using a variety of clinical, psychological, and technological approaches. These methods help healthcare professionals diagnose the condition, track its progression, and plan appropriate care.

One key method is cognitive assessment tools, such as memory and thinking tests. These tests measure abilities like recall, attention, language, and problem-solving. They are often used during initial diagnosis and repeated over time to monitor decline (NICE, 2018).

Another important approach is brain imaging techniques, including MRI and CT scans. These scans allow professionals to detect physical changes in the brain, such as shrinkage or damage to brain cells. This helps differentiate between different types of dementia, such as Alzheimer's disease and vascular dementia (NHS, 2023a).

Medical history and physical examinations are also essential. Doctors assess symptoms, lifestyle factors, and underlying conditions like diabetes or high blood pressure, which can increase the risk of dementia (NHS, 2023a).

In addition, biomarkers and laboratory tests are increasingly used. Blood tests and cerebrospinal fluid analysis can help identify proteins linked to dementia, improving early diagnosis (NICE, 2018).

Another approach is functional assessments, which evaluate how well individuals can perform daily activities such as dressing, eating, or managing finances. These assessments help determine the severity of the disease, and the level of support required (Alzheimer's Society, 2022).

Finally, monitoring systems and digital health records are used within healthcare services. These systems allow professionals to track patient progress over time and ensure continuity of care across different services (NHS England, 2019).

How Dementia is Controlled

Although there is currently no cure for dementia, there are several ways to control and manage the condition to improve quality of life.

One key approach is medication. Certain drugs can help manage symptoms such as memory loss, confusion, and mood changes. For example, medications such as cholinesterase inhibitors may improve communication between brain cells or slow down cognitive decline in some individuals (NHS, 2023b).

Another important method is lifestyle management. Research shows that maintaining a healthy lifestyle can reduce the risk or slow progression of dementia. This includes regular physical activity, a balanced diet, avoiding smoking and excessive alcohol consumption, and managing conditions such as hypertension and diabetes (NHS, 2022).

Psychological and cognitive therapies are also used. Activities such as memory training, cognitive stimulation therapy, and reminiscence therapy can help maintain mental function and improve wellbeing (NICE, 2018).

Person-centred care is essential in managing dementia. This approach focuses on understanding the individual's needs, preferences, and history to provide tailored support. It helps maintain dignity and independence for as long as possible (NHS England, 2019).

Additionally, environmental adjustments can help control the condition. For example, creating safe and familiar environments reduces confusion and anxiety. Simple changes such as clear signage, good lighting, and structured routines can make a significant difference (Alzheimer's Society, 2022).

Support for carers and families is also crucial. Since many individuals with dementia rely on informal care, providing education and support to carers helps improve overall management of the condition (NHS, 2023c).

Consequences of Dementia on People in the Community

Dementia has wide-ranging consequences that affect not only individuals but also families, carers, and the wider community.

One major consequence is loss of independence. As the disease progresses, individuals may struggle with basic tasks such as cooking, dressing, and personal hygiene. This increases reliance on others for support (NHS, 2023a).

Another significant impact is social isolation. People with dementia may withdraw from social activities due to confusion, embarrassment, or difficulty communicating. This can lead to loneliness and depression (Alzheimer's Society, 2022).

Dementia also has emotional and psychological effects. Individuals may experience anxiety, frustration, and mood changes as they become aware of their declining abilities. Behavioral changes, such as aggression or agitation, can also occur (NICE, 2018).

The condition places a heavy burden on families and carers. Many carers provide unpaid support, often spending several hours each day caring for their loved ones. This can lead to stress, burnout, and financial difficulties (NHS England, 2019).

There are also economic consequences. Dementia care is costly, involving healthcare services, social care, and informal care. Globally, dementia costs reached approximately \$1.3 trillion, highlighting its economic impact (World Health Organization, 2023).

In the wider community, dementia can lead to increased demand for health and social care services, including residential care homes and specialist support services. It is also associated with higher rates of hospital admissions and long-term care needs (NHS Digital, 2023).

Furthermore, individuals with dementia may face stigma and discrimination due to a lack of understanding about the condition. This can prevent them from seeking help and reduce their quality of life (Alzheimer's Society, 2022).

Health and Social Care Support Services Required

To meet the needs of individuals with dementia, a wide range of health and social care services are required within the community.

One essential service is primary healthcare, including GPs and memory clinics. These services provide diagnosis, treatment, and ongoing monitoring of the condition (NHS, 2023a).

Specialist dementia services are also important. These include neurologists, psychiatrists, and dementia nurses who provide expert care and support (NICE, 2018).

Home care services play a vital role in helping individuals remain in their own homes. Care workers assist with daily activities such as personal care, medication management, and meal preparation (NHS, 2023c).

Another key service is day care centres, which provide social interaction, structured activities, and respite for carers. These centres help reduce isolation and improve wellbeing (Alzheimer's Society, 2022).

Residential and nursing care homes are required for individuals with advanced dementia who need 24-hour support. These facilities provide a safe and supportive environment for those with complex needs (NHS England, 2019).

Support groups and counselling services are also important for both individuals and carers. These services provide emotional support, information, and coping strategies (Alzheimer's Society, 2022).

Additionally, assistive technologies such as alarms, GPS trackers, and medication reminders can help individuals maintain independence and safety (NHS, 2022).

Social services also play a key role in assessing needs and coordinating care packages. They ensure that individuals receive appropriate support based on their circumstances (NHS England, 2019).

Finally, community awareness and education programmes are important in reducing stigma and promoting understanding of dementia (Alzheimer's Society, 2022).

Conclusion

In conclusion, dementia is a progressive and life-changing condition that affects millions of people worldwide. It is characterized by a decline in memory, thinking, and the ability to carry out daily activities, with symptoms worsening over time. The growing number of cases globally

and within the UK highlights the seriousness of dementia as a major public health issue (World Health Organization, 2023; NHS Digital, 2023).

Various approaches are used to measure and monitor dementia, including cognitive tests, brain scans, and functional assessments. Although there is no cure, the condition can be managed through medication, lifestyle changes, and person-centered care (NICE, 2018). Dementia has significant consequences, including loss of independence, emotional distress, and increased pressure on families and healthcare services.

To support individuals effectively, a wide range of health and social care services is required, including medical care, home support, and community services. Addressing dementia requires a holistic and coordinated approach to improve quality of life for those affected and their families.

References

- Alzheimer's Society (2023) *What is dementia?* Available at: <https://www.alzheimers.org.uk/about-dementia> (Accessed: 14 April 2026).
- Alzheimer's Society (2023) *Dementia UK statistics*. Available at: <https://www.alzheimers.org.uk/about-us/policy-and-influencing/dementia-uk-statistics> (Accessed: 14 April 2026).
- National Institute for Health and Care Excellence (NICE) (2018) *Dementia: assessment, management and support for people living with dementia and their carers (NG97)*. Available at: <https://www.nice.org.uk/guidance/ng97> (Accessed: 14 April 2026).
- NHS (2023a) *Dementia - Overview*. Available at: <https://www.nhs.uk/conditions/dementia/> (Accessed: 14 April 2026).
- NHS (2023b) *Dementia - Treatment*. Available at: <https://www.nhs.uk/conditions/dementia/treatment/> (Accessed: 14 April 2026).
- NHS (2023c) *Dementia - Living with dementia*. Available at: <https://www.nhs.uk/conditions/dementia/living-with/> (Accessed: 14 April 2026).
- NHS (2022) *Dementia prevention*. Available at: <https://www.nhs.uk/conditions/dementia/prevention/> (Accessed: 14 April 2026).
- NHS Digital (2023) *Dementia profile*. Available at: <https://digital.nhs.uk/data-and-information/publications/statistical/dementia> (Accessed: 14 April 2026).
- NHS England (2019) *The NHS Long Term Plan*. Available at: <https://www.longtermplan.nhs.uk/> (Accessed: 14 April 2026).
- NHS England (2023) *Dementia care pathway*. Available at: <https://www.england.nhs.uk/dementia/> (Accessed: 14 April 2026).
- Public Health England (2019) *Dementia: applying All Our Health*. Available at: <https://www.gov.uk/government/publications/dementia-applying-all-our-health> (Accessed: 14 April 2026).
- World Health Organization (2023) *Dementia*. Available at: <https://www.who.int/news-room/fact-sheets/detail/dementia> (Accessed: 14 April 2026).

