



Prifysgol Cymru
Y Drindod Dewi Sant
University of Wales
Trinity Saint David

BMHP6029-Psychopathological Classifications and Provisions

Component 2: Written Assignment (Report)

Student name: Vannia Celia Delfino

Student number: 2322920

Lecture: Farah Wiita

Impulse Control Disorders: Classification, Treatment, and Emerging Issues

1. Introduction

Impulse control Disorders (ICDs) represent a clinically significant psychiatric disorder marked by an inability to control urges and drives or with temptation to perform an act that can be detrimental to the person, or to others (American Psychiatric Association, 2013). They have been comparatively poorly diagnosed and poorly treated in mental healthcare in the United Kingdom and other regions of the world, when they are a common issue among the general population. ICDs significantly interfere with an individual's everyday life. Occupational performance, social engagement, and personal relationships all suffer considerably. This disruption deepens existing vulnerabilities in affected individuals. It also worsens comorbid mental health conditions significantly.

The WHO has designated ICDs as a key area of concern. This is formally acknowledged within the ICD-11 framework (WHO, 2022). Accumulating research consistently associates these disorders with harmful outcomes. Documented consequences span both health and social functioning broadly.

Service outlines several distinct ICD presentations. These include pathological gambling and intermittent explosive disorder (Fariba and Gokarakonda, 2021). Kleptomania and pyromania are also formally recognised categories. Trichotillomania, commonly known as hair-pulling disorder, is equally included. Recent scholarship has broadened how ICDs are conceptually understood. Researchers now incorporate behavioural addictions within expanded frameworks. Compulsive internet use has attracted considerable academic attention recently.

Problematic pornography consumption is similarly being studied more rigorously. Binge-eating behaviour also features prominently in contemporary ICD discourse. The current report provides a detailed academic background of the Impulse control disorders, classification, diagnosis, existing evidence-based treatments and rehabilitation programs, existing clinical and ethical controversies, and recommendations to practice and policy that can be embraced in the United Kingdom. The literature sources used in the study are reputed to be authoritative, including the NHS, the National Institute of Health and Care Excellence (NICE), the WHO, peer-reviewed literature, and offer an integrated and critical perspective.

2. Classification and Diagnosis

Impulse control Disorders have seen a lot of representation in nomenclature as the issues of neurobiological, psychological, and epidemiological studies have evolved over the decades. The DSM-5 formally categorises ICDs under a specific diagnostic grouping. Topically, they are classified as Impulse-Control, Disruptive, and Conduct Disorders. This grouping brings together several behaviourally distinct conditions. Each shares core difficulties in regulating emotions and actions. Oppositional defiant disorder falls within this diagnostic category. Conduct disorder is equally recognised under this classification. Intermittent explosive disorder represents another formally listed condition. Kleptomania and pyromania complete the core diagnostic grouping. Pathological gambling and trichotillomania have been re-classified. Moreover, they are now classified under the chapters of obsessive-compulsive disorders. The disorders are also similar to addictive spectrum disorders. Impulse control disorders are also a diagnosis in the ICD-11 (WHO, 2022). There are three features of diagnostic principles.

To begin with, there is a chronic inability to regulate impulses or urges. Second, building up tension and affective excitement before the behaviour. Third, satisfaction, relief, or discharge during or after the behaviour.

Notably, the behaviour is not explained by some other mental or medical disorder. The ICD-11 has these disorders in a spectrum with the obsessive-compulsive and related disorders, and it is evident that impulsive and compulsive are the core transdiagnostic constructs that are dimensional and overlap.

In the United Kingdom, clinicians delivering services in the NHS tend to work with both ICD-11 and DSM-5, and the NICE (2014) guidelines inform the practice of assessment. Diagnosis typically arrived after a thorough psychiatric examination, structured clinical interviewing and through validated psychometric tests. The measurement of domains of impulsive symptomatology is done using such instruments as the Barratt Impulsiveness Scale (BIS-11) and “Minnesota Impulsive Disorders Interview” (MIDI) (NICE, 2014). The importance of the assessment process, frequency of co-morbidity of ICDs with other severe conditions, such as “attention deficit hyperactivity disorder” (ADHD), bipolar affective disorder, substance use disorders, and borderline personality disorder, are noteworthy (Royal College of Psychiatrists, 2020). Based on the type of disorder and procedure, ICDs are found to affect 1%-13% of the general population according to the epidemiological estimates (NHS, 2023). Pathological gambling, which has been redefined as the Gambling Disorder by the ICD-11 classification, has a prevalence of approximately 0.5-3 percent of adults in the United Kingdom, with young men, people of lower socioeconomic status, and people with other psychiatric disorders being disproportionately represented (Public Health England, 2019). Under-reporting of ICDs remains a significant issue, particularly because of shame, stigma and ignorance of such conditions in society.

3. Treatment and Rehabilitation

The NICE clinical guidelines, NHS commissioned services and an ever-growing amount of evidence in the form of pharmacological, psychological and psychosocial interventions guide the treatment of Impulse control disorders in the United Kingdom. Given the diversity of presentations in the ICD spectrum, no one treatment modality has ever been universally endorsed, but rather, personalised and formulation-based approaches have been preferred in all the clinical guidance documents.

3.1 Psychological Interventions

The most endorsed psychological treatment for different ICDs like pathological gambling, kleptomania and trichotillomania is Cognitive Behavioural Therapy (CBT). CBT is recommended as a first-line intervention by NICE (2014) with certain modifications to its use in terms of cognitive distortions, urge surfing, relapse prevention, and behavioural activation. As an adjunctive treatment, motivational interviewing (MI) is usually used to enhance the desire and readiness to change, especially when used in individuals with low insight or ambivalence towards the symptoms. The other type of therapy that has been known to be effective in treating impulsive behaviours is the mindfulness-based cognitive therapy (MBCT) as it offers them metacognitive insight and control of their feelings. The behaviour therapy, which is evidence-based and that can be used in the treatment of trichotillomania and skin-picking disorders is “Habit Reversal Training” (HRT). NICE (2014) acknowledges that HRT is an effective modality of treatment, which is the combination of training of awareness, competing response and social support. “Acceptance and commitment therapy” (ACT) has also been an appealing third-wave cognitive behavioural therapy, where the manifestation is in the form of experiential avoidance and cognitive fusion.

3.2 Pharmacological Intervention

Treatment of ICDs pharmacologically remains a topic of research and there is no pharmaceutical in the UK licensed to treat such a diagnosis. An opioid receptor antagonist called naltrexone can be used to decrease the desire and frequency of impulsive behaviour in the case of gambling disorder and kleptomania (NHS, 2023). In addition, SSRIs are also most commonly prescribed to treat ICDS of compulsive nature, e.g. trichotillomania and pathological skin picking, as suggested by the NICE on the obsessive-compulsive spectrum disorders. Besides, SSRIs are also generally used to treat ICDS with compulsive characteristics including trichotillomania and pathological skin picking, as recommended by NICE. N-acetylcysteine (NAC) is one of the examples of glutamate modulators that have proven to be initially effective in trichotillomania and gambling disorder. Nevertheless, larger-scale RCTs should be conducted to be able to make any conclusions regarding the clinical recommendations (NHS England, 2021).

3.3 Rehabilitation and Social Support

The ICDs rehabilitation is not only aimed at minimising the symptoms but also to recover the functions, reintegrate an individual back to social life and prevent relapse. The coordination of multidisciplinary care is around NHS community mental health teams (CMHTs) and specialist services, including gambling clinics and addictions services. Public Health England (2019) has highlighted the importance of peer support networks, participation of the voluntary sector and community-based interventions in addressing the broader social determinants that exacerbate ICD presentations. The NHS Long Term Plan (NHS England, 2019) will increase access to specialist psychological interventions and community mental health services to people with complex presentations, including people with comorbid ICDs and personality disorders, substance use, or neurodevelopmental disorders.

4. Recent Discussions and Problems

Impulse control disorders are a field that is marked by a number of major and unresolved controversies, which have significant implications for classification, treatment, and policy implications.

4.1 Impulsiveness vs. Compulsiveness

One of the main and least resolved issues is the correct position of ICDs in the taxonomic framework of mental disorders. Unlike the DSM-5, which positions ICDs as a major impulse regulation disorder, the ICD-11 (WHO, 2022) places many of these presentations on a continuum of obsessive-compulsive processes, which are also dimensions of similarity between impulsive and compulsive processes. The practical implications of this classification are controversial in treatment choice, commissioning of healthcare, and eligibility for psychological therapies using the NICE guidelines. To provide a more consistent and clinically useful nosology, the Royal College of Psychiatrists (2020) has suggested that further studies should focus on the transdiagnostic basis of impulsivity and compulsivity.

4.2 Technological Change Behavioural Addictions.

The sudden rise in the use of digital technologies has spawned a lot of discussion regarding the limits of ICDs and behavioural addictions. In ICD-11, gaming disorder was finally added (WHO, 2022) due to the increasing clinical support of pathological patterns of video game use. In the United Kingdom, NHS England (2021) created the National Problem Gambling Clinic and has gone forward with specialist services to cover technology-related presentations, such as compulsive internet use and social media addiction. Opponents, though, claim that medicalisation of normal digital practices poses a risk of diagnostic inflation and pathologisation of recreational activities, and demand greater epidemiological validation of new categories before they are institutionalised.

4.3 Stigma and Access to Services

Stigma related to ICDs is a significant hindrance to help-seeking behaviour in the United Kingdom. Gambling disorder, kleptomania, and compulsive sexual behaviour are often the presentations of gambling disorder and maliciously labelled as wilful, and this discourages the use of statutory mental health services (SeeMe, ND). Health UK highlights notable inequalities in specialist ICD service access (Mental Health UK, ND). NHS regions vary considerably in the provision of these services. Rural communities and economically deprived areas remain consistently underserved. This gap in provision raises serious equity concerns nationally. The marginalised groups experience an added obstacle to getting the right care. “Black, Asian and minority ethnic background” (BAME) people also face other culturally specific barriers to seeking help. Cultural stigma is the main setback. Language difficulties further restrict meaningful engagement with available services. Deep-rooted institutional mistrust also prevents timely access to treatment.

4.4 Pharmacological Research Gaps

Although ICDs are of clinical importance, the pharmacological evidence base is still disjointed and diffuse. Most pharmacological studies have been typified by limited size, limited time follow-up, and methodological constraints limiting the validity of effect size estimates and externalisation of results (NICE, 2014). The lack of pharmacological agents licensed in the United Kingdom to treat certain ICD presentations poses a prescription difficulty to clinicians, as they are left with off-label prescriptions with little evidence, accompanied by medicolegal and ethical consequences.

5. Recommendations

The following are recommendations that are proposed based on critical synthesis of the evidence and policy landscape to guide clinical practice, service development, and policy on the public health in the United Kingdom: To begin with, NHS England and Integrated Care Boards (ICBs) ought to contract specific specialist ICD pathways in community mental health systems, where geographical coverage of evidence-based assessment and treatment is equal. The existing excessive dependency on generic mental health care with limited specialist training on ICD presentation continues to perpetuate the under-identification and sub-optimal care.

Second, NICE ought to request new clinical guidelines that directly cover the assessment and treatment of ICDs as a different diagnostic category with new evidence available on gaming disorder, compulsive sexual behaviour disorder, and technology-related presentations. Current guidelines are still too general to take into account all presentations of ICD. Third, anti-stigma interventions against ICDs must be incorporated within national mental health promotion agencies and initiatives, with models like the Time to Change. Increasing social and professional knowledge of the neurobiological and psychological basis of ICDs is needed to oppose the moral narratives that demoralise help-seeking and undermine recovery.

Fourth, the funding organisations such as the National Institute of Health Research (NIHR), ought to focus on high-quality RCTs studying the effectiveness of pharmacological and integrated treatment methods of certain ICD presentations, especially underrepresented groups and long-term follow-up results. Fifth, NHS clinician training and ongoing professional development must include NHS-specific content that will enable practitioners working within

primary care, community mental health, and specialist services to identify, assess, and refer people with ICD presentations in a non-stigmatising way and in a timely fashion.

5.1 Conclusion

Impulse Control Disorders are an underdiagnosed group of clinically heterogeneous psychiatric disorders with a high individual, social, and economic burden. This report has followed the history of the ICD classification system, from the early categorical methods to the modern dimensional systems, codified in ICD-11 (WHO, 2022) and DSM-5, and has discussed the evidence base of psychological, pharmacological, and psychosocial treatment methods in the UK. Major gaps still exist in the area of specialist services, the establishment of a sound pharmacological evidence base, and the minimization of stigma that continues to hamper access to care. With the ongoing transformation of behaviour patterns and risk by digital technologies, and with the widening of the behavioural addictions taxonomy, the sphere of ICDs has to overcome complicated conceptual, ethical, and policy issues. The guidelines in this report indicate adherence to evidence-based, fair, and recovery-focused care among people with ICDs in the National Health Service and in the broader UK mental health system. It requires a long-term commitment to research, clinical training, and service development to help fulfil the unmet demand that is presently defining this valuable field of mental health practice.

References

Mental Health UK. (ND). *Obsessive Compulsive Disorder (OCD)*.

Available at: <https://mentalhealth-uk.org/help-and-information/conditions/obsessive-compulsive-disorder/> (Accessed: 19 April 2026).

SeeMe. (ND). *Experiencing Stigma and Discrimination*. Available at:

<https://www.seemescotland.org/stigma-discrimination/experiencing-stigma-and-discrimination> (Accessed: 19 April 2026).

Fariba, K. and Gokarakonda, S.B. (2021). *Impulse Control Disorders*. [online] PubMed.

Available at: <https://www.ncbi.nlm.nih.gov/books/NBK562279/>. (Accessed: 20 April 2026).

National Institute for Health and Care Excellence. (2014). *Obsessive-compulsive disorder and body dysmorphic disorder: Treatment (Clinical Guideline CG31)*. NICE.

Available at: <https://www.nice.org.uk/guidance/cg31> (Accessed: 17 April 2026).

NHS England. (2019). *The NHS long term plan*. Available

at: <https://webarchive.nationalarchives.gov.uk/ukgwa/20250506042016/https://www.longtermplan.nhs.uk/publication/nhs-long-term-plan/> (Accessed 14 April 2026).

NHS England. (2021). Adult mental health. Available at:

<https://www.england.nhs.uk/mental-health/adults/> (Accessed: 20 April 2026).

Public Health England. (2023). *Gambling-related harms evidence reviews: Summary*.

Available at: <https://www.gov.uk/government/publications/gambling-related-harms-evidence-review/gambling-related-harms-evidence-review-summary--2> (Accessed at: 14 April 2026).

Royal College of Psychiatrists. (2020). *Services for people diagnosed with personality*

disorder (PS01/20). Available at: https://www.rcpsych.ac.uk/docs/default-source/improving-care/better-mh-policy/position-statements/ps01_20.pdf?sfvrsn=85af7fbc_4 (Accessed: 20 April 2026).

Science Direct (2020) *Impulsive–compulsive problems in young people*. Available at:

<https://www.sciencedirect.com/science/article/pii/S1357303920302000> (accessed: 15 April 2026).

World Health Organization. (2022). *International classification of diseases, eleventh revision*

(ICD-11). <https://www.who.int/standards/classifications/classification-of-diseases> (Accessed: 20 April 2026).