

To explore whether exercise duration and the type of exercise has an impact on individual well-being and to provide a more in-depth insight to why this might be.

Charlotte Evans

2206694

Bachelor of the Arts (hons) in Applied Psychology.



Prifysgol Cymru
Y Drindod Dewi Sant
University of Wales
Trinity Saint David

Background literature



Aerobic vs Anaerobic (Wahid et al., 2016; Patel et al., 2017).



2.5 hours of moderate exercise per week (World Health Organisation, 2024).



80% of adolescents and 27% of adults don't meet the recommended timeframe (World Health Organisation, 2024).



Herazo-Beltran et al. (2017) study on barriers faced within exercise.

Studies

- COVID-19 led to reduced physical activity and a rise in depression and antidepressant use.
- Antidepressant prescriptions increased (Gray, 2022).
- Regular exercise promotes neurotransmitters like endorphins – supports Endorphin Theory (Craft & Perna, 2004).
- Enhances social interaction and self-identity (Hovland et al., 2023).
- Longer exercise duration (Basso et al., 2022).
- Dance and yoga groups showed greater well-being and community compared to control (Welford et al., 2022).
- Motivation differs: men are more intrinsically motivated, women extrinsically (Granero-Jiménez et al., 2022).
- Men prefer individual workouts; women prefer group settings for social support (Agius et al., 2024).





Rationale

- Exercise improves mental well-being
- Research gap: Effects of exercise type and duration underexplored
- Previous studies: small samples, limited methods
- This study uses a mixed-methods approach.
- Warwick-Edinburgh Mental Well-being Scale (Tennant et al., 2007)
- Open-ended questions for personal experiences
- Aims to deepen understanding of exercise's impact on well-being.

Hypotheses



The participants in the 150+ minutes group will have a higher score on the well-being score compared to the other groups.



The aerobic group will have higher scores on the well-being scale compared to the other groups.



The duration of aerobic exercise will have a positive effect on well-being, such that longer durations of aerobic exercise will lead to greater improvements in well-being scores.

Methodology

53 Participants

Aerobic group= 35 and Anaerobic group=18.

150+ group

60–150-minute group

less than 60-minute group

‘no exercise’ group

53= Quantitative 50=Qualitative

Mixed-method, web-based design

Quantitative - Measured exercise type, duration, and well-being


Used Warwick-Edinburgh Mental Well-Being Scale - Analyzed with 2x4 ANOVA (SPSS)

Qualitative - 5 open-ended questions - Thematic analysis: 67 codes - 4 themes


Materials-

Warwick Edinburgh Well-being Scale (Tennant et al., 2007).


Open-ended questionnaire used for qualitative data.



Procedure



Quantitative Results



Well-being Scores and duration

Highest: 60-150 minutes/week

Lowest: No exercise

Well-being scores and type

The type of exercise (aerobic vs. anaerobic) was not significant.

Interaction Between Exercise Type & Duration

Highest Well-being Score = *Anaerobic exercise (60–150 mins/week)*

Lowest Well-being Score = *Anaerobic / No Exercise*

Duration < impact > type of exercise.

Qualitative results

- Self-perception and mental well-being.
- Barriers to exercise.
- Personal preference in exercise.
- Physical health and fitness goals.



Discussion of the Quantitative results



Hypothesis 1 – Longer duration increased well-being scores – Supported Iwon et al. (2021).



Hypothesis 2 – Aerobic exercise would have higher well-being scores – Not supported Welford et al. (2022).




Hypothesis 3 – The 150+ minutes/week of aerobic exercise would have higher well-being scores – Not supported Basso et al. (2022).

Discussion of Qualitative results

- **Mental benefits** – Endorphin hypothesis and Cardiovascular Adaptation Theory (Craft & Perna, 2004; Vega et al., 2017).
- **Fitness goals** -muscle building or weight loss - boosted participants' self-esteem and motivation. (Iwon et al., 2021).
- **Personal preference** - group / solo training (Agius et al., 2024).
- **Barriers in exercise** (Herazo-Beltrán et al., 2017).
- **Implications** - Exercise plans – Reduce reliance on Anti-depressants.



Strengths



Used a mixed method – enhance validity and depth of findings.

Variation in primary exercise and duration/week
- compare differences – range of perspectives

Limitations and future research

- Equal participants in each group.
- Wording of question 11.
- Focus on age of participants





Conclusion

- Longer duration = increased well-being
- Type of exercise = not significant
- Benefits and barriers
- Prevention of exercise
- Acquiring a gym partner

Reference list

- Agius, G., Calleja, P., & Muscat-Inglott, M. (2024). Examining Individual Versus Group Exercise Preference Among Maltese Gym-Goers: Journal of Sports and Physical Education Studies. <https://doi.org/10.32996/jspes.2024.4.1.2>
- Basso, J. C., Oberlin, D. J., Satyal, M. K., O'Brien, C. E., Crosta, C., Psaras, Z., Metpally, A., & Suzuki, W. A. (2022). Examining the Effect of Increased Aerobic Exercise in Moderately Fit Adults on Psychological State and Cognitive Function. Frontiers in Human Neuroscience. <https://doi.org/10.3389/fnhum.2022.833149>
- Granero-Jiménez, J., López-Rodríguez, M. M., Dobarrío-Sanz, I., & Cortés-Rodríguez, A. E. (2022). Influence of Physical Exercise on Psychological Well-Being of Young Adults: A Quantitative Study. International Journal of Environmental Research and Public Health. <https://doi.org/10.3390/ijerph19074282>
- Gray, K. (2022, July 7). Medicines Used in Mental Health – England – 2015/16 to 2021/22 | NHSBSA. Wwww.nhsbsa.nhs.uk. <https://www.nhsbsa.nhs.uk/statistical-collections/medicines-used-mental-health-england/medicines-used-mental-health-england-201516-202122>
- Herazo-Beltrán, Y., Pinillos, Y., Vidarte, J., Crissien, E., Suarez, D., & García, R. (2017). Predictors of Perceived Barriers to Physical Activity in the General Adult Population: A CrossSectional Study. Brazilian Journal of Physical Therapy. <https://doi.org/10.1016/j.bjpt.2016.04.003>
- Hovland, J., Langeland, E., Ness, O., & Bente Ovedie Skogvang. (2023). Experiences with Physical Activity, Health and Well-being Among Young Adults with Serious Mental Illness. International Journal of Qualitative Studies on Health and Well-Being. <https://doi.org/10.1080/17482631.2023.2221911>
- Iwon, K., Skibinska, J., Jasielska, D., & Kalwarczyk, S. (2021, December 17). Elevating Subjective Well-being Through Physical Exercises: An Intervention Study. Frontiers. <https://www.frontiersin.org/journals/psychology/articles/10.3389/fpsyg.2021.702678/full#h3>
- Mahindru, A., Patil, P., & Agrawal, V. (2023). Role of Physical Activity on Mental Health and Well-Being: A Review. Cureus. <https://doi.org/10.7759/cureus.33475>
- Patel, H., Alkhawam, H., Madanieh, R., Shah, N., Kosmas, C. E., & Vittorio, T. J. (2017). Aerobic vs Anaerobic Exercise Training Affects the Cardiovascular System. World Journal of Cardiology. <https://doi.org/10.4330/wjc.v9.i2.134>
- Tennant, R., Hiller, L., Fishwick, R., Platt, S., Joseph, S., Weich, S., Parkinson, J., Secker, J., & Stewart-Brown, S. (2007). Warwick-Edinburgh Mental Well-Being Scale (WEMWBS) [Database record]. APA PsycTests. <https://doi.org/10.1037/t01628-000>
- Thygesen, L. C., Møller, S. P., Ersbøll, A. K., Santini, Z. I., Dahl Nielsen, M. B., Grønbæk, M. K., & Ekholm, O. (2021). Decreasing Mental Well-being During the Covid-19 Pandemic: A Longitudinal Study Among Danes Before and During the Pandemic. Journal of Psychiatric Research. <https://doi.org/10.1016/j.jpsychires.2021.09.035>
- Vega, R. B., Konhilas, J. P., Kelly, D. P., & Leinwand, L. A. (2017). Molecular Mechanisms Underlying Cardiac Adaptation to Exercise. Cell Metabolism. <https://doi.org/10.1016/j.cmet.2017.04.025>
- Welford, P., Östh, J., Hoy, S., Diwan, V., & Hallgren, M. (2022). Effects of Yoga and Aerobic Exercise on Well-being in Physically Inactive Older Adults: Randomized Controlled Trial. Complementary Therapies in Medicine. <https://doi.org/10.1016/j.ctim.2022.102815>
- Wicker, P., & Frick, B. (2015). The Relationship Between Intensity and Duration of Physical Activity and Subjective Well-being. The European Journal of Public Health. <https://doi.org/10.1093/eurpub/ckv131>
- World Health Organization. (2024, June 26). Physical Activity. World Health Organization. <https://www.who.int/news-room/fact-sheets/detail/physical-activity>

Basic Acknowledgement format of AI tool use

I acknowledge the use of Grammarly <https://www.grammarly.com/> on May 17th, 2025, to check my work by using the following prompt. I used the prompt(s):

1. Please check the following text for grammar.

I confirm that I have not presented any AI generated text as my own work. I confirm I have copies of my drafts, notes, and other resources I used before, and after the use of AI tools which I may be asked to provide in evidence.