

Introduction

Diabetes mellitus is a long-term chronic illness that happens when the body is unable to regulate blood glucose blood sugar levels effectively. Glucose is the body's major source of energy and originates from the food we eat. The hormone insulin, which is produced by the pancreas, helps transfer glucose from the bloodstream into the body's cells where it is used for energy. When the body does not create enough insulin or cannot use insulin efficiently, glucose builds up in the blood, resulting in high blood sugar levels (hyperglycaemia). If diabetes is not managed effectively, it can lead to major health concerns affecting the heart, kidneys, eyes, nerves and blood vessels

UK statistics(Prevalence)-6 million people in the UK are living with diabetes, including over 4.7 million diagnosed cases and approximately 1.3 million undiagnosed cases (Diabetes UK, 2025).

The Role of Insulin in Blood Sugar Control

Insulin is a hormone generated by the beta cells of the pancreas. After eating, carbohydrates are broken down into glucose, causing blood sugar levels to rise. In reaction, the pancreas produces insulin into the bloodstream. Insulin causes glucose to flow from the blood into body cells, where it can be used for energy. Excess glucose is stored in the liver and muscles as glycogen for later use. This mechanism decreases blood glucose levels back to a safe range.

The Role of Glucagon in Maintaining Blood Sugar Levels

Glucagon is another hormone generated by the pancreas that functions alongside insulin. When blood glucose levels fall too low, such as between meals or during physical exertion, the pancreas produces glucagon. Glucagon stimulates the liver to transform stored glycogen back into glucose and release it into the bloodstream. This improves blood sugar levels and avoids hypoglycaemia for example abnormally low blood glucose levels.

Blood Sugar Regulation in People with Diabetes

In people with diabetes, the normal regulation of blood glucose is impaired. In Type 1 diabetes, the pancreas produces little or no insulin, so insulin must be replaced through injections or other delivery methods. In Type 2 diabetes, the body may become resistant to insulin or produce insufficient insulin, causing glucose to remain in the bloodstream.

To maintain safe and effective blood glucose levels, people with diabetes are encouraged to monitor their blood sugar, follow their personalised treatment plan, maintain a balanced diet, participate in regular physical activity and attend healthcare appointments. Effective management helps reduce the risk of short-term problems, such as hypoglycaemia and

hyperglycaemia, and long-term complications including cardiovascular disease, kidney damage, nerve damage and vision problems.


How Oxygen and Glucose Are Used to Produce Energy Through Cell Metabolism

Cellular Respiration

Cellular respiration is a metabolic process that occurs inside cells, mainly within the mitochondria. It allows the body to convert the chemical energy stored in glucose into adenosine triphosphate (ATP), which is the usable energy source needed for activities such as muscle contraction, movement, growth, repair and maintaining body functions.

During cellular respiration, glucose is broken down using oxygen to release energy. The waste products produced are carbon dioxide and water, which are removed from the body through breathing and other processes.

Diagram: Aerobic Cellular Respiration



Link to Diabetes

In diabetes, the problem is not usually that glucose is unavailable, but that glucose cannot enter cells effectively because:

- The body does not produce enough insulin (Type 1 diabetes).
- Cells become resistant to insulin (Type 2 diabetes).

As a result, glucose remains in the bloodstream, causing high blood sugar levels, while body cells may not receive enough glucose efficiently for energy production. This explains why controlling blood glucose levels is important for maintaining normal cell metabolism and preventing complications.

How Recording and Tracking Blood Sugar Levels Helps Individuals with Type 2 Diabetes Safely Manage Their Condition

Blood Glucose Patterns

Recording and tracking blood sugar levels helps individuals with Type 2 diabetes understand how their glucose levels change over time. Blood glucose levels are affected by many factors, including food intake, physical activity, stress, illness and medication. By keeping a record of readings, patients can identify patterns, such as whether their blood sugar increases after certain meals or decreases after exercise. This knowledge allows individuals to make informed choices about their lifestyle and diabetes management.

Effective Diabetes Management

Regular monitoring allows individuals to check whether their diabetes treatment plan is controlling their blood glucose effectively. Type 2 diabetes occurs when the body becomes resistant to insulin or does not produce enough insulin, causing glucose to build up in the bloodstream. Tracking blood sugar levels provides information that can help patients and healthcare professionals evaluate whether lifestyle changes, medication or other treatments are working effectively.

Preventing Hyperglycaemia (High Blood Sugar)

Recording blood glucose levels can help identify periods when blood sugar is consistently too high. Long-term hyperglycaemia can damage blood vessels and increase the risk of complications such as cardiovascular disease, kidney disease, nerve damage and vision problems. By recognising high blood sugar patterns early, individuals can take action, such as adjusting their diet, increasing physical activity or discussing medication changes with their healthcare professional.

Reducing the Risk of Hypoglycaemia (Low Blood Sugar)

Although hypoglycaemia is more commonly associated with insulin treatment or certain diabetes medications, some individuals with Type 2 diabetes may experience low blood sugar levels. Monitoring helps patients recognise symptoms and identify situations that may cause low blood glucose, such as skipping meals, increased exercise or medication effects. Early recognition allows appropriate action to be taken to restore blood glucose levels safely.

Healthcare Professionals Make Informed Decisions

Accurate blood glucose records provide valuable information during diabetes reviews and appointments. Healthcare professionals, including practice nurses, can use these records alongside other measurements, such as HbA1c results, to assess overall diabetes control. This information supports decisions about medication adjustments, lifestyle advice and personalised care plans.

Patient Engagement and Self-Management

Tracking blood glucose encourages individuals to take an active role in managing their condition. Understanding the relationship between lifestyle choices and blood sugar levels can increase confidence and motivation. For example, patients may observe that regular physical activity improves their readings or that reducing high-sugar foods helps maintain better control. This supports shared decision-making between patients and healthcare professionals.

Improving Long-Term Health Outcomes

Consistent monitoring and recording of blood glucose levels can help individuals maintain safer glucose control and reduce the risk of diabetes complications. It supports early identification of problems, encourages adherence to treatment plans and enables patients to work effectively with healthcare professionals.

Link to the Case Study

In the case study, the practice nurse identified that many patients diagnosed with diabetes would benefit from increased understanding of their condition. Teaching patients the importance of recording and tracking blood sugar levels can empower them to manage their Type 2 diabetes more effectively. Improved knowledge allows patients to make safer decisions about medication, diet and physical activity, supporting better blood glucose control and improving their quality of life.

Type 2 Diabetes Management

1. Adopt a Healthy, Balanced Diet to Improve Blood Glucose Control

A healthy diet is an important part of managing Type 2 diabetes because the food a person eats directly affects blood glucose levels. Individuals should be encouraged to follow a balanced diet that includes high-fibre carbohydrates, vegetables, fruits, lean proteins and healthy fats while reducing the intake of foods high in added sugars, saturated fats and processed carbohydrates.

High-fibre foods, such as whole grains, beans, vegetables and pulses, are digested more slowly, which can help prevent rapid increases in blood glucose after meals. Maintaining appropriate portion sizes and choosing healthier carbohydrate sources can support better glucose control and help with achieving or maintaining a healthy weight.

2. Increase Regular Physical Activity to Improve Insulin Sensitivity

Regular physical activity is recommended for people with Type 2 diabetes because exercise helps the body use glucose more effectively. During physical activity, muscles require more energy and take up more glucose from the bloodstream, which can help lower blood sugar levels. Exercise also improves insulin sensitivity, meaning the body's cells respond better to insulin.

Individuals should be encouraged to include regular aerobic activities, such as walking, cycling or swimming, alongside strength exercises where appropriate. Increasing daily

movement and reducing long periods of sitting can contribute to improved blood glucose control, weight management and cardiovascular health.

Evidence-based guidance recommends that adults with Type 2 diabetes aim to include regular physical activity as part of their diabetes management plan. Exercise can also improve mood, energy levels and overall quality of life.

3. Stop Smoking to Reduce Diabetes-Related Health Risks

Smoking cessation is an important lifestyle change for people with Type 2 diabetes because smoking increases the risk of serious complications. Smoking can contribute to increased insulin resistance, making it more difficult for the body to control blood glucose levels. It also damages blood vessels and increases the risk of cardiovascular complications, which are already higher in individuals with diabetes.

Basically, Supporting patients to stop smoking through behavioural support, smoking cessation services and appropriate treatments can improve overall health outcomes. Reducing exposure to tobacco smoke lowers the risk of heart disease, stroke and circulation problems associated with diabetes.

Link to the Case Study

In the case study, the practice nurse identified that many patients would benefit from greater understanding of how lifestyle factors influence their diabetes management. Providing evidence-based education about diet, physical activity and smoking cessation can help patients make informed decisions and actively participate in their care plans. These lifestyle changes, combined with blood glucose monitoring and medical treatment where required, can improve blood sugar control and reduce the risk of long-term diabetes complications

Conclusion

In my opinion, In order to maintain acceptable blood glucose levels and lower the risk of major health problems, diabetes is a complicated long-term condition that needs to be effectively managed. The case study emphasises the growing number of primary care patients receiving diabetes diagnoses and the significance of enhancing their comprehension of the illness. In order to assist people understand how diabetes impacts the body, how glucose is used for energy through cellular metabolism, and why maintaining stable blood sugar levels is vital for general health, practicing nurses must give patient education.

Overall, practice nurses play a critical role in empowering patients through monitoring, education, and continuous support. Healthcare providers can assist people with Type 2 diabetes in acquiring the knowledge and abilities necessary to safely and successfully manage their condition by giving them accurate information on blood glucose control and lifestyle choices. Effective diabetes self-management can lessen the burden of diabetic complications on people and healthcare providers, enhance quality of life, and encourage improved health outcomes.

References

American Diabetes Association (2024) *Standards of Care in Diabetes—2024*. Diabetes Care, 47(Supplement 1), pp. S1–S321.

Diabetes UK (2024) *Diabetes and healthy lifestyle management*. Available at: <https://www.diabetes.org.uk>

National Institute for Health and Care Excellence (NICE) (2022) *Type 2 diabetes in adults: management (NG28)*. Available at: <https://www.nice.org.uk/guidance/ng28>

National Health Service (NHS) (2024) *Type 2 diabetes: causes, symptoms and treatment*. Available at: <https://www.nhs.uk/conditions/type-2-diabetes/>

World Health Organization (WHO) (2023) *Diabetes*. Available at: <https://www.who.int/health-topics/diabetes>

International Diabetes Federation (IDF) (2021) *IDF Diabetes Atlas*. 10th edn. Brussels: International Diabetes Federation.

Davies, M.J., Aroda, V.R., Collins, B.S. et al. (2022) 'Management of hyperglycaemia in type 2 diabetes, 2022: a consensus report', *Diabetologia*, 65, pp. 1925–1966.

Lean, M.E.J., Leslie, W.S., Barnes, A.C. et al. (2018) 'Primary care-led weight management for remission of type 2 diabetes', *The Lancet*, 391(10120), pp. 541–551.

Colberg, S.R., Sigal, R.J., Yardley, J.E. et al. (2016) 'Physical activity/exercise and diabetes: a position statement of the American Diabetes Association', *Diabetes Care*, 39(11), pp. 2065–2079.

Pan, A., Wang, Y., Talaei, M., Hu, F.B. and Wu, T. (2015) 'Relation of smoking with risk of type 2 diabetes: a meta-analysis of prospective cohort studies', *The Lancet Diabetes & Endocrinology*, 3(12), pp. 958–967.

Evert, A.B., Dennison, M., Gardner, C.D. et al. (2019) 'Nutrition therapy for adults with diabetes or prediabetes: a consensus report', *Diabetes Care*, 42(5), pp. 731–754.

Diabetes UK (2025) How many people in the UK have diabetes? Available at: <https://www.diabetes.org.uk>