



[23RD OF FEBRUARY]

[LCHS4003: INTRODUCTION TO HEALTH AND SOCIAL CARE]

[PORTOFOLIO: TWO TASKS (POSTER & ESSAY)]

SAMIRA-FLORENTINA SERBANA


2514813

DANIEL HARUNA



A Brief History: From Asylums to the Community

The development of community mental health must be understood in the context of the historical shift from institutionalisation to deinstitutionalisation. This transition was driven by changes in medical knowledge, human rights advocacy, and economic policy, and represented a fundamental rethinking of ethical mental healthcare (Shorter, 1997; Thornicroft & Tansella, 2004).



Critical Challenges Facing Community Mental Health Services

Despite progress, serious challenges remain. One major issue is the “severity gap,” where individuals are deemed too unwell for primary care but not unwell enough for specialist services (NHS England, 2019). This often results in delayed care until crisis point, creating a “revolving door” pattern of admission and discharge (Appleby et al., 2017; Thornicroft & Tansella, 2004).

A Brief History: From Asylums to the Community

The development of community mental health must be understood in the context of the historical shift from institutionalisation to deinstitutionalisation. This transition was driven by changes in medical knowledge, human rights advocacy, and economic policy, and represented a fundamental rethinking of ethical mental healthcare (Shorter, 1997; Thornicroft & Tansella, 2004).


The Era of the Asylum (1800s–1950s)

During the 19th and early 20th centuries, asylums were the dominant model of care for people with mental illness. Initially, these institutions were intended to provide “moral treatment” in calm, structured environments. However, by the early 20th century, many had become overcrowded and custodial, offering little therapeutic benefit (Scully, 2015).

Modern Functions of Community Mental Health Services (2026)

In 2026, community mental health services operate as hubs for long-term recovery, aiming to stabilise individuals within their own environments (Thornicroft et al., 2011). According to the WHO (2021), modern CMH services are structured around four core pillars.

At the core of community mental health is the philosophy of recovery. Recovery emphasises hope, personal agency, and the belief that people with severe or enduring mental health conditions can still achieve fulfilling lives when adequately supported (Anthony, 1993). CMH services typically rely on multidisciplinary teams that combine clinical interventions with social, physical, and practical support. This holistic approach ensures that treatment addresses the “whole person,” including their relationships, housing, employment, and physical health, rather than focusing narrowly on diagnosis or symptoms (Barker, 2009).



Deinstitutionalisation and Its Challenges (1970s–1990s)

While deinstitutionalisation reduced reliance on hospitals, it also exposed major weaknesses in service provision. Many community services were underfunded or never fully developed. This led to “transinstitutionalisation,” where individuals moved from hospitals into prisons, nursing homes, or homelessness rather than receiving effective community care (Lamb & Bachrach, 2001; Torrey, 1997).



The Evolution and Impact of Community Antenatal Midwifery Services

The maternity care landscape has evolved dramatically over the last century, shifting from a home-based, woman-centered foundation to a highly clinical institutionalisation before returning to a community-based model (Oakley, 1984; Downe, 2015). Community antenatal midwifery services are a crucial component of modern healthcare, emphasising accessibility, continuity of care, and the general health of the mother and child (Sandall et al., 2016).

Origins and Historical Context

The history of community midwifery extends back to the dawn of society. In the past, the "handywoman" or lay midwife provided help in the home and was a mainstay of the neighbourhood (Towler and Bramall, 1986). However, the formalisation of these services began in the late 19th and early 20th centuries. The passage of the Midwives Act in 1902 marked a major turning point in the UK, requiring formal training and establishing midwifery a protected profession (UK Parliament, 1902).

How these services were developed was significantly influenced by the sociopolitical environment of the post-war period. There was a push for hospital births when the National Health Service (NHS) was established in 1948. This was predicated on the Peel Report (1970), which claimed that hospital delivery was safer (Department of Health, 1970). As a result, prenatal care became more standardised and clinically focused, leading to the medicalisation of pregnancy (Scamell, 2014).

But by the 1990s, a movement against it had started. The important Changing Childbirth (1993) report called for a return to woman-centered care, saying that pregnancy is a normal part of life and not a medical condition (Department of Health, 1993). This change in policy set the stage for the modern community-based antenatal framework that is used today.

Critical Evaluation of Modern Functions

Community prenatal services are currently the primary point of contact for expectant mothers in the NHS (NHS England, 2016). These services are typically offered through home visits, general practitioner offices, or community clinics. A crucial element of the modern model is the Continuity of Carer approach, which aims to ensure that women receive support from the same midwife or small team during pregnancy, delivery, and the postnatal phase (Sandall et al., 2016).

When adequately resourced, community antenatal midwifery enables:

Early risk screening, which includes spotting issues such gestational diabetes and pre-eclampsia (NICE, 2021).

Psychosocial support, which enables midwives to spot domestic violence, mental health issues, and safeguarding concerns that would not be mentioned in hospital settings (RCM, 2018).

But there are still major obstacles. Rising caseloads and nationwide staffing shortages frequently disrupt continuity by limiting appointments to task-focused, "tick-box" interactions rather than comprehensive evaluations (RCM, 2022). Additionally, despite its inclusive goals, socioeconomic disadvantage, digital exclusion, and language barriers may prevent vulnerable populations from accessing community prenatal care (Marmot et al., 2020).

Enhancing Quality of Life

It is commonly known that community-based prenatal care improves quality of life. Decentralising services improves equity of access to maternity care by cutting down on travel expenses and time (NHS England, 2016).

According to Downe et al. (2018), community care promotes psychological autonomy and empowerment during pregnancy, a time that is frequently linked to vulnerability and anxiety. According to research, women who receive midwife-led continuity of care are more satisfied with their treatment, have fewer preterm births, and need less intervention (Sandall et al., 2016).

Conclusion

The profession's ability to recover maternity care from over-medicalization is demonstrated by community antenatal midwifery programs. The return to community-based, midwife-led models is crucial for comprehensive maternal and newborn health, despite the fact that past policy changes forced births into hospital settings (WHO, 2016). The ongoing challenge lies in securing sustainable funding and staffing to ensure that all individuals can access personalized, respectful, and continuous care.

Reference List (Harvard)

- Department of Health (1970) *Domiciliary and Maternity Bed Needs (Peel Report)*. London: HMSO.
- Department of Health (1993) *Changing Childbirth*. London: HMSO.
- Downe, S. (2015) *Normal Childbirth: Evidence and Debate*. 2nd edn. London: Elsevier.

- Downe, S. et al. (2018) 'The impact of continuity of midwifery care', *Midwifery*, 62, pp. 47–55.
- Marmot, M. et al. (2020) *Health Equity in England: The Marmot Review 10 Years On*. London: IHE.
- NICE (2021) *Antenatal Care (NG201)*. London: NICE.
- Oakley, A. (1984) *The Captured Womb*. Oxford: Blackwell.
- RCM (2018) *Maternal Emotional Wellbeing*. London: Royal College of Midwives.
- RCM (2022) *State of Maternity Services Report*. London: RCM.
- Sandall, J. et al. (2016) 'Midwife-led continuity models', *Cochrane Database of Systematic Reviews*, CD004667.
- Towler, J. and Bramall, J. (1986) *Midwives in History*. London: Croom Helm.
- UK Parliament (1902) *Midwives Act 1902*. London: HMSO.
- WHO (2016) *Standards for Improving Quality of Maternal and Newborn Care*. Geneva: WHO.