

Name: Shazia Yasin

Student Number: 2424946

Module: Introduction to Physiology

Module Code: BMHP5023

Internal Verifier: Chloe Harries

Submission Date: 23rd February 2026

Case Study Report: Pathophysiology and Clinical Management of Type 2 Diabetes

1. Introduction: The Clinical Challenge of T2DM (PART A)

As a practice nurse at a local GP surgery, I am seeing a worrying rise in patients with Type 2 Diabetes Mellitus (T2DM). T2DM is a long-term health condition where blood sugar levels remain too high. It happens because the body's system for managing sugar breaks down. Usually, the body becomes resistant to insulin, and the pancreas, the organ that makes insulin cannot keep up with the demand. While Type 1 diabetes is caused by the body attacking itself, T2DM is a metabolic issue where the body stops responding correctly to its own hormones. This increase in cases is a major crisis for public health, putting a lot of pressure on local doctors and the NHS. In our own clinic, the number of people on the diabetes register has jumped significantly, which shows we need better ways to teach and support our patients.

Epidemiological Context and Risk Factors: The rise in cases at our surgery is part of a global problem. Several factors, some we can change and some we cannot, combine to cause this:

The Environment and Body Fat: Having too much body fat, especially around the stomach, is a major factor. Fat is not just stored energy; it acts like a busy chemical factory. It releases substances that cause inflammation, such as TNF alpha and IL-6. These chemicals block the "signal" that insulin sends to your cells. This means the "locks" on your cells won't open to let sugar in, leading to chronic inflammation and insulin resistance.

Genetics: While lifestyle is often the trigger, some people are born with genes that make them more likely to get T2DM. This "thrifty genotype" helped our ancestors survive when food was scarce, but today, when high-calorie food is everywhere, it leads to health problems. Family history is the strongest risk factor we cannot change.

Socio-Economic Factors: A person's health is often linked to their income and where they live. Many of our patients live in areas where it is hard to find fresh, healthy food. Instead, they rely on cheap, processed foods that are full of sugar and bad fats. As a nurse, I must understand these struggles to create a care plan that actually works for the patient's real life. Also, the stress of money troubles can raise cortisol levels, a hormone that makes it even harder for the body to control blood sugar.

2. Homeostatic Maintenance of Blood Glucose (PART B)

Homeostasis is the physiological cornerstone of human health, representing the body's ability to maintain a stable internal environment despite external fluctuations. For the human body to function correctly, blood glucose must be kept within a narrow physiological range (typically to 4.0 to 7.0 mmol/L post-meal). Maintaining this balance is vital because the central nervous system, particularly the brain, relies almost exclusively on a constant supply of glucose, as it cannot store its own energy or effectively metabolise fatty acids. The brain consumes approximately 20% of the body's total glucose, and even transient drops in blood sugar can lead to impaired cognitive function, confusion, or more severe neurological consequences.

The Pancreatic Control Centre and the Negative Feedback Loop: The Islets of Langerhans within the pancreas act as the primary sensor and control centre for glucose regulation through a sophisticated negative feedback loop:




Figure 1: The homeostatic regulation of blood glucose via the antagonistic action of insulin and glucagon.

The Hyperglycaemic Response (Insulin): Following a meal, blood glucose levels rise as carbohydrates are broken down into monosaccharides. Beta cells in the pancreas detect this increase and secrete Insulin. Insulin is an anabolic hormone that facilitates the movement of glucose from the blood into the cells. Once in the bloodstream, insulin binds to specific receptors on target cells, primarily in skeletal muscle and adipose tissue. This binding initiates a signalling cascade that results in the translocation of GLUT4 glucose transporters from intracellular vesicles to the plasma membrane. This "opens the channel" for glucose to exit the blood. In the liver, insulin promotes glycogenesis, where glucose is synthesized into glycogen for future use.

The Hypoglycaemic Response (Glucagon): Conversely, during periods of fasting, sleep, or intense physical exertion, blood glucose levels drop. Alpha cells respond by secreting Glucagon. This catabolic hormone signals the liver to initiate glycogenolysis (the breakdown of stored glycogen into glucose) and gluconeogenesis (the synthesis of glucose from non-carbohydrate sources like amino acids and glycerol). This prevents the onset of hypoglycaemia, which could otherwise lead to loss of consciousness or neurological

damage. This delicate hormonal dance ensures that every cell in the body receives the fuel it needs while preventing the toxic effects of excessive circulating sugar.

3. Cellular Metabolism: Glucose, Oxygen, and ATP (PART C)

A major component of the education provided in our clinic is explaining the "why" behind the clinical symptoms of T2DM. Many patients present with profound, heavy fatigue. To understand this, we must examine the biochemistry of Aerobic Cellular Respiration, the process by which cells generate Adenosine Triphosphate (ATP), the universal energy currency.

The Biochemical Stages of Energy Production: For a cell to produce energy, it requires both glucose (the fuel) and oxygen (the oxidizing agent). This process occurs in three main stages:

1. **Glycolysis:** This occurs in the cytosol of the cell. One glucose molecule is cleaved into two molecules of pyruvate. This process is anaerobic and yields a net gain of 2 ATP and two electron carriers (NADH).
2. **The Krebs Cycle (Citric Acid Cycle):** Pyruvate is transported into the Mitochondria, often called the "powerhouse" of the cell. Through a series of redox reactions, the carbon bonds are broken, releasing CO₂ as a waste product and generating high-energy electron carriers (NADH and FADH₂).
3. **The Electron Transport Chain (ETC):** Located on the inner mitochondrial membrane, the ETC uses the electrons from NADH and FADH₂ to drive a proton pump. Oxygen acts as the final electron acceptor, combining with hydrogen protons to form water (H₂O). This process, known as oxidative phosphorylation, yields the bulk of the cell's energy, approximately 32-34 ATP molecules.




Figure 2: The biochemical pathway of aerobic respiration, showing the production of ATP within the mitochondria.

The complete chemical reaction for this process is: $\text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2 \rightarrow 6\text{CO}_2 + 6\text{H}_2\text{O} + 36\text{-}38 \text{ ATP}$

The "Diabetic Disconnect" and Fatigue: In a patient with T2DM, the insulin resistance "locks" the cell. Despite the bloodstream being saturated with glucose (hyperglycaemia), the glucose cannot be transported into the cell. Consequently, the mitochondria lack the

substrate necessary to perform the Krebs Cycle and the ETC efficiently. This leads to cellular starvation. When cells cannot produce ATP, the patient experiences the persistent, overwhelming fatigue characteristic of the condition. The body is essentially starving in a land of plenty because the fuel cannot reach the engine. This metabolic failure also forces the body to utilise alternative fuels, such as fatty acids, through beta-oxidation. While this provides some energy, it is less efficient and can lead to the production of ketone bodies, which in extreme cases can lead to metabolic acidosis.

4. Analysis of Recording and Tracking Data (PART D)

HbA1c (The Long-term Average): This is our most important tool for tracking your progress. While a regular finger-prick test shows your sugar at that exact second, the HbA1c test looks back over the last three months. It works by measuring how much sugar has "stuck" to your red blood cells. Since these cells live for about 120 days, the test gives us a reliable average of your overall control. A result of **48 mmol/mol** or higher confirms a diagnosis of diabetes. We use this number to decide if your current plan is working or if we need to adjust your medication to keep you safe.

Daily Testing (The Snapshots): We use home finger-prick tests to get "snapshots" of your day. These are helpful for seeing how your body reacts to specific things, like a high-carb meal, a stressful situation, or a bout of exercise. By seeing these immediate results, you can learn which parts of your routine cause your sugar to spike and which parts help keep it steady.

Preventing Damage (The Team Approach): We use all this data to find patterns before they become problems. If sugar levels stay high for too long, they act like "sand in a machine," causing wear and tear on your small and large blood vessels. This is also why we track your blood pressure and cholesterol. These three factors; sugar, pressure, and fat, often travel together. By managing all of them at once, we work as a team to protect your heart, kidneys, and eyes, ensuring you stay healthy for the long term.

5. Pathophysiological Complications of Hyperglycaemia

High blood sugar acts as a slow-acting toxin to the body over time. It causes the basement membranes of small blood vessels (capillaries) to become thick and fragile. This leads to several specific types of **Microvascular Damage**:

Retinopathy: Damage to the delicate blood vessels in the retina, which can lead to permanent blindness.

Nephropathy: Scarring of the nephrons, which are the kidney's vital filtering units.

Neuropathy: Nerve damage that results in a loss of "protective sensation". This typically follows a "stocking-glove" pattern, meaning it affects the longest nerves in the feet and hands first. This is particularly dangerous because injuries like blisters can go unnoticed, leading to deep ulcers, severe infection, and even amputation.

In addition to small vessel damage, T2DM causes **Macrovascular Damage** by speeding up atherosclerosis, the buildup of fatty plaques in large arteries. This significantly raises the risk of life-altering events such as heart attacks and strokes. Furthermore, chronic high sugar leads to the overproduction of reactive oxygen species (ROS), causing oxidative stress that damages cellular DNA. It also triggers the formation of **Advanced Glycation End-products (AGEs)**, harmful compounds that stiffen blood vessels and accelerate the ageing of internal organs.

6. Evidence-Based Lifestyle Recommendations (PART E)

The management of T2DM must be rooted in physiological interventions that address the underlying insulin resistance:

Exercise (The Insulin-Independent Pathway): Physical activity is a powerful non-drug treatment. When muscles contract, they trigger the movement of GLUT4 transporters to the cell surface through a "back door" called the AMPK pathway. This process works even if a patient is highly insulin-resistant, allowing sugar to enter the cells and lowering blood glucose levels. Even a simple ten-minute brisk walk after a meal can noticeably prevent a sugar spike.

Dietary Quality and the Glycaemic Index (GI): We advise a diet rich in fibre and low-GI carbohydrates, such as beans and whole grains. These foods are broken down slowly, causing a gentle rise in glucose rather than a sharp spike. This takes the pressure off the pancreas and reduces oxidative stress.

The DIRECT Trial and Remission: Clinical evidence, specifically the **DIRECT Trial**, shows that T2DM is not necessarily a "life sentence". By losing a significant amount of weight (around 15kg), patients can remove "ectopic" fat stored in the liver and pancreas. This allows the beta cells to "reboot" and begin secreting insulin normally again, potentially putting the disease into remission.

Smoking Cessation: Smoking introduces nicotine and carbon monoxide, which cause body-wide inflammation and damage the lining of blood vessels. For someone with diabetes, smoking acts as a "multiplier" for heart disease risk. Quitting is essential to protect the vascular system from the combined assault of toxins and high sugar.

7. The Nurse-Patient Partnership and Health Literacy

The success of T2DM management depends on health literacy. As a practice nurse, I do not just tell patients to eat better; I explain the "lock and key" model of insulin. When a patient understands that their fatigue is caused by a cellular energy deficit, they are more motivated to engage with lifestyle changes. By using the routine data collected in the surgery (HbA1c, BMI, blood pressure), we can create a visual "map" of the patient's progress. This empowers the patient, fostering a sense of self-efficacy, the belief that they have the power to influence their own health outcomes. In our surgery, we have implemented group consultation sessions where patients can share their experiences and learn from one another, improving the collective health literacy of our patient population.

8. Conclusion (PART F)

Type 2 Diabetes is a complex problem, but it can be managed well through understanding the science and using data. By looking at how the pancreas works and how cells make energy, we can see why the condition affects the whole body. My goal as a nurse is to use health data to guide better lifestyle choices. Through movement, better food, and quitting smoking, we can help patients avoid serious complications and even move toward reversing the disease entirely. With the right support, people with diabetes can live long, healthy lives.

Total Word Count (excluding titles) = 2028

9. Reference List (PART G - Harvard Style)

British Medical Association (BMA) (2026) *Managing Diabetes in Primary Care*. Available at: <https://www.bma.org.uk> (Accessed: 14 February 2026).

Diabetes UK (2025) *The State of the Nation: Diabetes Challenge*. London: Diabetes UK Publishing. (Accessed: 14 February 2026).

Hall, J.E. and Hall, M.E. (2020) *Guyton and Hall Textbook of Medical Physiology*. 14th edn. Philadelphia: Elsevier. (Accessed: 14 February 2026).

Lean, M.E., Leslie, W.S., Barnes, A.C. and Taylor, R. (2018) 'Primary care-led weight management for remission of type 2 diabetes (DiRECT)', *The Lancet*, 391(10120), pp. 541-551. (Accessed: 14 February 2026).

Marieb, E.N. and Hoehn, K. (2022) *Human Anatomy & Physiology*. 12th edn. London: Pearson Education. (Accessed: 14 February 2026).

National Institute for Health and Care Excellence (NICE) (2022) *Type 2 diabetes in adults: management (NG28)*. Available at: <https://www.nice.org.uk/guidance/ng28> (Accessed: 15 February 2026).

National Institute for Health and Care Excellence (NICE) (2023) *Diabetes - type 2: Prevention (PH38)*. London: NICE Guidelines. (Accessed: 15 February 2026).

Porth, C.M. (2022) *Essentials of Pathophysiology*. 5th edn. Philadelphia: Wolters Kluwer. (Accessed: 15 February 2026).

Stanfield, C.L. (2023) *Principles of Human Physiology*. 7th edn. London: Pearson. (Accessed: 15 February 2026).

Tortora, G.J. and Derrickson, B. (2023) *Principles of Anatomy and Physiology*. 16th edn. Hoboken: Wiley & Sons. (Accessed: 15 February 2026).

World Health Organisation (WHO) (2024) *Diabetes: Key Facts*. Available at: <https://www.who.int/news-room/fact-sheets/detail/diabetes> (Accessed: 16 February 2026).

Zimmet, P.Z., Magliano, D.J., Herman, W.H. and Shaw, J.E. (2019) 'Diabetes: a 21st century challenge', *The Lancet Diabetes & Endocrinology*, 7(1), pp. 56-66. (Accessed: 16 February 2026).