

Module Name: Introduction to Public Health

Module Code: LCHS4008

Student Name: Lucas Rocha da Silva

Student ID: 2440423

Lecturer Name: Caroline Salami

Table of contents

1. **Introduction**
 - 1.1 Definition of Stroke
 - 1.2 Signs and Symptoms
 - 1.3 Incidence and Severity
2. **Measurement and Monitoring of Stroke**
 - 2.1 Clinical Assessment and Diagnosis
 - 2.2 Population Data and Surveillance
3. **Control and Prevention of Stroke**
 - 3.1 Risk factors
 - 3.2 Prevention and Medical Control
4. **Impact of Stroke on individuals and Communities**
 - 4.1 Physical and Psychological Effects
 - 4.2 Social and Community Impact
5. **Health and Social Care Support**
 - 5.1 Rehabilitation Services
 - 5.2 Social Care and Community Support
6. **Conclusion**
7. **References**

1. Introduction

1.1 Definition of Stroke

Stroke is a serious condition that occurs when the blood supply to part of the brain is suddenly interrupted. As a result, the brain does not receive enough oxygen and nutrients, which can quickly lead to damage of brain cells. There are two main types of stroke. The most common one is ischaemic stroke, which is caused by a blockage in a blood vessel. On the other hand, haemorrhagic stroke happens when a blood vessel bursts and causes bleeding in the brain (NHS, 2023). In both cases, the damage can happen very quickly, which is why stroke is treated as a medical emergency. If not treated in time, it can lead to permanent disability or even death.

1.2 Signs and Symptoms

The symptoms of stroke usually appear suddenly and can vary depending on the area of the brain affected. In most cases, people may notice weakness or numbness on one side of the body. They may also have difficulty speaking or understanding what others are saying. Other symptoms can include blurred vision, confusion and a severe headache. In the UK, campaigns such as FAST are used to help people recognise these signs early. This is important because early treatment can make a big difference in recovery (WHO, 2023). Sometimes, symptoms may be mild at first, but they can quickly become more serious.

1.3 Incidence and Severity

Stroke continues to be one of the leading causes of death and long-term disability worldwide. According to the World Health Organization (WHO, 2023), millions of people are affected each year. In the UK, around 100,000 people have a stroke annually, which shows how common the condition is (Stroke Association, 2022). In addition, many people who survive a stroke are left with long-term health problems, which increases the pressure on healthcare services. The risk of stroke also increases with age, although it can still affect younger individuals.

2. Measurement and Monitoring of Stroke

2.1 Clinical Assessment and Diagnosis

Stroke is usually diagnosed in hospital using imaging techniques such as CT scans or MRI scans. In most cases, a CT scan is done first because it is quicker and can show if there is bleeding in the brain. MRI scans can provide more detailed information, especially in less obvious cases. Doctors also use assessment tools to measure how severe the stroke is and to monitor recovery over time. This helps them decide on the most appropriate treatment for each patient (NICE, 2019). Early diagnosis is essential to reduce long-term complications.

2.2 Population Data and Surveillance

At a national level, stroke is monitored using healthcare data. In the UK, NHS England collects information about stroke cases, including hospital admissions, recovery outcomes and death rates. This data is useful for identifying trends and understanding which groups are most affected. It also helps in planning healthcare services and improving prevention strategies over time (NHS England, 2023). Monitoring data over time also allows comparisons between regions and helps highlight inequalities in health outcomes.

3. Control and Prevention of Stroke

3.1 Risk Factors

There are a number of risk factors that can increase the chances of having a stroke. High blood pressure is considered the most important one, as it can damage blood vessels over time. Other risk factors include smoking, poor diet, lack of exercise and drinking too much alcohol. In addition, medical conditions such as diabetes and atrial fibrillation can also increase the risk (CDC, 2023). Many of these factors are linked to lifestyle, which means they can often be reduced with changes in behaviour. For example, reducing salt intake can help lower blood pressure.

3.2 Prevention and Medical Control

Preventing stroke mainly involves managing these risk factors. For example, eating a balanced diet, staying physically active and stopping smoking can all reduce the risk. Public health campaigns in the UK often focus on raising awareness about these changes.

However, prevention is only one part of the process. Stroke is also controlled through medical treatment and ongoing care. For instance, if a patient arrives at hospital early, they may receive thrombolytic treatment to break down blood clots. In some cases, a procedure such as thrombectomy may be used. After a stroke, patients are usually given medication to control blood pressure and cholesterol levels, which helps reduce the risk of another stroke (NICE, 2019). Regular follow-ups with healthcare professionals are also important.

4. Impact of Stroke on Individuals and Communities

4.1 Physical and Psychological Effects

The impact of stroke on an individual can be quite serious. Some people may lose movement on one side of their body, while others may struggle with speech or memory. These changes can make everyday activities much more difficult and affect independence.

In addition to physical problems, many people also experience emotional difficulties. It is common for stroke survivors to feel frustrated, anxious or depressed as they adjust to their new situation (Stroke Association, 2022). Recovery can take a long time, and progress is often gradual rather than immediate.

4.2 Social and Community Impact

Stroke also affects families and communities. In many cases, family members take on the role of caregivers, which can be challenging both physically and emotionally. This can also lead to financial pressure, especially if the person is unable to return to work.

From a wider perspective, stroke places a significant burden on healthcare systems. Long-term care, rehabilitation and support services require a large amount of resources. This means that stroke has both personal and social consequences (NHS England, 2023). It also highlights the importance of prevention at a population level.

5. Health and Social Care Support

5.1 Rehabilitation Services

Rehabilitation plays an important role in helping people recover after a stroke. Physiotherapy can help improve movement and strength, while occupational therapy focuses on daily activities such as dressing and eating. Speech and language therapy is also important for those who have communication difficulties.

These services are usually provided through the National Health Service and are adapted to each person's needs. Early support can make a noticeable difference in recovery (NICE, 2019). Consistent rehabilitation can also improve long-term outcomes.

5.2 Social Care and Community Support

In addition to healthcare, social support is also very important. Some people may need help at home, including support with daily tasks. Equipment such as walking aids can also help improve independence.

Organisations like the Stroke Association provide support groups and advice for both patients and their families. This kind of support can help people feel less isolated and more confident in managing their condition (Stroke Association, 2022). Community support can also improve mental wellbeing.

6. Conclusion

In conclusion, stroke is a serious condition that has a major impact on both individuals and society. It can lead to long-term health problems and affect a person's independence and quality of life. However, many of the risk factors can be reduced through lifestyle changes.

Early recognition of symptoms is very important, as it allows for quicker treatment and better outcomes. In addition, rehabilitation and social support play a key role in helping individuals recover and adjust after a stroke.

Overall, reducing the impact of stroke requires a combination of prevention, medical treatment and ongoing support. By improving awareness and access to healthcare services, it is possible to reduce the number of cases and support those affected more effectively.

7. References

World Health Organization (WHO) (2023) *Stroke, cerebrovascular accident*. Available at: <https://www.who.int> (Accessed: 18 March 2026).

NHS (2023) *Stroke overview*. Available at: <https://www.nhs.uk> (Accessed: 21 March 2026).

NHS England (2023) *Stroke statistics and services*. Available at: <https://www.england.nhs.uk> (Accessed: 24 March 2026).

Stroke Association (2022) *State of the Nation: Stroke statistics*. Available at: <https://www.stroke.org.uk> (Accessed: 26 March 2026).

National Institute for Health and Care Excellence (NICE) (2019) *Stroke and transient ischaemic attack in over 16s*. Available at: <https://www.nice.org.uk> (Accessed: 19 March 2026).

Centers for Disease Control and Prevention (CDC) (2023) *About stroke*. Available at: <https://www.cdc.gov> (Accessed: 22 March 2026)