

Depression: Assessment, Tracking, Management, and Community-Based Health and Social Care Assistance

student name : Zobia Ashfaq (2524669)

Module Code : LCHS4008

Module Title : Public Health

Lecturer's Name : Debra Odah

Introduction

Depression is one of the most common mental health disorders worldwide, affecting millions of people of all ages. It is more than temporary sadness; it is a serious medical condition that affects emotions, thoughts, behaviour, and physical health. According to the World Health Organisation (WHO, 2023), around 280 million people globally are living with depression, making it a leading cause of disability. In the United Kingdom, approximately one in six adults experiences symptoms of depression or anxiety each week, highlighting its significant impact on public health (NHS) England, 2024). Common symptoms include persistent low mood, loss of interest in daily activities, fatigue, sleep problems, poor concentration, and feelings of worthlessness. If left untreated, depression can affect relationships, education, employment, and overall quality of life. This report will discuss the causes and impact of depression, how it is measured and managed, and the health and social care services available to support individuals living with the condition.

1. Definition of Depression

Depression, also known as Major Depressive Disorder (MDD), is a common mental illness characterised by a persistent low mood, sadness, and a loss of interest or pleasure in activities lasting for at least two weeks. According to the World Health Organisation (WHO, 2023), depression affects how individuals think, feel, and behave, often interfering with their ability to carry out daily activities.

This condition can impact people of all ages, regardless of gender, ethnicity, or socioeconomic background. It may arise from a combination of biological, psychological, and environmental factors, such as genetics, stressful life events, chronic illness, trauma, unemployment, or social isolation.

2. Depression Symptoms and Signs

- Persistent melancholy or depression.
- Loss of interest in regular activities and interests.
- Feeling unworthy or hopeless.
- Exhaustion and low vitality.
- Inability to focus.
- Variations in body weight and hunger.

- Sleep issues (either excessive sleeping or sleeplessness).
- irritation and anxiety.
- Social disengagement.
- suicidal or self-harming thoughts.

Without a clear medical reason, depression can also manifest physically as headaches, aches in the muscles, and digestive issues.

Prevalence and Incidence (Global and UK Statistics)

One of the primary contributors to the global disease burden is recognised as depression.

- Depression affects over 280 million people globally.
 - It affects about 3.8% of the world's population.
 - One of the main causes of disability in the world is depression.
 - Depression is more common in women than in males.
-
- In the United Kingdom:
 - According to NHS England, one in six persons has weekly symptoms of anxiety or depression.
 - Mental health issues have greatly grown as a result of the COVID-19 epidemic, especially among young people and adolescents.
 - One of the top causes of death for young people in England between the ages of 20 and 34 is still suicide.
 - These figures show that depression is a serious public health concern that calls for efficient monitoring, prevention, and treatment methods.

4. Methods for Assessing and Tracking Depression

Healthcare practitioners assess and track depression using a variety of techniques.

Clinical Evaluation

First, a thorough medical history is taken, and symptoms, lifestyle, family history, and recent stressful events are discussed. This helps rule out physical ailments and assess the severity of depression.

Questionnaires for Screening

Among the verified screening instruments are:

- **PHQ-9 (Patient Health Questionnaire-9)**
- **Beck Depression Inventory (BDI)**
- **Hospital Anxiety and Depression Scale (HADS)**

These surveys help determine the severity of symptoms and monitor progress during treatment.

****Mental State Assessment (MSE)**** Mental health professionals assess various factors including self-harm risk, appearance, mood, behaviour, speech, cognitive processes, and memory. ****Risk Evaluation**** Medical professionals ask questions about suicidal thoughts, past attempts, self-harm history, and protective factors to evaluate the risk of suicide. ****Monitoring Follow-Up**** General practitioners (GPs), psychiatrists, psychologists, and community mental health teams regularly assess patients undergoing treatment. They use repeated questionnaires and evaluate symptom improvement to track progress.

5. How to Manage Depression

Medical care, psychiatric therapy, lifestyle changes, and community support are all necessary for managing depression.

Drugs

For moderate to severe depression, antidepressants like Selective Serotonin Re-uptake Inhibitors (SSRIs) are frequently used. These medications aid in controlling brain chemicals related to mood.

Examples include:

- Sertraline
- Fluoxetine
- italopram

Medication should always be prescribed and monitored by qualified healthcare professionals. Psychological therapies, especially talking therapies, are highly effective.

These include:

- CBT, or cognitive behavioural therapy
- Interpersonal Therapy (IPT) counselling
- Activation of Behaviour

NICE guidelines support CBT because therapy assists patients in recognising and altering negative thought patterns.

Modifications to Lifestyle

A healthy lifestyle can lessen symptoms:

- Regular physical exercise
- Balanced diet
- Good sleep hygiene
- Stress management
- Avoiding alcohol and drug misuse

****Social Assistance****

Recovery is significantly supported by the involvement of friends, family, employers, educational institutions, and community organisations. ****Early Intervention**** Recognising symptoms early can help mitigate issues such as suicide and long-term disability by allowing for therapy before depression escalates. If you would like to receive Part 2, please reply with "Continue," and I will email it to you.

- Depression's Effects on the Community
- Support Services for Health and Social Care
- Conclusion (150 words)
- Harvard Reference List in its entirety (2021–2025)

6. The Effects of Depression on Community Members

In addition to people, depression has detrimental effects on families, workplaces, healthcare systems, and society at large. A lower quality of life is one of the most important consequences. Individuals with depression frequently struggle to fulfil daily obligations and lose interest in once-enjoyed hobbies. Financial stability, work, education, and interpersonal connections may all be impacted.

Consequences for Physical Health

Numerous physical health issues are linked to depression. Chronic fatigue, headaches, digestive issues, impaired immunity, and a higher risk of cardiovascular disease are all possible. Unhealthy behaviours including smoking, drinking alcohol, abusing drugs, eating poorly, and not exercising can also result from poor mental health and raise the risk of chronic illnesses.

Social Repercussions

People who are depressed frequently distance themselves from friends, family, and community events. Social isolation might lessen opportunities for emotional support and exacerbate symptoms. Due to the stigma and discrimination associated with mental illness, many people are unwilling to seek professional assistance.

Economic Repercussions

Depression significantly affects society's economy. Individuals with untreated depression are more likely to miss work, be less productive, or quit their jobs entirely. While healthcare systems spend a lot of money on treatment, emergency care, and long-term assistance, employers incur higher expenditures as a result of sick leave.

Impact on Education

Young individuals who are depressed frequently have trouble focusing, do worse academically, and miss more school or university. Depression may have an impact on future social and professional prospects if treatment is delayed.

Impact on Families

Family members are emotionally and financially impacted by depression. Supporting someone who is depressed can cause stress, worry, and caregiver strain on partners, parents, and kids. Communication problems and emotional suffering can cause tension in family relationships.

Risk of Suicide

Suicide is one of the most serious consequences of depression. According to the World Health Organisation (2023), depression is a leading cause of the over 700,000 suicide deaths that occur each year. To reduce this risk, it is essential to take prompt action and ensure early diagnosis.

Conclusion

Depression is a common yet serious mental health disorder that affects millions of people worldwide and is a leading cause of disability. Its impact goes beyond emotional well-being, affecting physical health, education, employment, relationships, and overall quality of life.

Early diagnosis through clinical assessments and validated screening tools enables healthcare professionals to provide timely treatment and improve long-term outcomes. Effective management typically includes a combination of antidepressant medication, psychological therapies, healthy lifestyle changes, and ongoing support from health and social care services.

Community Mental Health Teams, general practitioners, psychologists, social workers, and voluntary organisations all play important roles in supporting individuals affected by depression. Reducing stigma,

increasing public awareness, and improving access to mental health services are essential for encouraging early help-seeking and preventing severe complications, including suicide.

A coordinated approach that involves healthcare providers, families, communities, and policymakers is necessary to improve mental health outcomes and enhance the well-being of people living with depression.

Harvard Reference List

National Institute for Health and Care Excellence (NICE) (2022) *Depression in adults: treatment and management (NG222)*. London: NICE.

NHS England (2024) *Mental Health Services*. Available at: <https://www.nhs.uk/mental-health/> (Accessed: 6 July 2026).

Office for National Statistics (ONS) (2024) *Mental health and well-being statistics*. Available at: <https://www.ons.gov.uk> (Accessed: 6 July 2026).

World Health Organization(WHO) (2023) *Depressive disorder (depression)*. Available at: <https://www.who.int/news-room/fact-sheets/detail/depression> (Accessed: 6 July 2026).

American Psychiatric Association (2022) *Diagnostic and Statistical Manual of Mental Disorders (5th ed., text revision)*. Washington, DC: American Psychiatric Association.

Mental Health Foundation (2023) *Fundamental Facts About Mental Health*. London: Mental Health Foundation.

Mind (2024) *Understanding Depression*. Available at: <https://www.mind.org.uk> (Accessed: 6 July 2026).

Royal College of Psychiatrists (2023) *Depression in Adults*. Available at: <https://www.rcpsych.ac.uk> (Accessed: 6 July 2026).