

Complete, fold and seal
(see note 1 below)

84%

SURNAME:

Dand

FORENAME(S):

Alexander

Martin

Candidate's Exam Number:

Book No.

**Total Books
Handed In:**

Programme Title:

Year/Level/Stage:

Examination Paper Title:

Examination Paper Ref No.:

Desk Letter/Number:

Date: ____ / ____ / 20____

Time: _____ am/pm

Room Name or No.:

Instructions to Candidates:


1. Please give the information requested above and enter your name in the space provided at the top right of this page. If your answers are to be marked under a scheme whereby the examiners will not know your identity, fold over and seal down the top right-hand corner of the page (using the gummed label provided) in order to conceal your name.
2. Commence each answer on a fresh page.
3. Do all rough work in the answer book and cross through it.
4. No part of this booklet is to be torn out.
5. Supplementary answer books if used must:-
 - (a) bear the candidate's name (concealed, where appropriate), reference number and paper title;
 - (b) be numbered consecutively;
 - (c) be fastened together by the tags available from the invigilator.
6. Every book must be handed in. In no circumstances may answer books, used or unused, be taken out of the exam room.
7. Your answers must be legibly written **IN INK**.
The Examiners can either ignore illegible script or have it typed and the expense of typing charged to the candidate.

Summary of Marks

Question Numbers Answered	1st Examiner	2nd Examiner (optional)	3rd Examiner (optional)
1	33		
2	34		
	67		
TOTAL			
RESULT			

Q₁.

a) i) $E = mgh$
 $\frac{E}{mh} = g$




1

ii) $T = 2\pi\sqrt{\frac{L}{g}}$

$$\frac{T}{2\pi} = \sqrt{\frac{L}{g}}$$

$$\left(\frac{T}{2\pi}\right)^2 = \frac{L}{g}$$

$$\frac{L}{\left(\frac{T}{2\pi}\right)^2} = g$$



4

iii) $V^2 = u^2 + 2gs$

$$V^2 - u^2 = 2gs$$

$$\frac{V^2 - u^2}{2s} = g$$




2

b) i) $(pq)^0 = 1$ (anything to power 0 = 1)




1

ii) $(\sqrt{q})^4 = (q^{\frac{1}{2}})^4$
 $= q^2$



2

iii) $(p+q)(3p-q) + (2p+q)(p-q)$
 $3p^2 - pq + 3pq - q^2 + 2p^2 - 2pq + pq - q^2$
 $5p^2 + pq - 2q^2$



3

C) $\frac{2}{x-1} + \frac{3}{x-4}$ multiply by both denominators.

$$\frac{5x-5}{(x+1)(x-4)} = \frac{A}{x+1} + \frac{B}{x-4}$$

multiply by both denominators.

$$= A(x-4) + B(x+1)$$

Sub $x=4$

$$= 4(0) + B5$$

$$d) i) 2^{(3x-5)} = 4^{(5x-6)}$$

$$(3x-5) \log 2 = (5x-6) \log 4$$

0.6020

$$0.3010$$

$$0.3010(3x-5) = 0.6020(5x-6)$$

$$0.9031x - 1.505 = 3.0105x - 3.6126$$

Transpose

$$2.1076 = 2.1074x$$

$$x = 1 \text{ (3dp).}$$

✓

4

$$ii) \ln(x^2-4) - \ln(x+8) = \ln x$$

$$\ln\left(\frac{x^2-4}{x+8}\right) = \ln x$$

Take out logs.

$$\frac{x^2-4}{x+8} = x$$

Transpose.

$$x^2+4 = x(x+8)$$

$$x^2+4 = x^2+8x$$

$$4 = 8x$$

$$\frac{4}{8} = x$$

$$0.5 = x$$

3

$$e) \text{ if } y = \frac{2-3x}{1-5x^2} = 0$$

$$\frac{dy}{dx} = \frac{(1-5x^2)(-3) - (2-3x)(-10x)}{(1-5x^2)^2}$$

-(-20x+30x^2)

$$= \frac{15x^2-3+20x-30x^2}{(1-5x^2)^2}$$

$$\frac{dy}{dx} = \frac{-15x^2+20x-3}{(1-5x^2)^2}$$

$$\text{When } x = -1$$

$$\frac{dy}{dx} = \frac{-15(-1)^2+20(-1)-3}{(1-5(-1)^2)^2}$$


$$= \frac{-15-20-3}{4^2}$$

$$= \frac{-38}{16}$$

$$\frac{dy}{dx} = -2.375$$

5

F)



$$p^2 + r^2 = q^2$$

$$(2x)^2 + (3x)^2 = 256$$

$$4x^2 + 9x^2 = 256$$

$$13x^2 = 256$$

$$x^2 = \frac{256}{13}$$

$$x = \sqrt{\frac{256}{13}}$$

$$x = 4.4376 \text{ cm}$$

$$p = 8.9 \text{ cm (1dp)}$$

$$r = 13.3 \text{ cm (1dp)}$$

6

33

②

$$a) i) 4 - q^2 = (2+q)(2-q)$$

Difference of 2 squares.

2

$$ii) 3ab + 4cd - cb - 12ad$$

$$b(3a-c) + 4d(c-3a)$$

2

$$iii) p^2 - 2p - 63$$

$$\begin{array}{r} 1 \ 63 \\ 3 \ 21 \\ \hline 7 \ 9 \end{array}$$

$$(p+9)(p-7)$$

$$\times p^2 - 7p + 9p - 63$$

$$iv) q^2 - 4qr$$

$$q(q-4r)$$

$$(q+2r)(q-2r)$$

$$q^2 + 2qr + 2qr - 4r^2$$

1

$$b) i) \text{Solve } (2x-1)(3x+5) = (6x-5)(x-9)$$

$$6x^2 + 10x - 3x - 5 = 6x^2 - 54x - 5x + 45$$

$$7x - 5 = 45 - 59x$$

$$66x = 50$$

$$x = \frac{50}{66}$$

$$x = 0.76 \quad (2dp)$$

✓

5

$$ii) \frac{x-4}{5} = \frac{2x-1}{3}$$

Cross multiply

$$3x-12 = 10x-5$$


$$-7x = 7$$

$$x = -1.00 \quad (2dp)$$

✓

4

c)



$$A + B + C = 180$$

$$A = 95^\circ$$

Sin rule $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

$$10.0382 = \frac{b}{\sin 40}$$

$$10.0382 \sin 40 = b$$

$$6.54 = b$$

$$AC = 6.5m \text{ (1dp).}$$

$$10.0382 = \frac{c}{\sin 45}$$

$$10.0382 \sin 45 = c$$

$$7.1m = c$$

$$AB = 7.1m \text{ (1dp)}$$

5

d). i)

$$4 \sin x = -3$$

$$\sin x = -\frac{3}{4}$$

$$x = \sin^{-1} 0.75$$


$$= 48.59$$

as $\sin$ is $-ve$, in $Q + 7$ quadrants

$$x = 228.59^\circ$$

$$x = 311.41^\circ$$

$$0 \leq x < 360$$



3

ii) $\cos x = -0.5$


$$x = \cos^{-1} 0.5$$

$$= 60$$

$$x = 120^\circ$$

$$x = 240^\circ$$

$$0 \leq x < 360$$



2

$$e) i) y = x^3 e^{2x}$$

using Product rule.

$$\frac{dy}{dx} = x^3(2e^{2x}) + e^{2x}(3x^2)$$

2

$$ii) y = \sin 5x \cos 3x$$

$$\frac{dy}{dx} = \sin 5x(-3\sin 3x) + \cos 3x(5\cos 5x)$$

1

F)

$$Z = 2x^3y - 7y^3x + \sin x \cos y$$

$$\frac{dz}{dx} = 6x^2y - 7y^3 + \cos x \cos y$$

$$\frac{dz}{dy} = 2x^3 - 21y^2x - \sin x \sin y$$

$$① \frac{\partial z}{\partial x} = 12xy - 5\sin x \cos y$$

$$③ \frac{\partial z}{\partial x} = 6x^2 - 21y^2 - \cos x \sin y$$

$$② \frac{\partial z}{\partial y} = 6x^2 - 21y^2 - \cos x \sin y$$

$$④ \frac{\partial z}{\partial y} = 42y - \sin x \cos y$$

7

34

Question No.

Question No.

Examiner Only

Rhif y Cwestiwn

Arholwr yn u

Question No.

Examiner On

Arholwr yn unig

fy Cwestiwn