

Managing Chronic Heart Failure: A Comprehensive Clinical and Operational Guide for Health and Social Care Professionals

Page 1: Table of Contents

Page 2: Introduction to Chronic Heart Failure

Page 3: Definition, Classification, and Epidemiology

Page 4: Aetiology, Risk Factors, and Pathophysiology

Page 5: Biopsychosocial Impact on Individuals and Families

Page 6: Evidence-Based NHS Treatments and Pathways

Page 7: Theoretical Frameworks and Models of Care

Page 8: Political Agendas and Implementation Challenges

Page 9: Recommendations for Person-Centred and End-of-Life
Care

Page 10: References

Page 11: References

Page 12: References



Introduction to Chronic Heart Failure

Chronic Heart Failure (CHF) is one of the most complex, formidable, and resource-intensive long-term conditions managed globally and across the UK. It is characterized by the heart's inability to pump blood at a rate commensurate with the body's metabolic requirements, leading to high hospital admission rates and high mortality.

This brochure has been commissioned for health and care professionals, including general practitioners, specialist heart failure nurses, clinical pharmacists, and social care commissioners and will aim deliver an exhaustive, evidence-based synthesis integrating cardiac pathophysiology, psychosocial impacts, theoretical care models, and NHS-aligned clinical strategies.

Modern healthcare requires a shift away from reactive acute care toward proactive, integrated, and person-centred delivery due to the unpredictable trajectory of CHF.

Managing CHF requires robust cross-sector collaboration between health and social care to address multi-morbidity, frailty, and caregiver support.



Definition, Classification, and Epidemiology

Definition: CHF is a clinical syndrome resulting from structural or functional cardiac impairment affecting ventricular filling or blood ejection.

Heart Failure with Reduced Ejection Fraction (HFrEF): Defined as an LVEF $\leq 40\%$, characterized by impaired myocardial contractility and systolic dysfunction.

Heart Failure with Mildly Reduced Ejection Fraction (HFmrEF): Defined as an LVEF between 41% and 49% .

Heart Failure with Preserved Ejection Fraction (HFpEF): Defined as an LVEF $\geq 50\%$ with diastolic dysfunction, accounting for nearly half of all hospitalizations.

Global Burden: Over 64 million people live with heart failure worldwide, scaling exponentially with age.

UK Burden: Over 1 million individuals are diagnosed in the UK, consuming over £2 billion annually in NHS expenditures and accounting for 5% of emergency medical admissions. Five-year mortality remains near 50% .

Aetiology, Risk Factors, and Pathophysiology

Cellular Mechanism: Sustained neurohormonal activation of the Sympathetic Nervous System (SNS) and Renin-Angiotensin-Aldosterone System (RAAS) initially maintains blood pressure but eventually accelerates myocardial cell apoptosis and interstitial fibrosis.

Risk Factor Category	Specific Clinical Determinants	Pathophysiological Mechanism
Non-Modifiable	Advanced age, Male sex (HFrEF), Female sex (elderly HFpEF), Genetic cardiomyopathies.	Age-related arterial stiffening and cardiomyocyte loss reduce cardiac reserve.
Modifiable	Hypertension, Coronary Artery Disease (CAD), Type 2 Diabetes, Obesity, Chronic Kidney Disease, Smoking.	Chronic pressure overload, ischemic scarring, microvascular dysfunction, and lipotoxic remodeling.

Biopsychosocial Impact on Individuals and Families

Biopsychosocial Implications

Physical Impact: Exertional dyspnoea, orthopnoea, paroxysmal nocturnal dyspnoea (PND), profound skeletal muscle fatigue, and peripheral oedema. Cardiac cachexia occurs in advanced stages due to chronic systemic inflammation.

Psychological Impact: High rates of anxiety, autonomic panic during breathlessness episodes, and comorbid major depression affecting 20\% to 40\% of patients, which independently predicts worse survival.

Social and Caregiver Burden: Informal caregivers experience severe physical exhaustion, depression, and strain from managing complex fluid restrictions and polypharmacy. Patients face social isolation due to loss of driving privileges and mobility.

Evidence-Based NHS Treatments and Pathways

Evidence-Based NHS Management

Four-Pillar Pharmacotherapy for HFrEF: Modern management relies on concurrent initiation and titration of:

- A. ARNI (Sacubitril/Valsartan) or ACE inhibitors/ARBs.
- B. Evidence-based beta-blockers (Bisoprolol, Carvedilol).
- C. Mineralocorticoid Receptor Antagonists (MRAs).
- D. SGLT2 Inhibitors (Dapagliflozin/Empagliflozin) for cardiorenal protection.

Diuretics: Loop diuretics (Furosemide) are used strictly for symptom relief and fluid overload management.

Device Therapy: Implantable Cardioverter-Defibrillators (ICDs) and Cardiac Resynchronization Therapy (CRT).

Cardiac Rehabilitation: Structured exercise training improves muscle function and quality of life, though participation barriers persist among older and deprived cohorts.

Theoretical Frameworks and Models of Care

Care Models

UK Model (Specialist Nurse-Led Pathways): Specialist Heart Failure Nurses (HFSNs) operate across primary care networks and secondary care to provide expert assessment, aggressive pharmacological titration, and rapid post-discharge surveillance.

International Model (Chronic Care Model - CCM): Developed by Wagner et al. (2001), the CCM identifies six systemic components:

- A. Health System Organization.
- B. Delivery System Design.
- C. Decision Support.
- D. Clinical Information Systems.
- E. Self-Management Support.
- F. Community Resources.

Combining the structural breadth of the CCM with the clinical agility of UK HFSN teams maximizes patient outcomes.

Political Agendas and Implementation Challenges

Policy and Implementation

The NHS Long Term Plan: Identifies cardiovascular disease as a primary clinical priority, mandating expansion of specialist heart failure nursing and cardiac rehabilitation.

National Heart Failure Audit (NHFA): Measures clinical performance and medication optimization across trusts to drive public accountability.

Get It Right First Time (GIRFT): Aims to eliminate unwarranted clinical variations in diagnostic access and specialist staffing ratios.

Implementation Challenges: Severe workforce shortages among specialist nurses, primary care echocardiography bottlenecks leading to emergency admissions, and socioeconomic health inequalities requiring multi-agency collaboration.



Recommendations for Person-Centred and End-of-Life Care

Operational Recommendations

Self-Management: Educate patients on daily morning weight monitoring to catch sudden increases ($>2\text{ kg}$ over 3 days) and provide written diuretic action plans.

Digital Integration: Combine remote telemonitoring and symptom-tracking apps with face-to-face domiciliary reviews to ensure digital equity.

End-of-Life Care: Initiate early Advance Care Planning (ACP) regarding patient values, treatment ceilings, and the deactivation of ICDs as death approaches.

Palliative Support: Collaborate with specialist palliative care teams to manage refractory breathlessness using low-dose opioids and anxiolytics.

References

- Alkhalafan, F., et al. (2022) 'Post-discharge transition care models in heart failure: A systematic review and meta-analysis', *European Journal of Heart Failure*, 24(4), pp. 678-690.
- Anker, S. D., et al. (2003) 'The wasting cachexia of chronic heart failure: Mechanisms and therapeutic implications', *International Journal of Cardiology*, 85(1), pp. 51-60.
- Blue, L., et al. (2001) 'Randomised controlled trial of specialist nurse intervention in heart failure', *BMJ*, 323(7316), pp. 715-718.
- Bodenheimer, T., Wagner, E. H. and Grumbach, K. (2002) 'Improving primary care for patients with chronic illness', *JAMA*, 288(14), pp. 1775-1779.
- Bozkurt, B., et al. (2021) 'Universal definition and classification of heart failure', *Journal of Cardiac Failure*, 27(4), pp. 387-413.
- Briggs, T. (2020) *Get It Right First Time (GIRFT) Programme National Specialty Report for Cardiology*. London: NHS England.
- British Heart Foundation (BHF) (2023) *Heart Failure Statistics 2023: Facts and Figures*. London: BHF.
- Buckley, B. S., et al. (2007) 'Self-management education for patients with chronic heart failure: A systematic review', *Patient Education and Counseling*, 68(2), pp. 144-155.
- Cook, C., et al. (2014) 'The annual global economic burden of heart failure', *International Journal of Cardiology*, 171(3), pp. 368-376.
- Cowie, M. R., et al. (2014) 'The ESC Heart Failure Long-Term Registry: Characteristics and 1-year outcomes in patients with heart failure', *European Journal of Heart Failure*, 16(8), pp. 816-827.
- Dalal, H. M., et al. (2015) 'Individual personalized home-based heart failure cardiac rehabilitation', *European Journal of Heart Failure*, 17(7), pp. 720-729.
- Dickstein, K., et al. (2008) 'ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2008', *European Heart Journal*, 29(19), pp. 2388-2442.
- Dracup, K., et al. (2003) 'Quality of life in patients with heart failure', *Journal of Cardiac Failure*, 9(4), pp. 295-303.
- Felker, G. M., et al. (2011) 'Diuretic strategies in patients with acute decompensated heart failure', *New England Journal of Medicine*, 364(9), pp. 797-805.
- Freedland, K. E., et al. (2015) 'Depression and heart failure: Comorbidity, mechanisms, and clinical implications', *Heart Failure Clinics*, 11(4), pp. 543-558.
- Goldstein, N. E., et al. (2004) 'Management of implantable cardioverter-defibrillators in end-stage heart failure', *Annals of Internal Medicine*, 141(11), pp. 836-838.
- Goodlin, S. J. (2009) 'Palliative care for patients with heart failure', *Journal of the American College of Cardiology*, 54(5), pp. 386-396.
- Harkness, K., et al. (2012) 'Effect of a web-based educational program on anxiety, depression, and quality of life in heart failure', *Patient Education and Counseling*, 88(3), pp. 445-450.
- Heidenreich, P. A., et al. (2022) '2022 AHA/ACC/HFSA guideline for the management of heart failure', *Journal of the American College of Cardiology*, 79(17), pp. e263-e421.

References

- Inglis, S. C., et al. (2017) 'Structured telephone support or telemonitoring programmes for patients with chronic heart failure', *Cochrane Database of Systematic Reviews*, (8), art. no. CD007228.
- Jaarsma, T., et al. (2011) 'Self-care management in heart failure: Are we failing patients?', *European Journal of Heart Failure*, 13(4), pp. 367-375.
- Jaarsma, T., et al. (2020) 'Symptom burden and quality of life in heart failure', *Current Heart Failure Reports*, 17(5), pp. 235-244.
- Jenkinson, D., et al. (2013) 'The psychological impact of chronic heart failure: A qualitative study', *Journal of Advanced Nursing*, 69(8), pp. 1801-1811.
- Johnson, M. J., et al. (2013) 'Use of opioids for refractory dyspnoea in heart failure', *Journal of Pain and Symptom Management*, 45(2), pp. 292-301.
- Krumholz, H. M., et al. (2002) 'Readmission after hospitalization for congestive heart failure among Medicare beneficiaries', *Archives of Internal Medicine*, 162(1), pp. 99-105.
- Luttik, M. L., et al. (2007) 'Caregivers in heart failure: Managing the burden', *European Journal of Cardiovascular Nursing*, 6(3), pp. 224-230.
- Mann, D. L. and Bristow, M. R. (2005) 'Mechanisms and models in heart failure: The biomechanical model and beyond', *Circulation*, 111(21), pp. 2837-2849.
- Marmot, M., et al. (2020) *Health Equity in England: The Marmot Review 10 Years On*. London: Institute of Health Equity.
- McDonagh, T. A., et al. (2021) '2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure', *European Heart Journal*, 42(36), pp. 3599-3726.
- Moss, A. J., et al. (2009) 'Cardiac-resynchronization therapy for the prevention of heart-failure events', *New England Journal of Medicine*, 361(14), pp. 1329-1338.
- Mosterd, A. and Hoes, A. W. (2007) 'Clinical epidemiology of heart failure', *Heart*, 93(9), pp. 1137-1146.
- National Health Service (NHS) (2019) *The NHS Long Term Plan*. London: NHS England.
- National Institute for Cardiovascular Outcomes Research (NICOR) (2025) *National Heart Failure Audit Annual Report*. London: NICOR.
- National Institute for Health and Care Excellence (NICE) (2018) *Chronic heart failure in adults: diagnosis and management* (NICE Guideline NG106). London: NICE.
- Packer, M., et al. (2020) 'Cardiovascular and renal outcomes with empagliflozin in heart failure', *New England Journal of Medicine*, 383(15), pp. 1413-1424.
- Piepoli, M. F., et al. (2006) 'Peripheral muscle disuse and inflammation in chronic heart failure', *European Heart Journal*, 27(21), pp. 2588-2597.

References

PAGE 12

- Ponikowski, P., et al. (2023) 'Global epidemiology of heart failure: A contemporary perspective', *Nature Reviews Cardiology*, 20(4), pp. 235-251.
- Rao, S., et al. (2020) 'Socioeconomic status and heart failure outcomes in the UK', *Heart*, 106(12), pp. 901-908.
- Roger, V. L. (2021) 'Epidemiology of heart failure: A contemporary perspective', *Circulation Research*, 128(10), pp. 1421-1436.
- Royal College of Nursing (RCN) (2023) *Specialist Nursing Workforce in the UK: Heart Failure Services Audit*. London: RCN.
- Rutledge, T., et al. (2006) 'Depression in heart failure: A meta-analytic review of prevalence, intervention effects, and associations with clinical outcomes', *Journal of the American College of Cardiology*, 48(8), pp. 1527-1533.
- Savarese, G., et al. (2023) 'Global burden of heart failure: A comprehensive update', *Cardiovascular Research*, 119(17), pp. 3272-3284.
- Schulz, R. and Sherwood, P. R. (2008) 'Physical and mental health effects of family caregiving', *American Journal of Nursing*, 108(9), pp. 23-27.
- Shah, S. J., et al. (2020) 'Phenotypic mapping of heart failure with preserved ejection fraction', *Circulation*, 141(12), pp. 1007-1032.
- Stewart, S., et al. (2003) 'More 'malignant' than cancer? Five-year survival rates in patients with heart failure in the UK', *European Journal of Heart Failure*, 5(3), pp. 315-325.
- Taylor, C. J., et al. (2022) 'Trends in incidence and prevalence of heart failure in primary care: A 20-year population-based study in the UK', *British Journal of General Practice*, 72(716), pp. e234-e242.
- Taylor, R. S., et al. (2014) 'Exercise-based rehabilitation for heart failure', *Cochrane Database of Systematic Reviews*, (4), art. no. CD003331.
- Virani, S. S., et al. (2020) 'Heart disease and stroke statistics—2020 update: A report from the American Heart Association', *Circulation*, 141(9), pp. e139-e596.
- Wagner, E. H., et al. (2001) 'Quality improvement in chronic illness care: A collaborative approach', *Joint Commission Journal on Quality Improvement*, 27(2), pp. 63-80.