

Name: Jozi Daniel

Student ID: 2521417

Module Code: LCHS4008

Module: Introduction to Public Health

Lecturer: Dr Betsy Jose

Submitted on: 21st July 2026

Depression: Measuring, Monitoring and Supporting Individuals in the Community

Introduction:

Depression is a common mental health condition that affects how a person thinks, feels and behaves. It is more than feeling sad or low in mood, as it can last for weeks, months or even longer and significantly impact daily life. Depression can affect relationships, work, education and a person's ability to carry out everyday activities. It is considered one of the leading causes of disability worldwide. Common symptoms include persistent low mood, loss of interest in activities, tiredness, sleep problems, changes in appetite, poor concentration and feelings of worthlessness or guilt. Depression can affect anyone, but certain factors increase the risk such as stressful life events, trauma, genetics and long-term health conditions.

Globally, depression affects over 280 million people, making it one of the most widespread mental health disorders. In the UK, around 1 in 6 adults experience symptoms of depression at any one time. This report will explore how depression is measured, monitored and managed within health and social care services, along with its impact on individuals, communities and the NHS.

Approaches used to measure and monitor depression:

Depression is measured and monitored using a combination of clinical tools, patient feedback and national data collection systems. In the UK, one of the most commonly used tool is the **Patient Health Questionnaire (PHQ-9)**. This is a short questionnaire that asks individuals about their mood, sleep, energy levels, appetite and thoughts of self-harm. It helps healthcare professionals understand how severe the depression is and whether symptoms are improving over time.

Another tool used is the **Hospital Anxiety and Depression Scale (HADS)**, which is often used in hospitals to identify anxiety and depression in patients who may also have physical health conditions. These tools are useful because they allow professionals to track changes in mental health and adjust treatment if needed.

In addition to screening tools, depression is monitored through GP records and referrals to mental health services such as **NHS Talking Therapies (formerly IAPT)**. GPs play an important role because they are often the first point of contact. They assess symptoms, rule out physical causes and decide whether a referral or treatment is needed. On a national level, the UK uses data from systems such as the **NHS Digital Mental Health Services Dataset**, which collects information about referrals, diagnoses, waiting times and treatment outcomes. This helps the NHS understand demand and whether services are meeting people's needs.

Large population surveys like the **Adult Psychiatric Morbidity Survey (APMS)** are also used to measure how common depression is across different groups of people. These surveys help identify trends, such as higher rates in young adults or people experiencing financial hardship.

Monitoring depression is important because it helps ensure people receive the right treatment, reduces the risk of symptoms getting worse, and supports long-term management of the condition. It also helps healthcare services plan resources more effectively.

Consequences of depression for individuals, community and NHS:

Depression can have a serious impact on individuals. It can affect a person's ability to function in everyday life, including basic activities such as getting out of bed, eating properly, washing, or going to work or college. Many people also experience social withdrawal, where they avoid friends and family, which can increase feelings of isolation. In more severe cases, individuals may experience self-harm or suicidal thoughts, which requires urgent support from healthcare services. Depression can also affect physical health, as it is linked with fatigue, sleep disruption and reduced immune function.

For families and carers, depression can be emotionally difficult and sometimes stressful. Loved ones may not always understand the condition, which can lead to communication difficulties or strain in relationships. Children in families affected by depression may also be emotionally impacted if a parent or caregiver is struggling.

In the wider community, depression can reduce workplace productivity due to sickness absence and reduced performance at work. This has an economic impact on employers and the wider economy.

For the NHS, depression is one of the most common mental health conditions treated in primary care. It places pressure on GP services, mental health teams and crisis services. Many individuals require long-term support, which increases demand on already stretched services. It is also linked with higher use of emergency departments, especially during mental health crises.

The economic cost of depression in the UK is significant due to healthcare costs and lost productivity. This shows why early intervention and effective treatment are so important.

How depression is controlled and managed:

Depression is managed using a combination of psychological, medical and social approaches. One of the main treatments is **Cognitive Behavioural Therapy (CBT)**, which helps individuals identify negative thinking patterns and replace them with more positive and realistic thoughts. CBT is widely used in NHS Talking Therapies and has been shown to be effective for mild to moderate depression.

Medication is another common treatment option. **Antidepressants**, such as selective serotonin reuptake inhibitors (SSRIs), are often prescribed to help regulate mood. These are usually used alongside therapy rather than on their own.

Lifestyle changes can also support recovery. This may include regular exercise, improved sleep routines, healthy eating and reducing alcohol intake. While these do not replace treatment, they can improve overall wellbeing.

Early intervention is very important in controlling depression. If symptoms are identified early, treatment can begin sooner, which reduces the risk of the condition becoming more severe or long-term. Public health strategies also play a role. Awareness campaigns help reduce stigma around mental health, encouraging more people to seek help. Schools, workplaces and community organisations may also provide mental health education and support programmes.

Health and social care support services:

A wide range of services is available in the UK to support people with depression. The first point of contact is usually a **GP**, who can assess symptoms and refer individuals for treatment or prescribe medication.

NHS Talking Therapies provides free psychological treatments such as CBT, counselling and guided self-help. These services are designed to support people with mild to moderate depression.

For individuals with more severe or complex needs, **Community Mental Health Teams (CMHTs)** provide specialist support. These teams may include psychiatrists, mental health nurses, psychologists and social workers.

Crisis support services are also available for people experiencing severe distress or suicidal thoughts. This includes NHS urgent mental health helplines, A&E services and crisis intervention teams. Charities such as **Mind**, **Rethink Mental Illness**, and **Samaritans (116 123)** offer emotional support, advice and 24/7 listening services. These organisations are important because they provide support outside of NHS services.

Social care services may also be involved if depression affects daily functioning. This can include support with housing, benefits, employment or independent living.

Conclusion:

Depression is a serious and common mental health condition that affects millions of people in the UK and worldwide. It can significantly impact a person's emotional wellbeing, physical health and ability to carry out daily activities. The condition also affects families, communities and places pressure on health services such as the NHS.

Depression is measured and monitored using tools like PHQ-9 and HADS, as well as national data systems and surveys. These systems help healthcare professionals understand how common depression is and ensure that services are meeting patient needs. Although depression can be long-term and challenging, it can be effectively managed through therapies such as CBT, medication, lifestyle changes and social support. Early intervention and access to support services are essential for improving outcomes. Overall, a wide range of health and social care services exist in the UK to support individuals living with depression, but continued awareness and investment are important to reduce the impact of this condition.

Reference List:

American Psychiatric Association (2022) *Diagnostic and statistical manual of mental disorders (DSM-5-TR)*. 5th edn. Washington, DC: American Psychiatric Publishing.

British Medical Association (2023) *Mental health and wellbeing in the UK*. Available at: <https://www.bma.org.uk> (Accessed: 5 July 2026).

Department of Health and Social Care (2023) *Mental health and wellbeing plan*. Available at: <https://www.gov.uk/government/organisations/department-of-health-and-social-care> (Accessed: 5 July 2026).

Mental Health Foundation (2023) *Mental health statistics: depression in the UK*. Available at: <https://www.mentalhealth.org.uk> (Accessed: 5 July 2026).

Mind (2024) *Depression*. Available at: <https://www.mind.org.uk> (Accessed: 5 July 2026).

NHS (2024) *Depression overview*. Available at: <https://www.nhs.uk/mental-health/conditions/depression/> (Accessed: 5 July 2026).

NHS Digital (2023) *Mental Health Services Monthly Statistics*. Available at: <https://digital.nhs.uk> (Accessed: 5 July 2026).

Public Health England (2022) *Common mental health disorders: prevalence and data*. Available at: <https://www.gov.uk/government/organisations/public-health-england> (Accessed: 5 July 2026).

Royal College of Psychiatrists (2023) *Depression in adults*. Available at: <https://www.rcpsych.ac.uk> (Accessed: 5 July 2026).

World Health Organization (2023) *Depression fact sheet*. Available at: <https://www.who.int/news-room/fact-sheets/detail/depression> (Accessed: 5 July 2026).

