

Cover Page

Student's Name: Segun Victor Ikudehinbu

Student ID: 2120079

Module Name: Introduction to Public Health

Module Code: LCHS4004

Lecturer Name: Bianca Sanfilippo

Dementia.

Introduction

Dementia is a collective name for progressive neurological disorders resulting from damage to the brain. Dementia can be caused by Alzheimer's disease, vascular dementia, and other conditions such as Parkinson's Disease or Huntington's Disease. Dementia affects an individual in

many ways; physical abilities such as mobility, speech, and memory are all affected, as well as emotional and social changes. Dementia is not a natural part of aging, but can begin to occur when individuals are in their 60s or 70s.

Dementia is a serious health concern worldwide, with an estimated 55 million people living with dementia across the world (WHO, 2022). Dementia affects individuals of all ages and is expected to increase as the population continues to age. In the UK, it is estimated that 850,000 people are currently living with dementia and this number is anticipated to rise to 1 million by 2025 (Public Health England, 2022). Dementia costs the UK economy £26 billion per year in healthcare and social care services (Alzheimer's Society, 2014). It is also estimated that by 2051, there will be over 2 million people affected by Dementia in the UK alone (Bowen et al., 2016). This demonstrates how important it is for measures to be taken in order to measure, monitor, and control the incidence of Dementia in communities. Therefore, dementia is a serious health concern worldwide, and its estimated levels of incidence demonstrate the severity of this issue (WHO, 2022). It is essential that approaches and strategies are implemented in order to measure, monitor, and control Dementia in communities. This will help to reduce the impact of Dementia on individuals and the community as a whole. The paper will discuss public health measures available, the key determinants/risk factors of dementia, Short-term and long-term public health consequences of the dementia in adult social care, and provide a conclusion.

Public health measures available.

Public health measures can be used to help identify and control the incidence of dementia in communities (Livingston et al., 2020). Dementia is an umbrella term for various age-related brain diseases, such as Alzheimer's and vascular dementia, that affect memory, thinking and behavior. Dementia is a progressive condition which typically affects older individuals but can

occur at any age (CDC, 2019). Dementia presents with both physical and psychological symptoms which can have a significant impact on the individual's quality of life, including cognitive impairment, impaired motor functions and impaired communication skills (Cerejeira et al., 2012).

There are different approaches and strategies used to measure, monitor, and control the incidence of dementia in communities (Livingston et al., 2020). For instance, establishing and monitoring the prevalence of dementia over time is critical in assessing impact and progress of interventions. To effectively measure and monitor the impact of interventions, it is important to collect comprehensive data on dementia prevalence, incidence, and other factors that may affect outcomes (Mangialasche et al., 2012). Evaluating interventions is essential for understanding the effectiveness of dementia prevention and control measures. Moreover, establishing partnerships between public health authorities, research communities, and other stakeholders is an important step in ensuring the effectiveness of interventions (National Academies of Sciences, Engineering, and Medicine et al., 2017).

The public health approaches to controlling dementia in communities include early diagnosis through screening programs; provision of appropriate education and training for healthcare professionals; increasing awareness among the general public about dementia; encouraging people to seek medical advice if they have any concerns regarding their cognitive health; and providing support services to people living with dementia and their family members (Surr et al., 2017).

Early diagnosis is essential in order to provide appropriate treatment, care and support for individuals with dementia (Robinson et al., 2015). Dementia screening programs are available where healthcare professionals can assess the individual's cognitive abilities, such as memory and language skills, in order to diagnose whether an individual may be suffering from dementia.

Screening can also detect other conditions that may cause similar symptoms, such as depression or drug/alcohol abuse, which can then be treated accordingly.

Provision of education and training for healthcare professionals is important in order to ensure that they are aware of the signs and symptoms of dementia and how best to provide support and care for those living with the condition (Livingston et al., 2020). Healthcare professionals need to be trained in how best to diagnose, assess and treat people with dementia, as well as providing advice on lifestyle modifications which may help reduce symptoms.

Increasing public awareness of dementia is also important in order to reduce stigma associated with the condition (Riley et al., 2014). This can be achieved through campaigns and educational materials that raise knowledge levels about dementia, as well as providing information about available support services for people living with dementia and their families.

Encouraging people to seek medical advice if they have any concerns regarding their cognitive health is also important, as this can facilitate early diagnosis and enable people to receive necessary treatment and support (Robinson et al., 2015). Furthermore, providing support services to individuals living with dementia and their families is essential in helping them cope with the challenges posed by the condition. Services such as home care, respite care and counseling can all be beneficial for those affected by dementia. Thus, a variety of public health measures are available to help reduce the incidence of dementia in communities.

The key determinants/risk factors of dementia.

The exact cause of dementia is unknown, however there are several risk factors which can increase an individual's likelihood of developing the condition. These include age; family history; lifestyle factors (such as diet and exercise); having certain medical conditions (e.g., stroke, head

trauma etc.); and use of tobacco, alcohol or other drugs. Dementia is usually more common in populations with a lower socioeconomic status, and research suggests that individuals who have had fewer educational opportunities are at higher risk for the condition (Ranson et al., 2021).

In addition to these specific risk factors, there is also evidence that suggests certain modifiable lifestyle behaviors can help reduce the likelihood of developing dementia (Ranson et al., 2021). These include engaging in regular physical exercise; eating healthy and balanced diet rich in vegetables, fruit and omega-3 fatty acids; maintaining good levels of social engagement; avoiding excessive alcohol use; not smoking; and engaging in mentally stimulating activities.

Therefore, by taking steps to reduce the risk factors associated with dementia and adopting a healthy lifestyle, individuals can reduce their risk of developing the condition (CDC, 2022). Furthermore, early detection and diagnosis is key to ensuring that individuals can access appropriate treatments and supports which are tailored to their needs.

As such, public health professionals play an important role in promoting good mental health and encouraging people to make positive changes in order to maintain cognitive functioning throughout life. Public health initiatives should focus on raising awareness of the risk factors associated with dementia as well as providing information about how best to reduce them (Heger et al., 2020). This will help ensure that individuals are aware of ways they can protect themselves against developing dementia so that they can continue living active and meaningful lives for longer. Thus, it is essential that society provides adequate support services for those affected by dementia and their families.

Short-term public health consequences of the dementia in adult social care.

The short-term public health consequences of dementia in a health or social care setting are far reaching and can be significant. Dementia can put an individual at risk of falls, as well as increasing their vulnerability to abuse. Dementia can also lead to an increased likelihood of hospitalization due to complications with mobility, memory impairment and confusion (Mick et al., 2015). Dementia also has profound effects on family members caring for the person living with the condition, who often have to make difficult decisions related to medical treatment and end-of-life care. For this reason, it is important that healthcare professionals are aware of the impact that dementia can have on both individuals and families and provide appropriate support services.

In addition to providing direct care, there is a need for increased public awareness in order to reduce stigma surrounding the condition and encourage people living with dementia and their families to seek help (Riley et al., 2014). Dementia-specific services should also be put in place to support those affected by the condition, including early assessment and diagnosis, education and training programs, medical treatments, family counselling, respite care and support groups.

Finally, it is essential that more research is conducted into the causes of dementia in order to increase the understanding of the condition and develop better treatment options (NHS Choices, 2019). This will ultimately improve outcomes for individuals living with dementia as well as helping to lessen the burden on healthcare systems.

Hence, dementia is a complex and distressing condition which impacts not only those diagnosed but also their families and caregivers. It is essential that healthcare professionals are properly trained in the diagnosis, assessment and treatment of dementia in order to provide appropriate care for those affected by the condition. In addition, it is important to increase public awareness around dementia and ensure that individuals have access to suitable services which can support them throughout their journey with the condition. Finally, further research into the causes

of dementia will be essential in order to improve our understanding of this debilitating condition and develop better methods for treating it.

Overall, it is clear that there are many short-term public health consequences associated with dementia; however, with increased knowledge and understanding, as well as improved access to supportive services and treatments, these risks can be minimized (Mangialasche et al., 2012). This will help ensure a better quality of life for those living with dementia, as well as enabling them to remain an active and participating member of their community.

Long-term public health consequences of dementia in adult social care.

The long-term public health consequences of dementia in adult social care are equally significant. Over time, individuals living with the condition often require increasingly complex and costly medical treatment and support services (Bennett et al., 2018). As dementia progresses, the individual's independence decreases, requiring more support from family members, friends and caregivers. This can put an immense strain on both physical and emotional resources. In addition, over time there is an increased risk of mental illness such as depression due to isolation associated with dementia.

In terms of direct healthcare costs, it has been estimated that the global cost of dementia will be approximately \$2 trillion by 2030; this includes direct hospital costs but also the indirect costs associated with providing long-term care for those affected by the condition (Nandi et al., 2022). Furthermore, as dementia progresses, individuals become more likely to require costly end-of-life care.

Overall, it is clear that the long-term public health consequences associated with dementia are significant. As such, it is critical that those affected by the condition have access to appropriate

support services and medical treatments in order to prevent or reduce these risks and improve quality of life for those living with dementia. In addition, increased public awareness of the condition will help to reduce stigma around dementia as well as encouraging people to seek help sooner rather than later (Kim et al., 2019). Finally, research into causes and better treatments for dementia will be essential in order to lessen the burden on healthcare systems and ensure a better quality of life for those living with the condition.

Conclusion.

In conclusion, it is evident that dementia has a significant impact on both the individual and society. The condition not only has considerable short-term public health consequences but also long-term implications for an individual's quality of life, their careers and healthcare systems. In order to minimize these risks, it is essential that those affected by dementia have access to suitable services and treatments which can enable them to maintain a better quality of life throughout their journey with the condition. In addition, increased public awareness around dementia will be key in helping to reduce its stigma as well as encouraging people to seek help sooner rather than later. Finally, further research into the causes of dementia must continue in order to improve our understanding of this debilitating condition and develop better methods for treating it. It is only with these combined efforts that we can hope to lessen the burden of dementia on individuals, families and healthcare systems across the world.

References

- Alzheimer's Society. (2014). *Dementia UK report*. Alzheimer's Society.
<https://www.alzheimers.org.uk/about-us/policy-and-influencing/dementia-uk-report>
- Bennett, H. Q., Norton, S., Bunn, F., Robinson, L., Rait, G., Goodman, C., Brayne, C., & Matthews, F. E. (2018). The impact of dementia on service use by individuals with a comorbid health condition: a comparison of two cross-sectional analyses conducted approximately 10 years apart. *BMC Medicine*, 16(1). <https://doi.org/10.1186/s12916-018-1105-8>

- Bowen, M., Edgar, D. F., Hancock, B., Haque, S., Shah, R., Buchanan, S., Iliffe, S., Maskell, S., Pickett, J., Taylor, J.-P., & O'Leary, N. (2016). Introduction. In *www.ncbi.nlm.nih.gov*. NIHR Journals Library. <https://www.ncbi.nlm.nih.gov/books/NBK374270/>
- CDC. (2019, April 5). *What Is Dementia?* Centers for Disease Control and Prevention. <https://www.cdc.gov/aging/dementia/index.html>
- CDC. (2022, September 13). *Reducing Risk of Alzheimer's Disease*. Centers for Disease Control and Prevention. <https://www.cdc.gov/aging/publications/features/reducing-risk-of-alzheimers-disease/index.htm>
- Cerejeira, J., Lagarto, L., & Mukaetova-Ladinska, E. B. (2012). Behavioral and Psychological Symptoms of Dementia. *Frontiers in Neurology*, 3(73). <https://doi.org/10.3389/fneur.2012.00073>
- Heger, I., Köhler, S., Boxtel, M. van, Vugt, M. de, Hajema, K., Verhey, F., & Deckers, K. (2020). Raising awareness for dementia risk reduction through a public health campaign: a pre-post study. *BMJ Open*, 10(11), e041211. <https://doi.org/10.1136/bmjopen-2020-041211>
- Kim, S., Werner, P., Richardson, A., & Anstey, K. J. (2019). Dementia Stigma Reduction (DESeRvE): Study protocol for a randomized controlled trial of an online intervention program to reduce dementia-related public stigma. *Contemporary Clinical Trials Communications*, 14, 100351. <https://doi.org/10.1016/j.conctc.2019.100351>
- Livingston, G., Huntley, J., Sommerlad, A., Ames, D., Ballard, C., Banerjee, S., Brayne, C., Burns, A., Cohen-Mansfield, J., Cooper, C., Costafreda, S. G., Dias, A., Fox, N., Gitlin, L. N., Howard, R., Kales, H. C., Kivimäki, M., Larson, E. B., Ogunniyi, A., & Orgeta, V. (2020). Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *The Lancet*, 396(10248), 413–446.
- Mangialasche, F., Kivipelto, M., Solomon, A., & Fratiglioni, L. (2012). Dementia prevention: current epidemiological evidence and future perspective. *Alzheimer's Research & Therapy*, 4(1), 6. <https://doi.org/10.1186/alzrt104>
- Mick, P., Reed, M., & Pichora-Fuller, M. (2015). Hearing, Cognition, and Healthy Aging: Social and Public Health Implications of the Links between Age-Related Declines in Hearing and Cognition. *Seminars in Hearing*, 36(03), 122–139. <https://doi.org/10.1055/s-0035-1555116>
- Nandi, A., Counts, N., Chen, S., Seligman, B., Tortorice, D., Vigo, D., & Bloom, D. E. (2022). Global and regional projections of the economic burden of Alzheimer's disease and related dementias from 2019 to 2050: A value of statistical life approach. *EClinicalMedicine*, 51, 101580. <https://doi.org/10.1016/j.eclinm.2022.101580>

National Academies of Sciences, Engineering, and Medicine, Health and Medicine Division, Board on Population Health and Public Health Practice, in, Baciú, A., Yamrot Negussie, Geller, A., & Weinstein, J. N. (2017, January 11). *Partners in Promoting Health Equity in Communities*. Nih.gov; National Academies Press (US).

<https://www.ncbi.nlm.nih.gov/books/NBK425859/>

NHS Choices. (2019). *Is there a cure for dementia? - Dementia guide*. NHS.

<https://www.nhs.uk/conditions/dementia/cure/>

Public Health England. (2022, July 1). *Dementia: Applying All Our Health*. GOV.UK.

<https://www.gov.uk/government/publications/dementia-applying-all-our-health/dementia-applying-all-our-health>

Ranson, J. M., Rittman, T., Hayat, S., Brayne, C., Jessen, F., Blennow, K., Duijn, C. van, Barkhof, F., Tang, E., Mummery, C. J., Stephan, B. C. M., Altomare, D., Frisoni, G. B., Ribaldi, F., José, Molinuevo, L., Scheltens, P., Llewellyn, D. J., Abramowicz, M., & Berthier, M. (2021). Modifiable risk factors for dementia and dementia risk profiling. A user manual for Brain Health Services--part 2 of 6. *Alzheimer's Research & Therapy*, 13(1), NA–NA. <https://doi.org/10.1186/s13195-021-00895-4>

Riley, R. J., Burgener, S., & Buckwalter, K. C. (2014). Anxiety and Stigma in Dementia: A Threat to Aging in Place. *The Nursing Clinics of North America*, 49(2), 213–231.

<https://doi.org/10.1016/j.cnur.2014.02.008>

Robinson, L., Tang, E., & Taylor, J.-P. . (2015). Dementia: timely diagnosis and early intervention. *BMJ*, 350(jun15 14), h3029–h3029. <https://doi.org/10.1136/bmj.h3029>

Surr, C. A., Gates, C., Irving, D., Oyebode, J., Smith, S. J., Parveen, S., Drury, M., & Dennison, A. (2017). Effective Dementia Education and Training for the Health and Social Care Workforce: A Systematic Review of the Literature. *Review of Educational Research*, 87(5), 966–1002. <https://doi.org/10.3102/0034654317723305>

WHO. (2022, September 20). *Dementia*. Wwww.who.int. <https://www.who.int/news-room/fact-sheets/detail/dementia#:~:text=Currently%20more%20than%2055%20million>