

The Role of Managing Hypertension as a Long-Term Condition

Programme: MSc Public Health and Social
Care in Practice

Module: Community Care and
Management of Longer-term Conditions

Module Code: LCHS7003

Lecturer: Dr Aishat Bakre

Student Name: Dil Afroze Khan

Student ID: 2508106

Contents

Part 1: The Problem	4
Part 2: The Impact	12
Case Study.....	12
Health Outcomes	13
Social and Economic Burden	15
Introducing Hypertension Policy and Practice.	17
Part 3: Managing Hypertension	20
Behavioral Approaches.....	20
Health and Therapeutic Services	21
Role of Technology	22
Part 4: Theory and Policy background.....	23
NHS and NICE Guidelines	23
International Frameworks	25
Part 5: Person-Centred Recommendations	27

Self-Care Strategies	27
Part 6: Future Directions	30
Conclusion	32
References	33

Part 1: The Problem

What is hypertension?



Figure 1 hypertension

Sources: <https://www.google.com/url?>

One of the most prevalent chronic conditions in the world among adults is hypertension, or the continued high blood pressure. It is a major contributor of cardiovascular morbidity as well as stroke or premature deaths (WHO, 2023). The NHS (2024) reports that approximately every fourth adult in the UK has a hypertension diagnosis and not all of them are properly diagnosed and managed.

It is also known as the silent killer as it may develop without any symptoms until the development of complications, including heart failure, kidney disease, and stroke (Riemann et al., 2023).

Treatment and management of hypertension condition is intimately connected with lifestyle and behavioural variables: diet and physical inactivity, stress, and, in particular, poor sleep hygiene (Buysse, 2014; Van Cauter et al., 2008). Poor or interrupted sleep has been reported to higher blood pressure, more active sympathetic nervous system, and poor metabolic control (Spiegel et al., 2005; Johnson et al., 2018). On the contrary, because sleep-wake schedules, low levels of stimulants, and an absence of disturbance in the sleeping environment are known as good sleep hygiene, they can positively affect blood pressure levels and

decrease cardiovascular workload (Irish et al., 2015).

The sleep hygiene is not, therefore, a preventive and therapeutic behaviour, but a long-term condition in itself, which is necessary in the prevention and treatment of hypertension and other cardiovascular risks. One of the significant health challenges among the population of the UK and the world as a whole is hypertension. It charges approximately 12 percent of the total GP consultations and is the cause of over half of all cardiovascular incidents (NHS, 2024). The risk is further aggravated by environmental stressors, work fatigue, and low quality of sleep-in shift workers, carers, and people who live in deprived regions (Johnson et al., 2018). The method of incorporating sleep hygiene into hypertension management will contribute to a decrease in resistance to treatment,

dependence on medications, and a decrease in the NHS load (Ruan et al., 2025).

Why It Is a Public Health Concern




Figure 2 Public Health

Sources: <https://www.google.com/url?>

Hypertension as a Community Health Issue. Hypertension has been noted to be a significant health issue affecting the general population in the United Kingdom and any other part of the world. It causes a significant burden of non-communicable diseases and is closely associated with unhealthy lifestyle habits, such as poor sleep, excessive salt consumption, obesity and inactive lifestyles

(Insel, 2006; Liu, 2016). The regulation of blood pressure is made worse by poor sleep hygiene, which is especially disruptive to circadian rhythms and activity of stress hormones. This relationship between sleep and hypertension underscores the need to have preventive public health interventions against the two conditions.

In the UK, hypertension is diagnosed in one in three adults, and a significant number of these patients are undiagnosed or not managed (NICE, 2023). This growing prevalence puts the NHS at a huge burden in terms of consultations, medication prescriptions, and referrals to cardiovascular and renal complications associated with the condition (NHS, 2024). It is also significant economically, with RAND Europe estimating that work productivity losses due to hypertension and related sleep deprivation have cost the UK economy billions of any given year in the form of absenteeism,

presenteeism, and medical spending (Irish et al., 2015).

Hypertension is equally disadvantaged in terms of vulnerable groups, such as shift workers, carers, and community members in deprived communities, who have chronic stresses and inadequate access to healthcare services (Johnson et al., 2018). The blood pressure increase in these groups is aggravated by environmental stressors, abnormal work schedules and poor sleep habits. In risky professions like healthcare, transport and construction, the uncontrolled hypertension levels pose risks of accidents and long-term disability (O'Donnell, 2010).

Population level preventive measure to hypertension is to incorporate preventive measures in the population health policy. Combination of sleep hygiene knowledge with known lifestyle modification programmes aimed at diet, physical activity

and stress management should be used as interventions. The NHS and the World Health Organization (WHO) support community-based and educational programs as effective ways of morbidity reduction and cardiovascular health outcomes (Ruan et al., 2025).

Prevalence and Burden in the UK & Globally

Adults worldwide have had to report some sleep disturbance as up to 40% and chronic insomnia as 10% (Buysse, 2014). The NHS statistics indicate that in the UK, one in every three adults has difficulty sleeping, and the prevalence of the problem is greater among women, older adults, and people with long-term conditions (NHS Inform, 2024). This problem was boosted by the COVID-19 pandemic, and a form of coronasomnia became a common phenomenon (McAlpine et al., 2024). It is a heavy economic burden. The UK economy spends about £40 billion

every year because of lost productivity as a result of poor sleep (Irish et al., 2015). The global issue is when in the US alone it is estimated that over 400 billion have been lost (Liu, 2016). In addition to the economic, bad sleep disorder affects learning, relationships, and the quality of life (Harvey, 2014).

The same tendencies are witnessed internationally. A study in Egypt revealed that poor sleep hygiene in medical students is very high, which reduces academic performance by a large margin (Talaia et al., 2025). These results prove that sleep hygiene is not just a western agenda but a worldwide health issue of concern.

Part 2: The Impact

Case Study

Mr A is a 42-year-old office worker who complained about constant fatigue, mood swings, and lack of concentration. He was a regular consumer of caffeine towards the end of the day, an electronic devices bed time user, and had irregular sleep patterns. Gradually his blood pressure increased, causing hypertension, and his weight increased as well as he experienced symptoms of anxiety. He lost his working performance and his family relationships became dysfunctional. It is an example of how inadequate sleep hygiene may contribute to worsening hypertension to start the cycle of physical disease, mental distress, and social disturbance (Harvey, 2014).

Health Outcomes

Insomnia has already been proven to be a causal factor of hypertension and other cardiovascular diseases. The chronic loss of sleep interferes with the endocrine and metabolic systems, increasing the cortisol level and the sympathetic nervous system, which increases blood pressure (Van Cauter et al., 2008). Poor sleep is also a cause of insulin resistance and obesity which are two key contributors to high blood pressure.

Also, the lower immune system exposes people to infection (Buysse, 2014). Sleep deprivation causes anxiety, irritability, and mood swings, which increase blood pressure due to stress (Blackwelder et al., 2021). This builds a two-way connection: lack of mental health deteriorates the quality of sleep, and the disruption of sleep also disrupts the emotional regulation (McAlpine et al., 2024).

Cognitive flexibility and slower response time also decrease sleep deprivation, which influences decision-making and productivity. Research indicates that long-term sleep deprivation causes the disconnection of the brain and leads to the loss of problem-solving and memory (Meisel et al., 2017; Lauharatanahirun et al., 2020). In the long term, insomnia leads to the rise of neurodegenerative diseases and chronic hypertension (Irish et al., 2015).

All in all, these results highlight that the enhancement of sleep hygiene is an important non-pharmacological approach to hypertension management and overall health.

Social and Economic Burden




Figure 3 Social and Economic Burden

sources: <https://www.google.com/url?>

Sleep deprivation and lack of control in high blood pressure are very expensive to individuals, employers and the healthcare system. Fatigue causes low efficiency at the workplace and high accident rates in high stakes workplaces like transport, construction and healthcare (O'Donnell, 2010). RAND Europe states that billions of pounds are lost a year through absenteeism and presenteeism related to sleep

deprivation and hypertension (Irish et al., 2015). In the NHS, sleep deprivation contributes to the worsening of hypertension-related complications, which in turn augments the GP rates, prescription rates, and hospitalization (Riemann et al., 2023). Indirect stress is also encountered by the families where the carer is taking care of hypertensive patients with sleep disruptions resulting in emotional strain (Buman, 2011). The occurrence of hypertension and sleep deprivation across the globe reflects the identical socioeconomic trends, as these issues are more common among the workers of the shift, low-income, and city-dwellers subjected to the chronic stress and poor environment (Johnson et al., 2018). This highlights the importance of combined sleep and hypertension prevention of the public health strategy.

Introducing Hypertension Policy and Practice.

Hypertension is among the most widespread chronic diseases in the world, and it is the cause of almost 13 percent of all deaths in the world (WHO, 2024). Although it can be easily prevented, it is usually not diagnosed until major cardiovascular complications arise. Sleep hygiene is another potential preventive force that is underexploited.

Studies continue to identify that poor or insufficient sleep leads to chronic stimulation of the sympathetic nervous system, elevated blood pressure at night, and decreased vascular recovery (Buysse, 2014; Van Cauter et al., 2008). In such a way, the health impacts of the inclusion of sleep education in the national strategies to address hypertension may be extensive.

In the UK, inclusion of sleep health in the management of hypertension supports the

reduction of cardiovascular risk by lifestyle modification in the NHS Long-Term Plan (Epstein, 2025). Inclusion of sleep hygiene assessment in primary care would enable general practitioners detect poor sleep as a risk factor early like the screening of smoking or dietary screening. This active management would allow minimizing the use of antihypertensive drugs and might save the NHS money, with hypertension-related treatment already costing it billions per year (NHS, 2024).

Another opportunity is a scaled opportunity offered by digital health interventions. Sleepio, which has been NHS approved and other CBT-I apps may be modified to incorporate blood pressure data, and therefore provide real-time information on the relationship between sleep habits and hypertension results (McAlpine et al., 2024). Those tools can give patients the power to

self-manage with personalized reminders, behaviour tracking and education modules that make the relationship between better sleep and cardiovascular health. Notably, policymakers should make sure that such technologies can be available to all social economic groups to avoid digital health inequity (Ruan et al., 2025).

On the international level, other nations like Japan and Finland have already initiated national sleep awareness to fight hypertension and metabolic diseases (Talaia et al., 2025). These programs show that prevention of fatigue at the workplace, public education and health programs in school may all help improve cardiovascular outcomes. This scheme applied to the UK with the creation of similar schemes especially in high-risk work such as the healthcare, logistics and manufacturing would not only enhance productivity, but

also minimize the levels of accidents and absenteeism (O'Donnell, 2010).

Part 3: Managing Hypertension

Behavioral Approaches

Mobile app-based lifestyle modification to overcome clinical inertia




Figure 4 Behavioral Approaches

Sources: <https://www.google.com/url?>

Hypnotic management of hypertension requires sleep hygiene interventions. Regular sleep and wakefulness, avoiding caffeine and screen time before bed, and a cool, dark, and quiet atmosphere make

restorative sleep (Buysse, 2014; Irish et al., 2015). The regulation of behaviour stabilizes circadian rhythms and reduces changing blood pressure.

The most evidence based one is Cognitive Behavioural Therapy of Insomnia (CBT-I), which involves the integration of behaviours and cognitive therapies including stimulus restriction, sleep restriction, and relaxation training (Edinger and Means, 2005; Perlis et al., 2005). These interventions are common throughout the NHS where they serve as alternative cost-effective means of long-term medication prescription.

Sleep hygiene should be used in hypertension management plans as it can better treat hypertension, regulate blood pressure, and increase the quality of life.

Health and Therapeutic Services

The lack of effectiveness of behavioural measures can be treated by short-term use

of pharmacological interventions (melatonin, low doses of hypnotics) but only with close supervision (Riemann et al., 2023). It is not encouraged because of the risks of dependency in the long run. Referrals to sleep clinics should be made to patients with hypertension whose sleeping problems include sleep apnea (Redline, 2010). Other complementary therapies, including acupuncture or herbal supplements, are used on an occasional basis but have a low level of evidence (Varadharasu et al., 2024).

Role of Technology

Clinical sleep management is now accompanied by digital interventions. CBT-I remote access services (such as Sleepio, which is approved by the NHS) are shown to positively influence sleeping behavior in hypertensive patients (McAlpine et al., 2024). Wearables and smartphone trackers help develop a better level of self-awareness

about sleep behaviour, whereas telehealth consultations expand the reach to patients in remote and resource-limited regions (Ruan et al., 2025). The combination of these technologies with the hypertension care contributes to the regular monitoring of the lifestyle and decreases the burden of the primary healthcare services.

Part 4: Theory and Policy background

NHS and NICE Guidelines




Figure 5 NHS and NICE Guidelines

Sources: <https://www.google.com/url?>

The NHS Long-Term Plan lists sleep health as one of the key lifestyle determinants of preventive and management of long-term conditions, such as hypertension (Epstein, 2025). Sleep is currently included as a behavioral determinant of chronic disease alongside diet and physical activity. NHS policy promotes early identification of sleep issues in GP appointments and the non-pharmacological treatment with the CBT-I and education of a patient.



Figure 6 NICE

Sources:

<https://www.google.com/url?sa=i&url=https%3A%2F%2Fwww.worcestershire.gov.uk%2F>

NICE guidelines (NG136) on hypertension emphasize lifestyle change as a first-line

intervention hypertension, and the quality of sleep, physical activities, and diet play a crucial role (NICE, 2023). CBT-I is also advised by NICE as a gold standard in the treatment of insomnia (Riemann et al., 2023). These models dishearten the hypnotic consumption in the long term and advance preventive and sustainable health programs.

Collectively, NHS and NICE introduce sleep hygiene as a hypertension management pillar, which enhances the ability to control blood pressure and healthcare expenditure.

International Frameworks

The World Health Organization (WHO) lists the lack of sleep as one of the risk factors of non-communicable diseases, including hypertension, diabetes, and cardiovascular disorders (Ruan et al., 2025). International regulations highlight population-based interventions, i.e., public education,

workplace health initiatives, and policy combination, to make sleep health resources equally accessible.

The American Academy of Sleep Medicine (AASM) is a world-renowned clinical direction, where behavioural therapy, such as CBT-I and multidisciplinary treatment, is recommended with the participation of physicians, psychologists, and sleep specialists (Buysse, 2014).

The combined efforts of WHO and AASM uses propose universal access to evidence-based culturally aware practices of sleep and hypertension management.

Part 5: Person-Centred Recommendations

Self-Care Strategies

Self-care practices are key to good management of individuals with hypertension. Sleep diaries can be used to educate patients to monitor bedtime habits, caffeine use, and sleep quality, which can be used to create awareness of the habits that impact BP (Irish et al., 2015). Physical activity, a healthy diet, and restful activities including meditation and breathing exercises are an addition to medical care and enhance general heart strength (Varadharasu, Kumar and Senthamarai, 2024).

The digital CBT-I platforms including Sleepio and Sleepier and NHS Trusts enable patients via education and remote access (Ruan et al., 2025). Sleep hygiene as a part of community care represents an effective

prevention approach that facilitates patient self-management of hypertension in a cost-effective way.

Carer Involvement



Figure 7 Carer Involvement

Sources: <https://www.google.com/url?>

Carers are very important in helping hypertensive patients and especially the old or patients with comorbidities. They are able to assist in creating regular routines, minimize

night interference, and guarantee drug compliance (Buman, 2011). In patients with mobility problems or cognitive impairments, involvement of the carers decreases anxiety,

improves treatment effects, and eliminates blood pressure peaks in the course of sleep (Redline, 2010).

Preventive Programmes

The reduction of the dual burden of poor sleep and hypertension depends on community-based prevention. The awareness of healthy sleep and blood pressure control is created through public health campaigns arranged by the NHS and local authorities (NHS Inform, 2024). Fatigue-related hypertension is eliminated through flexible hours and rest breaks at the workplace promoted by programmes (McAlpine, Mullan and Clarke, 2024). School education encourages life-long habits, which lower the risks of hypertension in the future (Harvey, 2014).

Together, preventive interventions focus on making lifestyle education an element of everyday existence and lowering the cost-levels as well as the healthcare costs.

Part 6: Future Directions

Research Gaps

Compared to CBT-I, the body of evidence regarding the CBT-I today has effectively helped to improve sleep, but further investigations are required regarding the long-term impacts of CBT-I on the management of hypertension (McAlpine, Mullan and Clarke, 2024). The effectiveness of digital interventions such as mobile CBT-I applications should also be evaluated in the context of their long-term efficacy, access, and safety. Moreover, it is necessary to conduct cross-cultural studies to learn the effect of socioeconomic variables on the adherence to the guidelines of sleep and hypertension management (Talaia et al., 2025).

Policy Recommendations

Sleep hygiene must be incorporated into screening of primary care in order to

enhance management of hypertension. The assessment of sleep quality made routinely in GP consultations can allow timely intervention and limit the use of medication (Ruan et al., 2025).

The physical access to digital tools of CBT-I and NHS-approved apps would be improved by scaling these tools on a national level, particularly in the underserved rural or low-income settings. Strong clinical evaluation and data security guidelines should be supported to ensure such initiatives are supported by policymakers.

Internationally, working with WHO systems will help coordinate cross-border policies to curb hypertension and support cardiovascular health across the world (Ruan et al., 2025).

Conclusion

Sleep hygiene is an evidence-based intervention, which is cost effective and is significant in prevention and management of hypertension. One can manage blood pressure, mental health, and even does not need to take medication with the assistance of better sleep. CBT-I and other behavioral interventions are based on guidelines of NHS and NICE and provide the patient with sustainable care. A complex intervention of sleep hygiene and lifestyle-modification, digital health solutions, and international frameworks is a complicated way of managing hypertension and improving the resilience of the general population health.

References

- Blackwelder, A., Hoskins, M. and Huber, L. (2021) "Effect of inadequate sleep on frequent mental distress," *Preventing chronic disease*, 18(200573), p. E61. Available at: <https://doi.org/10.5888/pcd18.200573>.
- Bu, Z. (2025) "Effects of various exercise interventions in insomnia patients: a comparative study," *BMJ Evidence-Based Medicine* [Preprint].
- Buman, M.P. (2011) "Sleep and Daily Functioning during a Naturalistic Stressor: The Case of Natural Disasters," *Journal of Behavioral Medicine*, 34(5), pp. 348–358.
- Buyssse, D.J. (2014) "Sleep health: can we define it? Does it matter?" *Sleep*, 37(1), pp. 9–17. Available at: <https://doi.org/10.5665/sleep.3298>.
- Edinger, J.D. and Means, M.K. (2005) "Cognitive-behavioral therapy for primary

insomnia,” *Clinical psychology review*, 25(5), pp. 539–558. Available at: <https://doi.org/10.1016/j.cpr.2005.04.003>.

Epstein, L. (2025) *Effects of sleep deprivation*, *Harvard Health*. Available at: <https://www.health.harvard.edu/staying-healthy/effects-of-sleep-deprivation>

Fucci, D. *et al.* (2020) “Need for sleep: The impact of a night of sleep deprivation on novice developers’ performance,” *IEEE transactions on software engineering*, 46(1), pp. 1–19. Available at: <https://doi.org/10.1109/tse.2018.2834900>.

Harvey, A.G. (2014) “The development and evaluation of a sleep education intervention to improve sleep for people with psychosis,” *Behaviour Research and Therapy*, 55, pp. 58–66.

Insel, B.J. (2006) “Sleep Duration and Incident Diabetes in Postmenopausal Women,” *Diabetes Care*, 29(2), pp. 293–297.

Irish, L.A. *et al.* (2015) "The role of sleep hygiene in promoting public health: A review of empirical evidence," *Sleep medicine reviews*, 22, pp. 23–36. Available at: <https://doi.org/10.1016/j.smr.2014.10.001>.

Johnson, D.A., Billings, M.E. and Hale, L. (2018) "Environmental determinants of insufficient sleep and sleep disorders: Implications for population health," *Current epidemiology reports*, 5(2), pp. 61–69. Available at: <https://doi.org/10.1007/s40471-018-0139-y>.

Lauharatanahirun, N. *et al.* (2020) "Flexibility of brain regions during working memory curtails cognitive consequences to lack of sleep," *arXiv [q-bio.NC]*. Available at: <http://arxiv.org/abs/2009.07233>.

Liu, Y. (2016) "Sleep duration and risk of stroke: a dose-response meta-analysis of prospective studies," *International Journal of Stroke*, 11(3), pp. 307–316.

McAlpine, T., Mullan, B. and Clarke, P.J.F. (2024) "Re-considering the role of sleep hygiene behaviours in sleep: Associations between sleep hygiene, perceptions and sleep," *International journal of behavioral medicine*, 31(5), pp. 705–717. Available at: <https://doi.org/10.1007/s12529-023-10212-y>.

Meisel, C. *et al.* (2017) "Decline of long-range temporal correlations in the human brain during sustained wakefulness," *Scientific reports*, 7(1), p. 11825. Available at: <https://doi.org/10.1038/s41598-017-12140-w>.

Nielson, S., Dautovich, N. and Dzierzewski, J. (2025) "0166 religious involvement and sleep health in middle-aged and older adults," *Sleep*, 48(Supplement_1), pp. A75–A75. Available at: <https://doi.org/10.1093/sleep/zsaf090.0166>.

O'Donnell, E. (2010) *Effects of sleep loss can be masked, not erased, Harvard Magazine*. Harvard Magazine Inc. Available at:

<https://www.harvardmagazine.com/2010/07/lost-sleep-is-hard-to-find>

Perlis, M.L. et al. (2005) *Cognitive Behavioral Treatment of Insomnia: A Session-by-Session Guide*. New York: Springer.

Redline, S. (2010) "The effect of age, sex, race, and body mass index on sleep apnea severity," *American Journal of Respiratory and Critical Care Medicine*, 183(5), pp. 736–746.

Riemann, D. et al. (2023) "The European Insomnia Guideline: An update on the diagnosis and treatment of insomnia 2023," *Journal of sleep research*, 32(6), p. e14035. Available at:

<https://doi.org/10.1111/jsr.14035>.

Ruan, J.Y. *et al.* (2025) "Effects of sleep hygiene education for insomnia: A systematic review and meta-analysis," *Sleep medicine reviews*, 82(102109), p. 102109. Available at: <https://doi.org/10.1016/j.smrv.2025.102109>.
Sleep hygiene (2024) *NHS inform*. Available at: <https://www.nhsinform.scot/mind-to-mind/sleeping-better/sleep-hygiene>

Sleep hygiene (no date a) *Nhs.uk*. Available at: <https://www.grmc.nhs.uk/sleep-hygiene>
Sleep hygiene (no date b) *Kingston and Richmond NHS Foundation Trust*. Available at: <https://www.kingstonandrichmond.nhs.uk/patients-and-families/patient-leaflets/sleep-hygiene>

"Sleep Hygiene - Patient Information" (no date) *University Hospital Southampton NHS Trust* [Preprint].

Spiegel, K. *et al.* (2005) "Sleep loss: a novel risk factor for insulin resistance and Type 2 diabetes," *Journal of applied physiology* (Bethesda, Md.: 1985), 99(5), pp. 2008–2019. Available at: <https://doi.org/10.1152/jappphysiol.00660.2005>.

Talaia, A. *et al.* (2025) "Sleep hygiene practices and sleep quality among medical students in Egypt: a cross-sectional study," *Sleep science and practice*, 9(1). Available at: <https://doi.org/10.1186/s41606-025-00125-y>.

Van Cauter, E. *et al.* (2008) "Metabolic consequences of sleep and sleep loss," *Sleep medicine*, 9 Suppl 1, pp. S23-8. Available at: [https://doi.org/10.1016/S1389-9457\(08\)70013-3](https://doi.org/10.1016/S1389-9457(08)70013-3).

Varadharasu, S., Kumar, U. and Senthamarai, R. (2024) "Sleep hygiene efficacy on quality of sleep and mental ability

among insomnia clients,” *Sleep Medicine* [Preprint].

Why sleep matters: Consequences of sleep deficiency (no date) *Harvard.edu*. Available at: <https://sleep.hms.harvard.edu/education-training/public-education/sleep-and-health-education-program/sleep-health-education-45>