



Student name: FLAVIA APIO

Student ID: 2023550

Module name: Introduction to Physiology

Module code: Introduction to Physiology

Lecturer name: Olusegun Awolaran

Assessment: Written Report

Contents Table

Introduction	3
Anatomy and Physiology of the health condition	4&5
Health and Lifestyle routine measures	6
Health and Social Care practice.....	7
Recommendations	8
Conclusion.....	9
Reference List	10

STROKE

Introduction


This is a well-organized report about stroke. It will outline common health and lifestyle metrics for learning about how people with strokes' bodies work, including details on the anatomy and physiology connected to it.

Additionally, this study will give instances of health and social care policy, practise, and/or planning from the UK to illustrate how campaigns, health education, and other care-planning initiatives may be influenced by data and measurements on how the body functions. This data will be associated with stroke. Additionally, advice made to stroke victims who reside in the UK will be included.

Anatomy and Physiology

When blood flow is interrupted to a portion of the brain, a potentially fatal medical condition known as a stroke may develop. Strokes are a medical emergency that require prompt care. The likelihood of permanent harm decreases with time after receiving treatment for a stroke. If you think you're having a stroke or that someone else is, call 999 immediately and ask for an ambulance. (NHS, 2022)

It helps to understand how blood flows to the brain in order to comprehend stroke (see Anatomy of the Brain). Internal carotid arteries and vertebral arteries work together to pump blood from the heart to the brain. (Fig. 1). The anterior (front) portions of the brain are supplied by the internal carotid arteries, while the posterior (back) portions are supplied by the vertebral arteries. The left and right vertebral arteries combine to produce the basilar artery after passing through the skull. The Circle of Willis, a ring near the base of the brain, serves as a conduit for communication between the internal carotid arteries and the basilar artery. The artery that is most frequently closed during a stroke is the middle cerebral artery. (Mayfield Brain & Spine, 2023).



Although anyone can have a stroke, there are specific factors that raise your risk. It's critical to understand the risk factors and take steps to lower your risk.

Strokes can happen to anyone at any time, despite the common misconception that they exclusively affect the elderly. Understanding how to spot a stroke in either yourself or another else is vital. You can identify the indicators with the help of the FAST test.

Although they may occasionally be younger, including children, stroke victims tend to be older. In the UK, one in four people of working age experience a stroke. You can take a variety of actions to lower your risk. Your doctor can assist you in understanding your stroke risk through basic exams and in making the necessary changes to lower that risk. (Stroke Association, n.d).

Stroke warning signs

To recognise the indicators, act quickly: Facial asymmetry: Are they able to smile? Have their eyes or mouth drooped? Can they raise both arms given their arm weakness? Speech issues: Can they grasp what you're saying and speak clearly? If any of these symptoms appear, it's time to dial 999 immediately. Because timing is so important if you're having a stroke, it's named F.A.S.T. Without treatment, you could lose millions of nerve cells per minute. The possibility that your voice, movement, and powers will return to normal decreases the longer you wait. Acting quickly can really save lives. (British Heart Foundation, 2019).

Health and Lifestyle routine measures

According to (British Heart Foundation, 2012) Lifestyle Questionnaire is made to make it possible for you to evaluate how nutritious your diet is. After selecting yes or no to the questions below, read the supplemental material to help you decide whether to adjust your diet.

- Does not eat the recommended fruits and vegetables per day or weeks.
- Prefers to consume fat-containing foods.
- Eats high contents of Starchy foods.
- Likes to drink high contents of sugary.
- Likes to add salt in the food.
- Drinks little amount of water and drinks plenty of alcohol over the recommended units per week.

Health and Social Care practice

With the help of the Stroke Association and NHS England, the National Stroke Programme was created after receiving substantial input from patients and medical professionals. The Long-Term Plans targets for stroke will be more easily attained with the help of community organisations, and 85,000 more people will receive better prevention, treatment, and care thanks to the plan. It will make sure that the foundation for transformation is set in every region of the nation. The program's goals are to:

- Improve the programmes for post-hospital rehabilitation for stroke victims.
- Bring about a ten-fold rise in the percentage of patients who get a thrombectomy to end their stroke, resulting in 1,600 more persons being independent every year.
- Teach other consultants at hospitals how to do thrombectomy more places in order to provide a national service.
- Provide treatment to twice as many patients by 2025 to ensure that 20% of stroke patients receive clot-busting thrombolysis, the greatest performance in Europe.
- Boost progress along the stroke route, including in rehabilitation, by enhancing the Sentinel Stroke National Audit Programme (SSNAP), which will help identify new requirements.
- Ensure that 90 percent of patients—rather than the present 29%—receive six-month assessments of their needs and advancement.

We operate in conjunction with the Stroke Association, NHS Right Care, getting it Right First Time (GIRFT), many professional bodies, other arm's length bodies (ALBs), and non-profit groups to carry out this nationwide programme of work. (NHS England, n.d).

In October 2016, the National Clinical Guidelines for Stroke were updated to their fifth version. Under the leadership of Professor Tony Rudd, it was created by the Intercollegiate Stroke Working Party of the Royal College of Physicians. The National Audit Programme for Stroke is under the control of this working group as well. The National Institute for Health and Care Excellence has accredited the 2016 revision of the recommendation. (NICE, 2019).

The recommendation gives a thorough review of stroke care, including every step of the stroke pathway from immediate assistance to long-term rehabilitation, including secondary prevention. In order to improve the quality of care for all stroke patients, regardless of age, gender, stroke type, or location, the edition also provides healthcare professionals with information about what should be provided to stroke patients and how this should be organised. (Royal College of Physicians, 2016).

Recommendations

You can prevent stroke by making healthy decisions and controlling any current medical conditions.

Many strokes may be prevented by leading a maintaining a healthy lifestyle and treating any diseases that increase your risk for stroke in conjunction with your medical team. You can lower your risk of stroke by making a healthy lifestyle choice.

By making wholesome food choices for your meals and snacks, you can lower your risk of stroke. Make sure you eat a lot of fruits and vegetables that are fresh. Consuming meals low in cholesterol, trans fats, and saturated fats can help lower cholesterol. Limiting sodium and salt intake can lower blood pressure. Your chance of having a stroke is increased by both high blood pressure and excessive cholesterol.

Your risk of stroke increases if you are obese or overweight. Physical activity has several advantages, including helping you keep a healthy weight and reducing your cholesterol and blood pressure. You run a far greater chance of suffering a stroke if you smoke. If you're a smoker, stop right away. If you smoke, quitting will lower your chance of experiencing a stroke. Alcohol can raise blood pressure, so try to limit your consumption. Women should limit their alcohol consumption to one drink per day, while males should limit their intake to two drinks per day.

The best measures to lower your risk of stroke should be discussed with your doctor. Having diabetes or high blood pressure, high cholesterol, or any of these problems, there are steps you may take to reduce your risk of stroke. Your physician should check your cholesterol at least once every five years. Your medical team must be aware of this simple blood test. You can reduce your risk of stroke if you have high cholesterol by taking medication and making lifestyle changes. Because high blood pressure frequently has no symptoms, check your blood pressure periodically. If your doctor suspects you have any of the symptoms, they can suggest getting tested for diabetes. If you have diabetes, regularly check your blood sugar levels. (CDC, 2022).

Your risk is probably increased if a close family members who have experienced a stroke. Strokes affect persons of colour and those from South Asian families more frequently. Any risk factors you may have, such as being overweight, you may need to get evaluated for diabetes early if you are black or South Asian. Certain risk factors are more prevalent in women, such as pregnancy and use of the combination pill. If you must take a blood thinner, this may occasionally result in extremely heavy periods. The genetic ailment sickle cell disease damages red blood cells, which carry oxygen throughout the body. It may cause painful episodes and other symptoms in addition to raising the risk of having a stroke. Stroke risk is somewhat increased if you experience a migraine with aura, even though it has not been proven that migraines cause stroke. (Stroke Association, n.d).

Conclusion

If you've already had a stroke, your chance of having another is significantly higher. To reduce your stroke risk factors, you typically need long-term pharmaceutical therapy.

For instance, medication to control high blood pressure, antiplatelet or anticoagulant medications to lower the risk of blood clots, and statins to lower cholesterol levels. In order to enhance your overall health and reduce your risk of stroke, you will also be urged to adjust your lifestyle. Examples of such changes include eating a nutritious diet, exercising frequently, quitting smoking if you smoke, and consuming alcohol within the advised limits.

References

1. British Heart Foundation, (2019). *Signs of a stroke*. Available at: <https://www.bhf.org.uk/information-support/conditions/stroke> (Accessed: November 2019).
2. British Heart Foundation, (2012). How healthy is your diet? Questionnaire. Available at: https://moodle.uwtsd.ac.uk/pluginfile.php/666645/mod_resource/content/2/BHF_how-healthy-is-your-diet-questionnaire.pdf?redirect=1 (Accessed: 2012).
3. Mayfield Brain & Spine, (2023). *Blood supply of the brain*. Available at: [https://mayfieldclinic.com/pe-stroke.htm#:~:text=A%20stroke%20is%20a%20sudden,red\)%20circulation%20of%20the%20brain](https://mayfieldclinic.com/pe-stroke.htm#:~:text=A%20stroke%20is%20a%20sudden,red)%20circulation%20of%20the%20brain). (Accessed: 2023).
4. National Centre for Chronic Disease Prevention and Health Promotion, Division for Heart Disease and Stroke Prevention, (2022). Prevent Stroke: What You Can Do. Available at: <https://www.cdc.gov/stroke/prevention.htm> (Accessed: April 5th 2022).
5. NHS, (2022). *Stroke*. Available at: <https://www.nhs.uk/conditions/stroke/> (Accessed: 13th September 2022).
6. NHS England, (n.d). *NHS England's work on stroke*: Available at: <https://www.england.nhs.uk/ourwork/clinical-policy/stroke/> (Accessed: n.d).
7. Royal College of Physicians, (2016). *Stroke guidelines*. Available at: <http://www.rcplondon.ac.uk/guidelines-policy/stroke-guidelines> (Accessed: 3rd October 2016).
8. Stroke Association, (n.d). *Are you at risk of stroke?* Available at: https://www.stroke.org.uk/what-is-stroke/are-you-at-risk-of-stroke?gclid=CjwKCAjwue6hBhBVEiwA9YTx8lehMMapHxI6r5qAUXT2HPKBgmlRtFLsnRNRK7RXuNLQSKpHciETRoCOAYQAvD_BwE (Accessed: n.d).
9. Stroke Association, (n.d). *Main risk factors for stroke*. Available at: <https://www.stroke.org.uk/what-is-stroke/are-you-at-risk-of-stroke#:~:text=Lifestyle%20factors,late%20to%20make%20a%20change> (Accessed: n.d).