

ACDB4011 CHEMISTRY QUIZ 3

12

NAME

Kahe Byrne

DATE: Nov 08, 2022

1. (6 points) Indicate whether each of the following statements is true or false:

T F

☒ ☐

Toxic gases are liberated when the oxygen balance for a given explosive substance is large and negative.

☐ ☒

Supercritical fluid is a highly compressed fluid that combines the properties of gases and liquids.

☐ ☒

Dipole-dipole forces are stronger than network covalent bonds

☐ ☒

0.5 mol of nitrogen gas, N_2 , will occupy less space than 0.5 mol of methane gas, CH_4 , at the same temperature and pressure.

☒ ☐



According to Boyle's law, the volume of a gas is directly proportional to the temperature.

☐ ☒

Heavier gas molecules have higher kinetic energy than lighter gases at the same temperature and pressure.

2. (4 points) Explain in terms of intermolecular forces why does ice float in water? Support your answer with a drawing.

Water



H-bonds !!!
more in ice
→ keep the H₂O molecules closer

The force between bonds in the water are a lot more reduced than the ice intermolecular forces. Because of the tightly packed molecules ice would be lighter than water therefore floating to the top.

6

3. a) (4 points) Op How many moles of oxygen gas will react with 0.5 mol of $C_4H_{10}O$ in incomplete combustion to carbon monoxide and water?



0.5 mol of $C_4H_{10}O = 2 \text{ moles}$
of Oxygen gas

- b) (6 points) $2p$ Calculate the volume occupied by the moles of oxygen gas calculated in a) at a pressure 220.0 kPa and a temperature of 21.0 deg C.

pressure = 220.0 kPa Temperature = 21°C 2 moles of oxygen

$$PV = nRT$$

$$V = \frac{nRT}{P}$$

$$V = \frac{2 \times 0.0821 \times 21}{220}$$

$$V = 0.01567 \times 100$$

$$V = 1.567$$

- c) (6 points) Op What will be the volume occupied by the moles of oxygen gas calculated in a) if the pressure decreased to 100.0 kPa and temperature increased to 50.0 deg C. DO NOT use the ideal gas equation!

~~Use the ideal gas equation~~


$$V = \frac{nRT}{P}$$

$$= \frac{2 \times 0.0821 \times 50}{100}$$

$$= 0.0821 \times 100$$

$$= 8.21$$


4. a) (4 points) *Op* For a high explosive substance HMX ($C_4H_8N_8O_8$, 296.5 g/mol) calculate the oxygen balance.



$= \frac{16}{296.5}$

$4 \times 12 + 8 \times 1 + 8 \times 14 + 8 \times 16 = 296.5 \text{ g/mol}$
 $= 0.05396 \times 100$
 $= 5.396$

- b) (4 points) *Op* Using Kistiakowsky-Wilson rules determine the decomposition products of HMX.



- c) (2 points) *Op* What is the volume of all gaseous products formed upon decomposition of 1 mol of HMX at STP?

$\frac{5.396}{296.5} \times 1 = 0.0182 \times 100$
 $= 1.82$