



Component 1: Written Assignment (Case Study Report)

Student Name: Liliana Vladut

Student ID 2411574

Module Name: BSC- Health and Social Care

Module Code: BMHP 5023

Lecturer: Soban Sadiq

Table of Contents

Introduction	2
Regulation of Blood Glucose Levels	2
Cellular Metabolism: The Role of Oxygen and Glucose in Energy Production	3
Importance of Recording and Tracking Blood Glucose Levels	4
Evidence-Based Lifestyle Recommendations	5
Conclusion	6
References	8

Introduction

Diabetes mellitus is a long-lasting metabolic condition that is caused by chronic hyperglycaemia caused by a lack of insulin secrets, insulin effects, or both. The most common type is Type 2, which is closely linked with the modifiable lifestyle conditions including obesity, lack of physical activity and poor eating habits (Ahmad et al., 2026). The increase in the number of diagnoses in the primary care environment underscores the actual importance of patient education and disease management plans. In healthy physiological states, the blood glucose level is tightly controlled within a high and low range via the action of several hormones.

Insulin helps in the absorption of glucose in the body cells to generate energy whereas glucagon helps to maintain the blood glucose levels normal. In Type 2 diabetes, there is the development of insulin resistance which decreases the responsiveness of the body to insulin. The functioning of pancreatic beta cells can decrease with time and result in inadequate insulin synthesis and chronic hyperglycaemic state (Alserhani et al., 2025). A prolonged TG condition leads to the development of major complications such as cardiovascular disease, neuropathy, nephropathy and retinopathy. Some of the main risk factors are an aging population, genetic predisposition, central obesity, sedentary lifestyle, cigarette and socioeconomic factors. These factors need to be understood as a way of preventing and early identification and treatment of the condition.

Regulation of Blood Glucose Levels

Blood glucose level is an important aspect that should be controlled to maintain homeostasis. The endocrine role of the pancreas is the major regulator of this process. The pancreas has groups of specialised cells referred to as the Islets of Langerhans. Beta cells and alpha cells can be found in these islands producing insulin and glucagon respectively (Barchiesi et al., 2025). These hormones coexist in a counter-regulatory relationship to ensure that the level of blood glucose is within a safe physiological range. When the blood glucose level increases, the insulin is released especially after consuming carbohydrates. It attaches to the insulin receptors of the target tissues like

skeletal tissue and fat. This communication activates the removal of glucose transporters to the cell membrane that enables glucose to be introduced into the cell. When glucose gets inside it can be utilized in immediate production of energy or it can be stored in glycogen in the liver and in the muscles through the process of glycogenesis. Insulin is also able to prevent gluconeogenesis, which decreases the synthesis of new glucose in the liver.

Glucagon is a counter regulating hormone. Glucagon is secreted in response to a reduction in blood glucose such as during fasting or strenuous exercise by alpha cells. It triggers glycogenolysis or the breakdown of glycogen into glucose and gluconeogenesis. These activities raise the amount of glucose in the blood, and this keeps the body constantly active with energy to essential organs, especially the brain. This control system works by way of negative feedback (Cao et al., 2024). Increase in glucose levels causes an increase in insulin secretion to reduce glucose levels. When the glucose levels drop glucagon secretion rises to elevate glucose levels. This is a finely balanced system that is disturbed in Type 2 diabetes. Insulin resistance refers to a situation where the cells are not responsive to insulin which causes low uptake of glucose. This results in glucose being left in blood hence hyperglycaemia. In the long run, the long-time reliance on the beta cells can lead to insulin production and deteriorate glycaemic control. The connection between the physiological functioning and the anatomical structure is obvious. The direct effect of structural changes in physiological processes is the dysfunction of pancreatic beta cells, which directly affects hormonal regulation.

Cellular Metabolism: The Role of Oxygen and Glucose in Energy Production

Most cells in the human body use glucose as their main energy substrate and the energy source is vital in ensuring normal physiological functioning. Cellular respiration is the biochemical process in which glucose is transformed into the primary energy currency in cells namely adenosine triphosphate (ATP). This process takes place through the involvement of oxygen and mostly happens in specialised organelles called mitochondria (Dwibedi et al., 2022). The precise generation of energy is the basis of maintaining cellular processes and homeostasis in general. The cellular respiration follows a series of three

steps. The initial phosphorylation occurs in the cytoplasm, which is referred to as the glycolysis stage. In the glycolysis process, a molecule of glucose is enzymatically cleaved to produce two molecules of pyruvate with a minor production of ATP and reduced coenzymes. This step is independent of oxygen and can take place in the anaerobic state. Nevertheless, to produce energy at its highest levels, oxygen is required to take place in the following stages.

The Krebs cycle is the second stage that is situated in the mitochondrial matrix. In this case, pyruvate is oxidized to the high-energy carriers of electrons that are stored as potential energy. The last step is the electron transport chain which occurs along the inner membrane of the mitochondrion. The use of oxygen as the last electron acceptor allows a much larger number of ATP to be produced using oxidative phosphorylation. ATP produced by this mechanism plays a role in the essential physiological processes such as muscle contraction, nerve impulse conduction, active membrane transport, protein synthesis and cell repair (Hakeem et al., 2024). Insulin resistance in patients with Type 2 diabetes affects the process of glucose uptake into the cells. Although the blood glucose level is high, there is a decrease in intracellular glucose supply leading to compromised efficiency in ATP production. The result of this metabolic imbalance may be fatigue and low physical performance. Also, persistent hyperglycaemia facilitates oxidative stress and vascular damage, which predisposes to the occurrence of the long-term complications. It is thus important to have the ability to effectively regulate glucose to achieve energy production as well as the prevention of progressive tissue injury.

Importance of Recording and Tracking Blood Glucose Levels

Regular data monitoring is one of the key components of the efficient management of Type 2 diabetes. The measurement of blood glucose helps patients and medical workers to assess the success of treatment methods and determine trends in glycaemic regulation. Regular monitoring is objective evidence that helps in the process of clinical decision-making and encourages patient involvement in self-care (Parker, 2020). Blood glucose self-monitoring is usually done through a glucometer which measures blood samples collected by pricking fingers into the capillary blood. The approach can give real-

time data on the current glucose levels and enables patients to react to the changes accordingly. CGM devices provide a more in-depth information as they monitor the changes in glucose levels during the day and night and reveal the patterns that might not be revealed during intermittent readings. Moreover, glycated haemoglobin (HbA1c) test indicates average blood glucose level during two to three months, which gives a vital indicator of the long-term metabolic control.

Monitoring and measuring of blood glucose helps patients to understand how lifestyle interventions like diet, physical activity, stress and drug adherence affect their state. Determining constant trends assists in the prevention of acute complications such as hypoglycaemia and the extreme hyperglycaemia. In the case of healthcare workers, it is possible to use this routine information and adjust medications, conduct individual care planning and identify patients who are more exposed to developing long-term complications (Shah et al., 2021). Routine monitoring has drawbacks although it has its advantages. Others can be anxious and uncomfortable or have a financial constraint that decreases the adherence. Unreliable readings may also be caused by inaccurate technique, and this may influence the treatment decisions. Thus, formal education and continuous professional care are necessary to achieve proper data interpretation and safe self-care. The methodical application of regular health data shows very well how the knowledge of physiology is applied in practice in clinical practice.

Evidence-Based Lifestyle Recommendations

Lifestyle change is a principle of Type 2 diabetes care, and it is well acknowledged as a key element of the long-term glycaemic management. The studies have shown that specific dietary, physical activity, smoking behaviour and weight management changes have a strong positive impact on insulin sensitisation and the risk of complications. In comparison, lifestyle interventions can be described as solutions to insulin resistance that do not rely on pharmacological intervention but the underlying metabolic dysfunction (Ahmad et al., 2026). Change of diet plays a central role in controlling the level of glucose in the blood. Whole grain, vegetarian, lean protein, and non saturated fat diets lead to slower glucose absorption rates as well as postprandial peaks. By avoiding high

glycaemic index and refined carbohydrates, one will avoid the rapid increase in blood glucose levels, so the pancreatic beta cells will not have to work hard. High dietary fibre slows down gastric emptying and enhances the glycaemic stability. Moreover, decreasing the amount of excess calories helps to maintain weight loss lowering adipose tissue-related inflammation and increasing the insulin receptor sensitivity at the cellular level.

Exercise is also very imperative to the physiology. Glucose uptake does not require insulin, and is stimulated, through alternative metabolic pathways, by skeletal muscle contraction during exercise. Frequent aerobic and resistance exercises lead to greater efficiency of mitochondria, greater retention of muscle glycogen and overall elevated insulin sensitivity. Physical activity also decreases blood pressure and improves lipid profiles in addition to the glycaemic gains, which reduces cardiovascular risk that is high in diabetes. Cessation of smoking is also relevant (Alserhani et al., 2025). Nicotine exposure helps in endothelial dysfunction as well as insulin resistance, which accelerates the process of vascular damage. Cessation of smoking enhances vascularization, lowers the levels of oxidative stress and diminishes the risk of macrovascular complications. Weight control is closely associated with better metabolic consequences. Small changes in weight cause significant changes in HbA1c and insulin sensitivity. Nevertheless, obstacles like socioeconomic obstacles, cultural eating habits and mental defence can interfere with compliance. Long term behavioural change, therefore, requires personalised care planning, education and long-term professional support.

Conclusion

Diabetes type 2 is a progressive and complicated metabolic illness that is characterised by a lack of insulin regulation caused by the resistance to insulin and dysfunction of the pancreatic beta cells. In usual physiological state, the glucose levels of the blood are well maintained by a series of regulated hormonal processes that include insulin and glucagon that make sure that glucose is used in the cells to produce energy, and that it does not build up in the body. The presence of glucose along with oxygen is necessary in the formation of adenosine triphosphate (ATP) which aids all other physiological functions such as muscle contraction, nerve conduction and tissue repair.

Interference with these regulatory processes in Type 2 diabetes causes sustained hyperglycaemia that in the long run causes the risk of microvascular as well as macrovascular disease. Management would involve proper knowledge of the physiology and realistic measures employed to manage and regulate the condition hence effective management can only be achieved by having a complete insight into the physiology and the practical measures employed in the treatment of the condition. Routine blood glucose and HbA1c tracking allows identifying tendencies early, facilitates prompt clinical interventions and allows planning personal care.

Lifestyle change has been the focus of enhancing better metabolic results. It has been demonstrated that dietary modifications, physical exercise, smoking quit and weight control are very helpful in improving the insulin sensitivity and decreasing the risks in long term health conditions. Structured education, interpretation of routine health data and facilitation of sustainable behavioural change are also important activities as a practice nurse. Educating patients on information and practical skills empowers them to have better self-management, increase compliance with treatment and maintain safer and more effective long-term management of their condition.

References

- Ahmad, I., Taimur, H., Poduri, G.V., Nawaz, A., Shiriyama, Y., Shabbir, S., Rahman, M.S., Uzakova, A., Ahmad, H.S., Okamoto, M. and Yuasa, M., 2026. A Systematic Review and Meta-Analysis of RCTs Assessing Efficacy of Lifestyle Interventions on Glycemic Control in South Asian Adults with Type 2 Diabetes. *Medical Sciences*, 14(1), p.48.
- Alserhani, M.F., Alrasheed, F., Al Qamshah, A.M., Alkharashi, Y., Aljarallah, M.F., Alarifi, F.F., Althaqeb, E., Alrasheed, M.I., Alkharashi, L.M., Al Eidan, E. and Alrasheed, M., 2025. Effectiveness of Self-Management of Blood Glucose in Improving Glycemic Control in Patients with Diabetes: A Systematic Review and Meta-Analysis. *Cureus*, 17(12).
- Barchiesi, M.A., Calabrese, A., Costa, R., Di Pillo, F., D'Uffizi, A., Tiburzi, L. and Zahid, E., 2025. Continuous glucose monitoring in type 2 diabetes: a systematic review of barriers and opportunities for care improvement. *International Journal for Quality in Health Care*, 37(3), p.mzaf046.
- Cao, Y., Shrestha, A., Janiczak, A., Li, X., Lu, Y. and Haregu, T., 2024. Lifestyle intervention in reducing insulin resistance and preventing type 2 diabetes in Asia Pacific region: a systematic review and meta-analysis. *Current diabetes reports*, 24(9), pp.207-215.
- Dwibedi, C., Mellergård, E., Gyllensten, A.C., Nilsson, K., Axelsson, A.S., Bäckman, M., Sahlgren, M., Friend, S.H., Persson, S., Franzén, S. and Abrahamsson, B., 2022. Effect of self-managed lifestyle treatment on glycemic control in patients with type 2 diabetes. *NPJ digital medicine*, 5(1), p.60.
- Hakeem, L., Ahmad, H., Aslam, A., Nawaz, K., Ali, U., Shaifq, H.R., Iqbal, N. and Ullah, M.K., 2024. A Comprehensive Review on Insulin Resistance. *International Journal of Immunological Nursing*, 10(1), pp.1-4p.
- Parker, J., 2020. Glucose metabolism, energy production and regulation of cellular and whole-body metabolism. *Journal of the Australasian College of Nutritional and Environmental Medicine*, 39(1), pp.29-33.
- Shah, S.Z., Karam, J.A., Zeb, A., Ullah, R., Shah, A., Haq, I.U., Ali, I., Darain, H. and Chen, H., 2021. Movement is improvement: the therapeutic effects of exercise and general physical activity on glycemic control in patients with type 2 diabetes mellitus: a systematic review and meta-analysis of randomized controlled trials. *Diabetes Therapy*, 12(3), pp.707-732.