

AI Acknowledgement Table

URL	Date Used	Prompts
https://app.grammarly.com/	30/03/26	Check my gramar
copilot.microsoft.com	28/03/26	Provide research on the different fitness tests

Dylan Arnold - 2501635

Word Count - 1536

Comprehensive Health and Fitness Assessment Report

Introduction

This report looks at the health and fitness of a 19 year old male who is 6ft2 and weighs 92kg. He plays football now and again but doesn't train much, so he is classed as moderately active. This means he does some physical activity, but not enough to gain the full health benefits that come from regular training.

Health-related fitness includes cardiovascular endurance, muscular strength, muscular endurance, flexibility, and body composition. These are important because they affect how well someone can move, exercise, and stay healthy. Low fitness levels are linked to problems such as heart disease, obesity, and type 2 diabetes (American College of Sports Medicine, 2021). These conditions can develop over time, so improving fitness early in life can help reduce long-term health risks.

Fitness testing helps identify what someone is good at and what they need to improve. It also gives a starting point to measure progress later on and helps when planning training programmes. Without testing, it would be difficult to know whether training is working or if changes need to be made.

The aim of this report is to carry out a range of fitness tests, compare the results to normal values, and give recommendations to help improve overall health and fitness.

Fitness Test Design

A set of tests was chosen to assess each part of health-related fitness. These tests are mostly used in schools, gyms, and sports settings, so they are familiar and easy to carry out.

The Queen's College Step Test was used to assess cardiovascular endurance. It measures heart rate after stepping for three minutes. It is simple, doesn't take long, and works well for people with average fitness levels (Heyward, 2018). It is also safer than maximal tests, which require someone to work at full effort.

Muscular strength was measured using a handgrip dynamometer. This test is easy to use and gives a good idea of overall strength (Baechle and Earle, 2016). Grip strength is often used because it links well to general strength and is quick to measure.

Muscular endurance was tested using the one minute sit-up test. This shows how many sit-ups can be completed in one minute and gives an idea of how long the abdominal muscles can keep working. Muscular endurance is important for posture, daily tasks, and performance.

Flexibility was measured using the sit and reach test. This looks at flexibility in the hamstrings and lower back, which are important for posture and movement (Heyward, 2018). Poor flexibility can lead to tight muscles and a higher risk of injury.

Body composition was assessed using waist measurement. This shows how much fat is stored around the stomach area, which is linked to health risks such as heart disease (NHS, 2023). Waist measuring is often used because it is quick and does not require specialist equipment.

These tests were chosen because they are simple, don't need expensive equipment, and are often used in fitness testing. They can also be repeated easily, which is useful for tracking progress.

There are some limitations. The Queen's Step Test only gives an estimate of cardiovascular endurance. The sit up test may use other muscles, which can affect accuracy. Waist measurement does not show body fat percentage and cannot tell the difference between fat and muscle. Even with these limitations, the tests are suitable for this assessment.

The Queen's Step Test looks at how well the heart and lungs work together. The sit-up test shows how long the abdominal muscles can work without stopping. Grip strength shows overall muscle strength. The sit and reach test shows how flexible the muscles and joints are. Waist measurement shows how fat is stored around the body.

These tests are realistic for everyday fitness settings because they are safe, easy to repeat, and can be done in places like gyms, sports halls, or schools. They are also suitable for people with different fitness levels, which makes them practical for general assessments.

Procedures

Before testing, the client completed a PAR-Q to make sure it was safe to take part. This is important because it checks for any medical issues that might make exercise unsafe. All tests were carried out using standard procedures to keep results fair and consistent (American College of Sports Medicine, 2021).

Waist measurement was taken using a flexible tape measure. The tape was placed around the stomach at the top of the hip bone while the client stood upright. The measurement was taken after a normal breath out. This helps avoid errors caused by breathing in or holding the stomach in.

For the Queen's Step Test, the client stepped up and down on the step platform for three minutes at a set pace using a metronome. A heart-rate monitor recorded the heart rate before and after the test. The pace and height of the step were kept the same throughout to make sure the test was fair.

For the sit up test, the client lay on their back with knees bent and feet held in place. Sit-ups were completed for one minute, only correct reps were counted. This helps make sure the results are accurate and not affected by poor technique.

Grip strength was measured using a handgrip dynamometer. The client squeezed as hard as possible for three attempts, and the average score was recorded. Taking the average helps stop one really good or really bad attempt from affecting the result too much.

The sit and reach test was done with the client sitting with legs straight and shoes removed. The client reached forward as far as possible, and the best score was recorded. The test was repeated to make sure the result was reliable.

Standardised procedures are important because they make results reliable and allow fair comparison with normal values. Validity is also important, as it shows whether a test measures what it is meant to measure. For example, the Queen's Step Test only estimates cardiovascular endurance. Reliability means getting similar results each time, which can be improved by using the same equipment and instructions.

Results

Results Compared to Normative Data Interpretation

Test	Result	Comparison	Meaning
Waist measurement	105 cm	Above 102 cm	High risk
Queen's Step Test	158 bpm	Average	Average fitness
Sit-ups	39 reps	Above average	Good
Sit and reach	23 cm	Average to above average	Good
Grip strength	42.2 kg	Low-average to average	Moderate

The client's waist measurement of 105cm is above the healthy level for males, which is below 94cm. Anything above 102 cm is classed as high risk (NHS, 2023). This suggests the client carries too much fat around the stomach area, which increases the risk of health problems. Reducing waist size would help lower these risks.

The Queen's Step Test result of 158 bpm shows an average level of cardiovascular endurance. This matches the client's low training levels. Someone who trains regularly would normally have a lower heart rate after the test because their heart and lungs would be more efficient.

The client completed 39 sit ups, which is above average for this age group (Heyward, 2018). This shows good muscular endurance and suggests the client has reasonably strong core muscles.

The sit and reach score of 23cm is within the average to above-average range, showing a good level of flexibility. Good flexibility can help reduce injury risk and improve movement in sport and daily life.

Grip strength measured at 42.2 kg falls within the low-average to average range, showing moderate muscular strength. Improving strength would help with daily tasks and overall fitness.

Overall, the client has good muscular endurance and flexibility but needs to improve cardiovascular endurance and body composition. These weaknesses could affect long-term health if not improved. Playing football occasionally may explain the good endurance and flexibility. The lack of regular training likely contributes to the average cardiovascular result and high waist measurement.

Recommendations

To improve overall fitness and reduce health risks, the client should increase their physical activity levels. Small changes, such as walking more or being active daily, can also help.

For cardiovascular endurance, the client should aim for at least 150 minutes of moderate exercise per week, such as running, cycling, or playing football more often (World Health Organisation, 2020; NHS, 2023). This can be broken into shorter sessions to make it easier to manage.

To reduce waist measurement, a mix of cardiovascular exercise and strength training is recommended. Strength training should be done two to three times per week and should target major muscle groups (Baechle and Earle, 2016). This will help increase muscle mass and improve metabolism.

To improve muscular strength, the client should include resistance exercises such as weight training or bodyweight exercises. These can be progressed over time by increasing weight or reps.

To maintain and improve muscular endurance, exercises like sit ups, planks, and circuit training should be done regularly. These exercises help the muscles work for longer without getting tired.

Flexibility can be maintained through regular stretching, especially after exercise. Stretching helps reduce muscle tightness and can improve posture.

A balanced diet and staying active throughout the day will also help improve health and fitness. Eating a mix of fruits, vegetables, lean protein, and whole grains can support training and help reduce body fat.

References

American College of Sports Medicine (2021) ACSM's Guidelines for Exercise Testing and Prescription. 11th ed. Philadelphia: Wolters Kluwer.

Baechle, T.R. and Earle, R.W. (2016) Essentials of Strength Training and Conditioning. 4th ed. Champaign, IL: Human Kinetics.

Heyward, V.H. (2018). Advanced Fitness Assessment and Exercise Prescription. 7th ed. Champaign, IL: Human Kinetics.

McArdle, W.D., Katch, F.I. and Katch, V.L. (2022). Exercise Physiology: Nutrition, Energy, and Human Performance. 9th ed. Philadelphia: Wolters Kluwer.

NHS (2023) Physical activity guidelines. Available at:
<https://www.nhs.uk/live-well/exercise/> (Accessed: 27 March 2026).

World Health Organisation (2020) WHO guidelines on physical activity and sedentary behaviour. Available at: <https://www.who.int/publications/i/item/9789240015128> (Accessed: 327 March 2026).