

Table 1*Means and SD for hyperactivity scores in the chocolate and no chocolate conditions*

<i>Condition</i>	<i>Mean</i>	<i>SD</i>
Chocolate	16	3.5
No chocolate	11	4.5

Table 2*AAS users and non-users across the 10 subscales of the MDI (M, SD and t value)*

<i>Group</i>	<i>Users M (SD)</i>	<i>Non-users M (SD)</i>	<i>t</i>
Body anxiety	16.10 (4.98)	9.54 (3.90)	7.329*
Compulsivity	8.34 (4.06)	10.28 (4.87)	2.159*
Illusory eating	3.76 (1.66)	3.54 (1.12)	.775
Inadequacy	13.34 (5.46)	8.32 (3.66)	5.399*
Inappropriate eating	3.74 (1.30)	3.54 (1.40)	.738
Increased muscularity	11.10 (4.34)	9.14 (3.55)	2.470*
Muscularity drive	13.02 (4.74)	8.78 (3.63)	5.015*
Persistence	10.82 (3.95)	8.12 (3.74)	3.506*
Preoccupation	13.40 (3.66)	8.62 (2.79)	7.337*
Social sacrifice	6.66 (2.93)	4.70 (1.05)	4.439*

* $p < .05$