



Prifysgol Cymru
Y Drindod Dewi Sant
University of Wales
Trinity Saint David

Component 1: Written Assignment (Case Study Report)

Student name: Olga Maculska-Migovska

Student ID number: 2242936

Programme: BSc Health and Social Care

Module code: BMHP5023

Module title: Introduction to Physiology

Lecturer: Shaila Hussain

Case Study 2: Asthma

Submission date: 9 April 2026

Case Study Report: Adolescent Asthma Management

Introduction

Asthma is a chronic inflammatory condition of the airways that affects normal breathing. It is characterised by variable airflow obstruction and bronchial hyper-responsiveness and is commonly associated with symptoms such as wheezing, shortness of breath, chest tightness and coughing (NHS, 2024; World Health Organization, 2024).

This case study focuses on adolescents attending a local GP surgery asthma clinic, where more serious exacerbations have been linked to lifestyle-related triggers including smoking, vaping and spending time in crowded environments such as bars and clubs. These factors may increase exposure to chemical irritants and respiratory infections, which can worsen airway inflammation and reduce asthma control (Global Initiative for Asthma, 2024).

A clearer understanding of respiratory structure, function and triggers may help individuals with asthma engage more effectively with their care plans and manage their condition more safely. This report examines the structure of the respiratory system, the processes of mechanical respiration and gas exchange, the impact of irritants on respiratory function, the importance of peak-flow monitoring, and evidence-based lifestyle recommendations to support better asthma management.

Anatomy and Physiology of the Respiratory System

Structure of the Respiratory System and Mechanical Respiration

The lower respiratory tract begins at the trachea, a tube that carries air from the upper airways to the lungs. The trachea divides into the right and left bronchi, with one bronchus entering each lung. Inside the lungs these bronchi continue to branch into smaller airways known as bronchioles. The bronchioles terminate in tiny air sacs called alveoli, which are the primary site of gas exchange (Tortora and Derrickson, 2023).

The bronchial tree forms a complex branching network that conducts air from the trachea to the alveoli. As described by Tortora and Derrickson (2023), the epithelial lining of the airways gradually changes along this pathway. In the trachea and bronchi, the epithelium is ciliated pseudostratified columnar epithelium containing mucus-secreting goblet cells. As the airways divide into smaller bronchioles the epithelium becomes simpler and transitions to ciliated simple cuboidal epithelium. At the same time the amount of cartilage decreases while the proportion of smooth muscle increases. This structural change is important in asthma because the smooth muscle surrounding the bronchioles becomes highly reactive to irritants such as cigarette smoke or e-cigarette vapour, which can trigger bronchoconstriction and airway narrowing (Tortora and Derrickson, 2023).

Mechanical respiration, also known as pulmonary ventilation, refers to the movement of air into and out of the lungs. During inspiration the diaphragm contracts and flattens while the external intercostal muscles lift the rib cage upwards and outwards. These actions increase the volume of the thoracic cavity. As thoracic volume increases the pressure inside the lungs falls below atmospheric pressure, allowing air to flow into the lungs.

During expiration the diaphragm and external intercostal muscles relax. The diaphragm returns to its dome-shaped position and the rib cage falls. This reduces thoracic volume and

increases pressure inside the lungs, forcing air out. Under normal conditions expiration is largely a passive process (Tortora and Derrickson, 2023).

In asthma the airways become narrowed due to inflammation, excess mucus production and contraction of the smooth muscle within the airway walls. This increases resistance to airflow, making it harder for air to move in and out of the lungs. As a result, breathing becomes more difficult and symptoms such as wheezing, coughing and shortness of breath may occur (Global Initiative for Asthma, 2024).

This increased airway resistance means that greater effort is required during breathing, particularly during expiration when air may become partially trapped in the lungs. Air trapping reduces effective ventilation and contributes to breathlessness. Exposure to environmental triggers such as cigarette smoke, e-cigarette vapour, air pollution and respiratory infections can worsen airway inflammation and increase the risk of asthma exacerbations (Global Initiative for Asthma, 2024).

Gas Exchange in the Lungs

Gas exchange occurs in the alveoli, which are surrounded by a network of pulmonary capillaries. The alveolar-capillary membrane is extremely thin, allowing gases to diffuse easily across it. Oxygen from inhaled air passes from the alveoli into the blood where it binds to haemoglobin in red blood cells and is transported to body tissues. At the same time carbon dioxide, a waste product of cellular metabolism, diffuses from the blood into the alveoli so that it can be removed during exhalation (Tortora and Derrickson, 2023).

In asthma narrowed airways may reduce effective ventilation and make gas exchange less efficient during exacerbations. Effective gas exchange depends on pressure gradients between gases in the alveoli and the blood in pulmonary capillaries. Oxygen moves from areas of higher partial pressure in the alveoli to lower partial pressure in the blood, while carbon dioxide diffuses in the opposite direction. When airway obstruction occurs ventilation of some alveoli may be reduced, limiting oxygen diffusion and contributing to symptoms such as breathlessness and fatigue (Tortora and Derrickson, 2023).

Additional factors can also influence the efficiency of gas exchange. Effective gas exchange depends not only on diffusion across the alveolar-capillary membrane but also on adequate ventilation and blood flow within the lungs. This relationship is commonly referred to as ventilation-perfusion matching. When ventilation and blood flow are well balanced, oxygen uptake and carbon dioxide removal occur efficiently. However, in asthma, airway narrowing may reduce airflow to certain alveoli, creating ventilation-perfusion mismatch. As a result, some areas of the lung receive less oxygen, which may contribute to fatigue, breathlessness and reduced exercise tolerance (Tortora and Derrickson, 2023).

Pathophysiology: The Impact of Irritants

Smoking and vaping expose the respiratory system to chemical irritants that can inflame the airway lining. These irritants trigger airway inflammation, leading to swelling of the airway walls and narrowing of the airway lumen. In asthma this inflammatory response worsens airway hyper-responsiveness and increases sensitivity to environmental triggers (Global Initiative for Asthma, 2024; Sinyor and Perez, 2019).

Irritants can also stimulate contraction of the smooth muscle surrounding the bronchioles. This bronchoconstriction further narrows the airways and increases airflow resistance. Irritation of

the airway lining may also increase mucus production, which can partially block the airways and make breathing more difficult (Sinyor and Perez, 2019).

Cigarette smoke and e-cigarette vapour may also damage the normal protective mechanisms of the respiratory tract. Toxic substances in smoke and aerosol can impair mucociliary clearance, meaning that particles, allergens and pathogens remain in the airways for longer. In adolescents who spend time in crowded environments this may increase exposure to respiratory infections that can trigger asthma exacerbations (Global Initiative for Asthma, 2024; Right Decisions NHS Scotland, 2024).

During an asthma exacerbation airway inflammation, bronchoconstriction and mucus production combine to reduce effective ventilation. Expiration becomes particularly difficult because narrowed airways can trap air in the lungs. As a result, individuals may experience wheezing, coughing, chest tightness and shortness of breath, reflecting worsening airflow limitation and respiratory function (Global Initiative for Asthma, 2024; Sinyor and Perez, 2019).

Recording and Tracking Respiratory Function

Monitoring respiratory function is an important component of asthma management because it helps individuals identify early deterioration and respond appropriately. One commonly used method is peak expiratory flow (PEF) measurement, which records the maximum speed at which a person can exhale air from the lungs. This measurement is obtained using a handheld device called a peak flow meter and is widely recommended for individuals with asthma to assess airway function at home (Global Initiative for Asthma, 2024).

Regular monitoring of respiratory function also plays an educational role in asthma self-management. Adolescents who track their peak flow readings become more aware of how their breathing changes in response to triggers such as smoke exposure, respiratory infections or physical activity. Over time, recognising these patterns can help individuals anticipate worsening symptoms and respond earlier. Healthcare professionals may use peak flow records during asthma reviews to assess treatment effectiveness and determine whether medication adjustments are required. This collaborative approach encourages adolescents to participate actively in managing their condition and supports better long-term asthma control (Global Initiative for Asthma, 2024; NHS, 2024).

Peak flow readings indicate how well air is moving through the airways. When the airways become inflamed, narrowed or obstructed by mucus, airflow is reduced, and peak flow values may fall below the individual's normal range. These physiological changes occur because asthma causes chronic airway inflammation, bronchial hyper-responsiveness and mucus production that restrict airflow through the bronchi and bronchioles (Sinyor and Perez, 2019).

Regular monitoring allows individuals to identify patterns or sudden decreases in peak flow that may indicate the beginning of an asthma exacerbation. Early detection enables individuals to follow their asthma action plan, adjust medication if required and seek medical advice before symptoms become severe (Global Initiative for Asthma, 2024).

In addition to peak-flow measurement healthcare professionals may assess respiratory function using spirometry. This clinical test measures lung function and airflow, including forced expiratory volume (FEV1), which helps clinicians evaluate asthma control and guide treatment decisions (NHS, 2024; British Thoracic Society, National Institute for Health and Care Excellence and Scottish Intercollegiate Guidelines Network, 2024).

For adolescents attending asthma clinics regular monitoring can support better self-management. Tracking peak flow readings helps individuals recognise early warning signs and identify potential triggers such as smoking, vaping or respiratory infections (Global Initiative for Asthma, 2024).

Evidence-Based Lifestyle Recommendations

Lifestyle changes play an important role in improving asthma control and reducing the risk of exacerbations. In adolescents, self-management is particularly important because symptoms may worsen when medication is used inconsistently or when triggers such as smoking, vaping and respiratory infections are ignored (Global Initiative for Asthma, 2024).

One of the most important recommendations is to avoid smoking and vaping. Tobacco smoke and e-cigarette vapour irritate the airways and increase the risk of asthma symptoms and attacks (Centers for Disease Control and Prevention, 2024). Adolescent asthma guidance recommends asking young people whether they smoke or vape and supporting them to stop. Avoiding these irritants may improve asthma control and reduce the likelihood of severe exacerbations (Right Decisions NHS Scotland, 2024).

Another recommendation is to reduce exposure to crowded environments where respiratory infections may spread easily. Environments such as bars or clubs may increase exposure to respiratory viruses or poor air quality, both of which can trigger asthma symptoms. Practising good hand hygiene and recognising personal triggers can help reduce the risk of infection-related exacerbations (Global Initiative for Asthma, 2024).

Adherence to prescribed medication is also essential. Preventer medication taken consistently can reduce airway inflammation and stabilise airway function. Adolescents may rely too heavily on reliever inhalers instead of regular preventer therapy, which can lead to poorer asthma control. A personalised asthma action plan helps individuals recognise worsening symptoms and respond appropriately by adjusting treatment or seeking medical advice (NHS, 2024).

Regular physical activity may support overall health when asthma is well controlled. Exercise is not a replacement for medical treatment, but moderate activity can support general fitness and respiratory health. Education about asthma triggers, symptom recognition and correct inhaler technique can further improve self-management and support safer long-term asthma control.

Conclusion

Asthma is a chronic inflammatory condition that affects the airways and can significantly impair normal breathing. Understanding the structure and function of the respiratory system explains how airflow moves through the trachea, bronchi, bronchioles and alveoli. Mechanical respiration moves air into and out of the lungs, while gas exchange in the alveoli allows oxygen to enter the bloodstream and carbon dioxide to be removed from the body.

This report has shown that irritants such as smoking and vaping can increase airway inflammation, bronchoconstriction and mucus production, which restrict airflow and worsen respiratory symptoms. Monitoring respiratory function using tools such as peak-flow measurements enables early detection of airway deterioration.

Evidence-based lifestyle recommendations including avoiding smoking and vaping, reducing exposure to respiratory infections, adhering to preventer medication and following personalised asthma action plans can improve asthma management. Together these strategies help adolescents manage asthma more safely and maintain better long-term respiratory health.

Reference List

British Thoracic Society, National Institute for Health and Care Excellence and Scottish Intercollegiate Guidelines Network (2024) *Asthma: diagnosis, monitoring and chronic asthma management (NG245)*. Available at: <https://www.nice.org.uk/guidance/ng245> (Accessed: 8 April 2026).

Centers for Disease Control and Prevention (2024) *Asthma and smoking*. Available at: <https://www.cdc.gov/asthma/> (Accessed: 8 April 2026).

Global Initiative for Asthma (2024) *Asthma management and prevention for adults, adolescents and children 6–11 years: A summary guide for healthcare providers*. Available at: <https://ginasthma.org> (Accessed: 8 April 2026).

NHS (2024) *Asthma*. Available at: <https://www.nhs.uk/conditions/asthma/> (Accessed: 8 April 2026).

Right Decisions NHS Scotland (2024) *Asthma in adolescents*. Available at: <https://rightdecisions.scot.nhs.uk/asthma-pathway-bts-nice-sign-sign-244/asthma-in-adolescents/> (Accessed: 8 April 2026).

Right Decisions NHS Scotland (2024) *Asthma pathway (BTS, NICE, SIGN) [SIGN 244]*. Available at: <https://www.rightdecisions.scot.nhs.uk/asthma-pathway-bts-nice-sign-sign-244/> (Accessed: 8 April 2026).

Sinyor, B. and Perez, L.C. (2023) *Pathophysiology of asthma*. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK551579/> (Accessed: 8 April 2026).

Tortora, G.J. and Derrickson, B. (2023) *Principles of Anatomy and Physiology*. 17th edn. Hoboken: Wiley.

World Health Organization (2024) *Asthma*. Available at: <https://www.who.int/news-room/fact-sheets/detail/asthma> (Accessed: 8 April 2026).

National Institute for Health and Care Excellence (2024) *Asthma: diagnosis, monitoring and chronic asthma management*. Available at: <https://www.nice.org.uk/guidance/ng245> (Accessed: 8 April 2026).